tension. If partial relapse should occur—and I have never known more than this—any bands that may have escaped division, or any which may since have become prominent by the extension, may be

Fig. 8.—Hand showing Dupuytren's Contraction of Middle, Ring, and Little Fingers. From a cast.

Fig. 9.—The same Hand as shown in fig. 7, thirteen years after operation. The Middle and Ring Fingers remained completely cured, with full power of Flexion; and all trace of Contraction in the Palm of the Hand had disappeared. The Little Finger, which never could be brought quite straight, from alterations in the joint, said to be the result of injury, remains contracted between the first and second Phalanges. From a cast.

divided subcutaneously, and the disposition to recontraction prevented. This contrasts very favourably with the relapsed cases after open wound, which from the nature of the cicatricial contraction are incapable of further relief.

It has been my object in the present paper to give to the profession an increased confidence in the subcutaneous operation for Dupuytren's contraction of the fingers, and to point out a number of details in the mode of performing the operation and in the after-treatment, upon which I believe much of the success will be found to depend.

THE TREATMENT OF ACNE.

By JAMES STARTIN, M.R.C.S.,
Surgeon to t. John's Hospital for Diseases of the Skin.

AUTHORS and writers, from the earliest ages up to the present day, seem to have considered the successful treatment of acne, particularly that form known as acne rosacea, to be of some considerable difficulty. I have been much rewarded lately, in hospital and private practice, with some very good results by the carrying out of the following mode of treatment.

In the first place, vegetable acids, mineral waters, purgatives, arsenical preparations, and other specifics, are not indicated in this eruption of the skin, nor have I ever found them of much avail. The diet should be carefully regulated, all indigestible food avoided; and little or no stimulant should be allowed, except in those cases where the powers of digestion should require slight excitation to perform their proper function; and the discriminate use of chalybeates, combined with mineral acids, vegetable bitters, and tincture of iodine, I consider to be the most useful internal remedies. These are considerably augmented by the sulphur-vapour douche or vapour-bath.

But it is the external treatment that is most important, and it is to this that I wish to draw more particular attention. If, at the onset of the affection, the sebaceous glands and follicles become overloaded, they should be relieved by pressure between the finger and thumb-nail, and by frequent washings with warm water and oatmeal; after which a good rubbing with a flesh-brush will remove the contents of a

number of the pimples; and, if the disease be more advanced and the indurated spots become enlarged and painful, the vapour-douche is then of much service, followed by an ointment of the ammonio-chloride of mercury and camphor, or the gentle application of iodide of sulphur ointment at bedtime, and the use of a cooling zinc or calamine lotion, to be painted upon the face with a camel-hair brush two or three times a day. With the judicious use of the above remedies and applications, I may probably nearly remove and cure the disease; but troublesome redness, with often great vascularity, still remains in spite of treatment, and it is to this stage of the affection my remarks tend. To those small indurations and superficial venous enlargements and plexuses, I generally apply the acid nitrate of mercury with care by means of a spun-glass brush. One or two applications, with the immediate use of bibulous or blotting paper afterwards, will usually suffice to effect complete obliteration without a scar; and I find, after a few days, the morbid states of the capillaries have undergone a complete change for the better. Nevertheless, we may still have some of the larger capillaries, which keep up the localised redness and induration, to contend against; then the mercurial acid treatment does not hold good. I therefore divide each engorged vesicle with the point of the lancet; and, should the hæmorrhage be much, I apply a small ring of silver or steel, which I have fixed into a handle at right angles, and about the eighth of an inch in diameter, over the two divided ends of the vessel. I then insert a minute grain of nitrate of silver, which at once stops the bleeding and obliterates the engorged conspicuous vesicle, leaving only a small black discoloured spot, which may be easily removed by the application of a solution of iodide of potassium.

The above treatment may appear somewhat heroic for so common an eruption. I can only state that I have made use of it in a large number of cases, and find it to be the most successful mode of cure.

GOITRE AND THE HÆMORRHAGIC TENDENCY.

By R. BRUCE LOW, M.D.,

Medical Officer of Health, Helmsley Rural Sanitary District.

IN the course of my reading, I have been unable to find any mention of the fact that goitrous persons are peculiarly subject to hæmorrhages. As I reside in a district where goitre abounds, I have had opportunities of remarking the frequency of flooding among the women, and more especially among the women with goitres. During the last few months, I have collected notes of one hundred and eighty-three cases of goitre. Of these, there were ninety women who had borne children, and, out of these, thirty-one were habitual flooders; besides these, a considerable number of others showed a hæmorrhagic tendency, especially at their menstrual periods. There appears to be a great predisposition to flooding in the district, so much so, that the medical men of the neighbourhood are united in their opinion that more care and attention are requisite after labour is over here than in other districts. Even with the greatest care, flooding sets in. For example: in a case where a goitrous woman was confined safely and the placenta removed, I grasped the contracted uterus through the abdominal wall, and held it firm for more than an hour; but, feeling exhausted, I transferred my post to an attendant, with the effect of seeing the woman flood and faint exactly four minutes after I left hold of the uterus. In cases, again, where labour is over before the arrival of the medical man, flooding is very apt to go on to a very serious extent. The tendency to flood is not entirely confined to goitrous women in the district, but these are usually the worst cases.

The etiology of goitre is still far from being clear; but it is now accepted that the disease is endemic in certain well-defined geological districts, and also that it may be produced by some bad hygienic conditions, *e.g.*, dampness, overcrowding, and bad ventilation. The water-supply of the Helmsley district is pure and good, but, owing to its situation on the oolitic lime-stone formation, the water contains carbonate and sulphate of lime and, in smaller quantity, magnesia. The permanent hardness is not great, the water being used for washing and other domestic purposes. It contains no trace of iron. The district is extensively wooded, and is intersected by numerous small valleys, which are constantly filled with damp foggy emanations. The dwellings are small, badly constructed, ill-ventilated, and generally overcrowded. Many of them are in bad repair. The people, for the most part, are engaged in agricultural pursuits. Intermarriage has hitherto prevailed to a great extent, the isolated position of the district having prevented the inhabitants from mixing with those of other neighbourhoods.

Many medical men have called attention to the evils which arise from the continued use of water impregnated with lime and magnesian salts. Dr. Murray, in an able paper in the *BRITISH MEDICAL JOURNAL* for