lene, they could not keep it—it very soon changes; therefore, it has a certain amount of impurity, a certain admixture of alcohol with it, by which alone it can be kept. If true pure bichloride of methylene could be made, it would cost about a guinea an ounce, and would not keep half an hour, so that this is a mixture of pure bichloride and spirit. I have seen it made. The chloroform and alcohol are placed in a retort with sheets of zinc and boiled; as distillation goes on, one atom of chlorine leaves the chloroform, and the zinc is converted into a chloride. Some people say it is only a mixture of chloroform and spirit; but that certainly is not true, because the vapour of it burns, while, as you very well know, the vapour of chloroform does not.

The nurses have sponges and water all ready—sponges of a certain size, thoroughly well cleansed and in sufficient number, neither too large nor too small. If they be too small they may be lost, and if they be too large they cannot be introduced. The assistants are ready. There are only two required: one stands opposite the operator to be prepared to assist him in tying any vessel, and more particularly in preventing the coming out of intestines after the escape of the cyst from the abdominal cavity. If the assistant be not careful as the cyst is drawn out, the intestines follow, and give a good deal of trouble; but, if he carefully hold up the abdominal wall, keeping the edges of the wound together, it is impossible that any intestine can follow the cyst as it escapes. He passes the middle finger inwards under the umbilicus, and the forefinger to the right and the thumb to the left of the wound, and holding the edges closely together as the tumour comes out of the abdomen. When I first saw ovariectomy done, and when I first did it myself, we were told always to have wet flannels and warm water to wrap up the intestines if they came out. I have seen all the intestines of a patient on the table wrapped up in warm wet flannels; the result was a quantity of wool from the flannels adhering to the intestines, which had to be cleaned off before they were returned.

Then, as to the instruments one uses. First, an ordinary scalpel—working rather with the point of the instrument than with the shoulder. Next we have a number of what are called my torsion-forceps, to hold any vessel in the abdominal wall. Supposing a vessel is bleeding, it is caught in a moment, and the forceps hang down holding the vessel, so that, when the peritoneal cavity is opened, no blood drops into it. I have them plated with nickel, so that they never rust. The bleeding vessels having been stopped, the next step is to divide the peritoneum, catching hold of it previously with forceps or by one of these little hooks. The advantage of the hook is that one is less likely to catch a bit of cyst with it. If the cyst be lying close to the abdominal wall when you are catching up the peritoneum with the intention of dividing it, you may catch the cyst also, and sometimes divide it as well as the peritoneum. That is avoided by using this hook. One or two flat touches of the scalpel on the peritoneum are sufficient to divide it. Then a broad director is passed into the opening, and, with a blunt-pointed knife, the peritoneum is very safely divided. I rather insist upon this blunt point, because, with a sharp-pointed instrument, supposing there is a bit of intestine adhering to the abdominal wall, it may be injured, or a sharp-pointed knife might enter the bladder if it were high up. Having laid bare the cyst by the incision of the peritoneum to the extent of three or four or five inches, it then becomes necessary to empty the cyst, and this is done by the form of trocar I have here, which is just one of the trocars I showed you for tapping, but of a larger size, and furnished with outer spring-hooks to fasten the cyst. It is passed into the cyst, then the point is withdrawn, and the fluid rushes through the cannula into the pail below the table. As that is done, the outer hooks are opened, the cyst is caught hold of and easily fastened to the cannula, and pulled out through the opening in the abdominal wall; the fluid passes out, the cyst is held by these grasping claws, and, if the cyst be free, it of course readily and easily follows the instrument. If any adhesion be noticed as the cyst comes out, it may be separated; but I must reserve any further description of that till the next lecture.

BASFORD RURAL DISTRICT.—Mr. Whitgreave commences his report by congratulating his sanitary authority because they will soon have what they have patiently waited for for years—viz., a plentiful supply of pure water. The birth-rate is calculated to have been 42.64, and the death-rate 19.58, or 19.36 if 518 colliers who came into the district in 1877 be included in the population. There were 1,986 births and 912 deaths, so that the births were more than double the deaths. Inflammatory diseases of the lungs caused a large mortality, especially in the Greaseley registration district, which Mr. Whitgreave cannot account for. The mortality amongst infants was high, as 35 per cent. of the total deaths occurred amongst children under one year old; and 16.4 deaths under one year were registered to each 100 births, which is much above the average of rural districts in 1877. There is a full account given as to the removal of nuisances by the inspector.

CONTRACTION OF THE FINGERS (DUPUYTREN'S CONTRACTION),

AND ITS SUCCESSFUL TREATMENT BY SUBCUTANEOUS DIVISIONS OF THE PALMAR FASCIA, AND IMMEDIATE EXTENSION.*

By WILLIAM ADAMS, F.R.C.S.,

Surgeon to the Great Northern Hospital; Consulting-Surgeon to the National Orthopaedic Hospital; etc.

CONTRACTION of the fingers takes place from a variety of causes; and the pathological conditions will be found to vary according to the nature of the producing cause, such, for example, as local injuries, with laceration of tendons, deep abscesses in the palm of the hand, burn-cicatrices, etc. But in the present paper I propose to direct attention to one form only of finger-contraction, commonly met with in men about the middle or beyond the middle period of life—very rarely at younger ages. I have never seen it in women.

The contraction may affect one finger alone; in which case, according to my observation, the ring-finger is most frequently drawn down towards the palm of the hand, but some authors state that the contraction most frequently commences in the little finger. The contraction having commenced in one finger, however, the adjacent fingers are in most cases gradually drawn down, though to a less extent. I have frequently seen the ring-finger severely contracted, so that its tip would nearly touch the palm of the hand, and the middle and little fingers contracted to about half this extent. I have seen the contraction sometimes limited to the little finger. The index-finger and the thumb usually escape. I have never seen them more than slightly contracted. In these contracted fingers, the articulations are generally in a healthy condition; the joints can be flexed freely, but any attempt at extension is painful, from the resistance offered in severe cases by a tense contracted cord passing from the finger into the palm of the hand, to which the skin of the palm is closely adherent. The skin in the palm of the hand near to the base of the contracted fingers is commonly drawn into thick knotty folds.

The pathology and treatment of this form of finger-contraction always has been and still is the subject of much difference of opinion; and Dupuytren appears to have been the first surgeon to investigate the anatomical conditions by dissection. In the report of his clinical lectures,† it is stated that when Dupuytren heard of the death of a man who was the subject of this affection, he “was determined that this remarkable disease should not remain unknown. The arm was given to him, and he made a careful dissection. The skin having been removed from the whole extent of the palm of the hand and the palmar face of the fingers, the fold or the puckering of this structure entirely disappeared; it was, therefore, very evident that the arrangement which presented during the disease did not depend on this cause, but was communicated to it; but how, and by what? The dissection being continued, the Professor discovered the palmar aponeurosis extended, retracted, and diminished in length, its inferior part being divided into cords, which passed on to the sides of the affected finger. In extending the finger, he observed that the aponeurosis underwent a kind of tension of crispation—this was a ray of light; so he considered the aponeurosis was something in the effects [to be the cause, W. A.] of the disease. He cut the prolongations on the sides of the fingers, and immediately the contraction ceased, the fingers returned to a state of semiflexion, and by slight force to complete extension. The tendons were natural; the sheaths were not open; the articulations, ligaments, synovial membranes, and bones, were in their normal or natural state.” Since Dupuytren made this important contribution to our knowledge of this affection, it has been sometimes spoken of “Dupuytren's finger-contraction”: a title as useful as it is also a just compliment to the great surgeon, distinguishing it from all other forms of finger-contraction.

M. G. Goyrand‡ of Aix also records the dissection of the hand of a man, seventy-two years of age, affected with this form of finger-contraction; and the description agrees very closely with that of Dupuytren, although he regards the lateral bands extending along the phalanges not as digital prolongations of the palmar fascia, but abnormal fibrous fasciculi, which extend from the fascia to the sheaths of the flexor tendons, into which they are inserted opposite the second phalanx.

* Read before the Royal Medical and Chirurgical Society, May 22nd, 1877.

† Selections from the clinical lectures delivered at the Hôtel Dieu, Paris; delivered in the session 1831 and 1832, by Baron Dupuytren. *London Medical and Surgical Journal*, vol. i, p. 267. London: 1832. Renshaw and Rust.—The observations of Dupuytren on this subject are fully recorded in his *Leçons Orales de Clinique Chirurgicale*, tome premier, p. 1. Bruxelles: 1832.

‡ *Gazette Médicale de Paris*, 1835, p. 481.

In the Museum of King's College is a dissected specimen of one of these finger-contractions, No. 1444-3 (see fig. 1), presented by the late Professor Partridge. In this specimen, the contraction is limited to the little finger, and shown to depend upon a strong contracted band of the palmar fascia *a*, by which the finger is drawn towards the palm of the hand, the fascial band stretching across like the string of a bow, and passing along the outer side of the little finger along the first phalanx to the beginning of the second. The flexor tendons *b* are seen lying deeply along the concavity of the curves close to the bones.

This dissection has also been figured in Druitt's *Surgeon's Vade-Mecum*, eleventh edition, fig. 301, and I am indebted to Dr. Druitt for permission to reproduce it here (see fig. 2). In this drawing, the direc-

Fig. 1.—Dissection of a contraction of Little Finger: from specimen in Museum of King's College. *a*. Contracted Band of Palmar Fascia, stretching across like the string of a bow; *b*. Flexor Tendons lying deeply along the concavity of the curve, close to the bones.

Fig. 2.—Another view of the same dissection as Fig. 1, showing the contraction to depend upon a band of the Palmar Fascia. (From Druitt's *Surgeon's Vade Mecum*, Fig. 301.)

tion and relations of the contracted band of the palmar fascia are more distinctly shown, but the relation of the fascia to the tendons had not been demonstrated by dissection at the time when this drawing was made.

There is a cast in the Museum of King's College apparently taken from the same subject previously to the dissection, as it corresponds in all respects with the contraction of the little finger, and shows the lateral direction of the contracted fascial band. (Fig. 3.) This cast has no number, but is engraved June 1853, and I am informed that Mr. Partridge always used it in lecture, together with the dissection.

It has never occurred to me to have the opportunity of dissecting one of these finger-contractions; but on April 15th, 1864, a gentleman, Mr. L., aged 50, residing at Hampton Court, who suffered from Dupuytren's contraction of the fourth and fifth fingers in each hand, and upon whom I had proposed to operate, met with an accident. In attempting to hold a restive horse, the contracted fingers on the right hand were suddenly torn open and the skin in the palm of the hand torn across. I saw him shortly after the accident, when the hand had been merely tied up by a handkerchief, and found a large gaping wound in the palm of the hand, reaching nearly half way across. The palmar fascia had been torn across, together with the skin, but the sheaths of the tendons were not torn, and it was evident that these structures had not been implicated in the contraction. The tendons in their sheaths were seen lying at a depth from the surface, running along the concavity of the curve in proximity with the bones, whilst the fascia had evidently been stretched across, like the string of a bow. After cutting away a few torn shreds of fascia, I found, on attempting to extend the fingers, that the previously transverse wound assumed a lozenge shape, in a perpendicular direction. I therefore approximated the edges laterally, and introduced sutures, so that when

sewn up it resembled a longitudinal instead of a transverse wound. The hand was firmly bound round with narrow strips of plaster, and bandaged to a splint, with the fingers in an extended position. On the third day, the wound presented a healthy appearance, without any suppuration, and I therefore removed the sutures; the healing process proceeded without interruption, and the fingers remained nearly straight, without their power of flexion being lost.

These, so far as I know, are the only facts to which we can appeal for the anatomical conditions presented in Dupuytren's finger-contraction; and they all agree in negating the supposition that the tendons or sheaths of the tendons are involved in these contractions. This is also confirmed by my own clinical experience, as I have invariably found that, after dividing subcutaneously a tense and prominent cord, which has generally been mistaken for a tendon, in the palm of the hand, the patient is immediately able to flex the finger as strongly as before the operation, at once disproving the opinion often expressed by surgeons that such fingers will remain stiff and inflexible after the operation.

The commencement of this form of contraction is sometimes to be traced in one hand, in patients in whom the contraction exists in a severe form in the other, and I have watched its commencement from a flattened nodular induration in the palmar fascia in or just above the transverse crease in the palm of the hand corresponding to the metacarpophalangeal articulations. In the beginning, the skin is not adherent to this flattened induration; but in one case now under my observation, the commencement of a fascial-cord contraction can be traced, leading from the transverse crease to the little finger in the left hand: a spot certainly not exposed to pressure.

In the *second stage*, the skin becomes adherent to the thickened fascia, and a puckered dimple is produced, extending from which towards the finger a thickened band of fascia can be distinctly felt.

The *third stage* consists of an increased thickening of the palmar fascia and the band leading to the finger; and also the formation of a thick cord-like band of fascia, leading from the central thickening towards the annular ligament of the wrist, accompanied with a gradually increasing drawing down of the finger or fingers towards the palm of the hand. Such I believe to be the gradual progress of the fascial contraction and its adhesion to the skin in Dupuytren's finger-contraction.

Cause.—With regard to the cause of this affection, most authorities agree in assigning it, in the great majority of cases, to a local cause, and believe it to be produced by pressure from the use of tools in various occupations; and it is said that carpenters, gardeners, and gunners, are specially liable to it. Sir James Paget refers to the elder men occupied in wire-drawing and lock and key-making, as being subject to this condition. It is also generally admitted that this form of contraction may take place from constitutional causes, independently of any local cause, and that it is then traceable to a gouty diathesis. Sir James Paget* points out the dependence of this contraction in some cases upon the gouty diathesis, and believes that the adhesion of the palmar fascia to the adjacent sheaths of tendons and the integuments forms a point of diagnosis. My own opinion is that it nearly always depends upon a constitutional rather than on any local cause, and essentially I regard it as depending upon a gouty diathesis. In favour of this opinion, I would refer, first, to the class of patients in whom it occurs. During a connection of more than twenty years with the Royal

Fig. 3.—Cast from the Hand represented in Fig. 1, previous to dissection.

* Abstract of Clinical Lectures delivered at St. Bartholomew's Hospital (BRITISH MEDICAL JOURNAL, May 22nd, 1875, p. 665).

Orthopædic Hospital, I have seen but very few cases of Dupuytren's contraction in the labouring class; and the cases that did present themselves generally occurred in butlers and in-door servants. It seems, however, to be an affection of common occurrence in the middle and upper classes of society. The cases which have fallen under my observation have occurred in clergymen, barristers, medical men, officers of the army and navy, and merchants; the only condition common to the whole series being a disposition to gout. Coexisting with the finger-contraction, in the cases which I have seen there have been generally other manifestations of a gouty tendency, more especially to that form we recognise as rheumatic gout affecting several articulations and often causing enlargement of the joints of the fingers rather than true inflammatory gout, affecting the great toe. Secondly, I would refer to the frequent occurrence of this affection in the left hand only, and to its occurrence in both hands, which we can hardly explain by any local cause. Thirdly, I would refer to the fact that in several instances I have known two brothers suffer from it; and in some cases, the father and son have been similarly affected; so that, for these reasons, I am disposed to attach far greater importance to the constitutional than to any local cause.

Treatment.—With regard to treatment, this must be either mechanical or operative; and there can be no doubt that gradual mechanical extension, by an apparatus worn night and day, or in slight cases only at night, would not only prevent increase of the contraction, but diminish, and possibly even cure it when slight and not of long duration. But in severe cases, and in those of long standing, mechanical treatment is useless, the only prospect either of benefit or cure being from operative treatment either by open wound or the subcutaneous method.

Dupuytren, in the year 1831, divided the contracted palmar fascia by open wound, making an incision ten lines in length; but a second and even a third incision was found to be necessary. Dry charpie was applied, and the fingers extended. Inflammation and suppuration followed; and at the end of nearly two months, it is reported that "the fingers regained their natural condition".* Two other successful cases are also reported in the same lecture.

Mr. South,† in his translation of Chelius, states that "Goyrand does not divide the skin transversely, as Dupuytren does, because in straightening the finger, the cut in the skin gapes too much; but he cuts through it longitudinally and through the bridge transversely".

Dr. Otto W. Madelung‡ has recently published an account of the treatment of these cases adopted in the Surgical Hospital at Bonn by Professor Busch. This operation belongs to the class of open wounds, and consists in dissecting up a triangular flap of skin from the contracted cord in the palm of the hand, and then dividing all the bands of the contracted fascia which can be reached; as the flap thus raised contracts, the lower points of the wound may be united by sutures. "A light bandage closes the wound; the hand is then kept by the patient in a sling, without the slightest attempt to preserve the extension. . . . Extension-movements with the fingers are only made when the wound has entirely granulated, and then only in a light gentle way. At first, wood cylinders of various sizes are laid in the hand; later, the hand is stretched on a back splint. Active and passive movements are now to be made in, and during the time the hand-bath is used for cleansing the wound. More complicated apparatus than the above are never required. The healing of the wound, accelerated perhaps by skin-grafting, is accomplished in three or four weeks."§ Successful cases under treatment are referred to, but one is added by the author in which, notwithstanding the use of Lister's antiseptic treatment, suppuration with sloughing of the flexor tendon of the little finger took place; but this is stated to be the only unfavourable case.

Professor Alfred C. Post|| of New York also operates upon these cases by open wound, making incisions at a number of points, but not longer than absolutely required, as he believes that the adhesion to the skin prevents a strictly subcutaneous section from being made. He advocates immediate extension on a metal splint; the dressings to be removed every two or three days, and passive motion applied. Of the four successful cases cited by Professor Post, only one seems to have been of Dupuytren's contraction, the others depending on abscesses and traumatic inflammation.

All operations by open wound for these cases should be condemned as unnecessarily severe, involving a long and tedious reparative process, with the risk of suppurative inflammation, and also a liability to failure, in which event the condition of the patient would be worse than before the operation; contraction from cicatrix being one of the most difficult conditions to relieve. Nevertheless, in England, the operation by open wound has long been practised by the majority of surgeons, and is still recommended in some of the text-books, when the subcutaneous operation has failed, as it is generally supposed to do. There is, however, in England, a very general distrust in all operative procedures in the treatment of this affection, which is regarded as incurable, the patients being advised to bear the ills they have, rather than run any risk from suppurative inflammation, with the doubtful gain of a stiff and useless finger.

The operation which I now propose to bring before the notice of the profession is not based upon any new principle in surgery, as the subcutaneous division of the palmar fascia for contracted fingers has been practised, both in this country and on the Continent, for many years. Within the last few years, I have materially modified the details, both with regard to the mode of performing the operation and the after-treatment. The success has been found much greater, and the treatment much less irksome to the patient, as well as of shorter duration, than the method of operating and the after-treatment which I adopted in the earlier part of my practice from the teaching of my late colleagues and predecessors at the Orthopædic Hospital, Mr. Tamplin and Mr. Lonsdale. They believed that both fascia and tendon were involved in the contraction, and divided the contracted cords in the palm of the hand in the same way as they divided tendons: subcutaneously, cutting from below upwards and making only one puncture. They also adopted the same method of after-treatment, by gradual mechanical extension; using complicated steel instruments, with cog-wheels opposite the joints of the fingers. Further study of these cases, and the opportunity which was accidentally afforded me of seeing that the tendons were not involved in a severe case of finger-contraction previously referred to, in which the palm of the hand was torn across by a horse, confirmed me in the belief that the tendons were not implicated in this form of contraction of the fingers, and that surgically we had to deal with contraction of the palmar fascia alone, as Dupuytren had described the affection. I therefore commenced the practice of making multiple subcutaneous divisions of the fascia, introducing the smallest tenotomy knife—smaller than any ordinarily used, as shown in fig. 4—and carrying it between the skin and the contracted cord, which I then divide by cutting downwards very slowly and cautiously, taking care not to dip the point or divide any structures, except the contracted band of fascia. In one case, where three fingers were contracted, I made as many as nine punctures and subcutaneous divisions; but where so much is required to be done, I think it generally advisable to confine the operation to one or two fingers, making not more than five or six punctures—in many cases, four will be found sufficient—and leave the other finger or fingers to be treated at a subsequent operation. As a rule, I make the *first puncture* at the greatest distance from the finger, in the palm of the hand, towards the annular ligament, a little removed from the point where the skin is adherent to the fascia, and where the skin is not tightly stretched over the contracted cord, so that the tenotomy-knife can be readily introduced between the two. The *second puncture* should divide the same cord as the first, but as near to the finger as possible, thus leaving the contracted band in the palm of the hand, when adherent to the skin, isolated and cut off from its connections at its upper and lower extremities. The *third and fourth punctures* divide the lateral bands or digital prolongations of the palmar fascia, which usually pass from the central cord in the palm to the adjacent sides of the fingers. These must be divided very carefully in order to avoid cutting the vessels and nerves along the sides of the fingers; the punctures should be made at the bifurcation of the cutaneous net between the fingers, and the incisions directed obliquely upwards and outwards towards the palm of the hand. These incisions will divide the strongest and most prominent bands, which produce the flexion of the first phalanx of the finger upon the hand, and if care be taken to avoid dipping the point of the knife, there will be no fear of wounding vessels or nerves. Sometimes lateral bands of contracted fascia require to be divided opposite the centre of the first phalanx, and this must be done by puncture at the edge of the contracted bands, the knife being directed transversely towards the bone; but this cut must be made very carefully, to avoid the artery and nerve; the surgeon remembering that the band, though tough and strong, is at the same time very thin. These bands are more readily cut through with the scissor-pointed tenotome, *i.e.*, with the point at the extremity of the straight cutting edge, represented in fig. 4. The central-pointed tenotome generally used does not divide

* *Op. cit.*, p. 267.

† *A System of Surgery*, by J. M. Chelius; translated by John F. South. Vol. ii, p. 194. London: Renshaw. 1847.

‡ *The Causes and Operative Treatment in Dupuytren's Finger Contraction*, by Dr. Otto W. Madelung; translated from the German. London: Trübner and Co. 1876.

§ *Op. cit.*, p. 11.

|| "On Contraction of the Palmar Fascia, and of the Sheaths of the Flexor Tendons", by Alfred C. Post, M.D., of New York (*Archives of Clinical Surgery*, August 1876)

the fascial bands so readily or with the same precision. Occasionally a lateral band may have to be divided between the first and second phalanx, as the sheath of the tendons, or the tendons themselves, may be readily divided by such incisions, and would lead to a loss of the power of flexing the finger after the operation. I

Fig. 4.—Small Knives used for division of the Bands of Fascia.

always avoid making any central incisions in front of either the first or second phalanx, as the sheath of the tendons, or the tendons themselves, may be readily divided by such incisions, and would lead to a loss of the power of flexing the finger after the operation.

Subsequently, I also adopted another important modification in the after-treatment of these cases; viz., the plan of immediate extension, bringing the finger or fingers as nearly as possible into the fully extended position at the time of the operation, and afterwards applying

Fig. 5.—Retentive Metal Splint to which the Fingers and Hand are bandaged after the operation.

as a retentive apparatus a well-padded metal splint bent to the shape of the wrist, hand, and fingers, and applied along the palmar surface; the finger or fingers, as well as the hand and wrist, being bandaged to the splint, as shown in fig. 5. This was a great change from the prolonged after-treatment required by the method of gradual mechanical extension made with the complicated steel instrument with cog-wheels.

I was led to make this alteration from the gradual to the immediate mechanical extension after considering the pathological conditions, and the objects to be accomplished with regard to the reparative process in fascia, as distinguished from the reparative process in tendon.

The object of immediate extension is to widen as quickly as possible the gaps made by the incisions in the fascia, with the view of preventing union of the divided fascia; or to have the union as feeble as possible: objects precisely the reverse of those we desire to obtain after the subcutaneous division of tendons.

Since I have adopted these modifications in the operation

and after-treatment, I have had good reason to be satisfied with the results, and in the great majority of the cases operated upon, the contraction has been completely and permanently removed.

Anæsthetics.—With regard to the employment of anæsthetics in these operations, which must necessarily be done slowly and very carefully, several punctures being made, I have always advised the use of chloroform or ether. I abandoned the use of ether-spray, after employing it some years ago, because it was found to harden the skin, so as to make the puncture difficult and obscure the anatomical relations of the parts. It occurred to me, however, that if I rapidly thawed the frozen skin by rubbing it with my own hand, the deep anæsthesia would probably remain long enough for a subcutaneous operation; and this I have found to occur. In the case of a medical friend, Mr. R., upon whom I operated for contracted fingers in January 1877, and who objected to take either chloroform or ether from the condition of his circulation, I made four punctures, and he only felt one of them, and that very slightly. The objection of hardening the skin, therefore, being done away with, the local anæsthesia by the ether spray is now found to be applicable to tenotomy and some other subcutaneous operations, as its inventor, Dr. Richardson, originally thought it would

be. The hand of the gentleman alluded to, Mr. R., is represented in fig. 6, showing severe contraction of the middle and ring fingers of the right hand previous to operation; and its improved condition one

Fig. 6.—Hand showing Dupuytren's Contraction of Middle and Ring Fingers, with prominent Cord in Palm of Hand. From a cast.

Fig. 7.—The same Hand as shewn in fig. 6, one year after operation. From a photograph.

year after the operation is shown in fig. 7. The fingers are perfectly straight and useful, the voluntary power not having been interfered with by division of tendons.

A severe degree of contraction of three fingers is represented in fig. 8; and the same hand is shown in fig. 9, thirteen years after the operation. The middle and ring-fingers remain completely cured, with full power of flexion, and all trace of contraction in the palm of the hand has disappeared. The little finger, which never could be brought quite straight from alterations in the joint said to be the result of injury, remains contracted between the first and second phalanges. This gentleman held and still continues to hold a consular position, and the contraction of three fingers in the right hand so much interfered with his ability to write that his signature was scarcely legible. His writing powers were completely restored by the operation.

Summary.—The operation and treatment which I now practise may be described as follows.

1. The subcutaneous division of all the contracted bands of fascia which can be felt; the bands to be divided by several punctures, with the smallest tenotomy-knife passed under the skin and cutting from above downwards: a pledget of lint being at once placed over each puncture and retained in position by a strip of plaster.

2. Immediate extension to the full extent required for the complete straightening of the fingers, where this is possible, and the application of a retentive well-padded metal splint from the wrist along the palm of the hand and the fingers; the fingers and hand to be bandaged to the splint.

3. The bandage not to be removed until the fourth day, when the lint and plaster may also be taken off, as the cutaneous punctures are always found to be healed by the fourth day. The retentive metal splint to be reapplied, and the hand and fingers bandaged to it.

4. Extension to be kept up by the splint worn continuously night and day for two or three weeks; but the splint and bandage to be changed every two or three days. After this, the extension splint is to be worn at night only, for an additional three or four weeks, free motion being encouraged during the day.

Relapse of the deformity. I believe, is now guarded against by the plan of dividing all the contracted bands of fascia by as many punctures as may be necessary; and, in severe cases, not attempting too much at one operation; and also by the adoption of the plan of immediate ex-

tension. If partial relapse should occur—and I have never known more than this—any bands that may have escaped division, or any which may since have become prominent by the extension, may be

Fig. 8.—Hand showing Dupuytren's Contraction of Middle, Ring, and Little Fingers. From a cast.

Fig. 9.—The same Hand as shown in fig. 7, thirteen years after operation. The Middle and Ring Fingers remained completely cured, with full power of Flexion; and all trace of Contraction in the Palm of the Hand had disappeared. The Little Finger, which never could be brought quite straight, from alterations in the joint, said to be the result of injury, remains contracted between the first and second Phalanges. From a cast.

divided subcutaneously, and the disposition to recontraction prevented. This contrasts very favourably with the relapsed cases after open wound, which from the nature of the cicatricial contraction are incapable of further relief.

It has been my object in the present paper to give to the profession an increased confidence in the subcutaneous operation for Dupuytren's contraction of the fingers, and to point out a number of details in the mode of performing the operation and in the after-treatment, upon which I believe much of the success will be found to depend.

THE TREATMENT OF ACNE.

By JAMES STARTIN, M.R.C.S.,
Surgeon to t. John's Hospital for Diseases of the Skin.

AUTHORS and writers, from the earliest ages up to the present day, seem to have considered the successful treatment of acne, particularly that form known as acne rosacea, to be of some considerable difficulty. I have been much rewarded lately, in hospital and private practice, with some very good results by the carrying out of the following mode of treatment.

In the first place, vegetable acids, mineral waters, purgatives, arsenical preparations, and other specifics, are not indicated in this eruption of the skin, nor have I ever found them of much avail. The diet should be carefully regulated, all indigestible food avoided; and little or no stimulant should be allowed, except in those cases where the powers of digestion should require slight excitation to perform their proper function; and the discriminate use of chalybeates, combined with mineral acids, vegetable bitters, and tincture of iodine, I consider to be the most useful internal remedies. These are considerably augmented by the sulphur-vapour douche or vapour-bath.

But it is the external treatment that is most important, and it is to this that I wish to draw more particular attention. If, at the onset of the affection, the sebaceous glands and follicles become overloaded, they should be relieved by pressure between the finger and thumb-nail, and by frequent washings with warm water and oatmeal; after which a good rubbing with a flesh-brush will remove the contents of a

number of the pimples; and, if the disease be more advanced and the indurated spots become enlarged and painful, the vapour-douche is then of much service, followed by an ointment of the ammonio-chloride of mercury and camphor, or the gentle application of iodide of sulphur ointment at bedtime, and the use of a cooling zinc or calamine lotion, to be painted upon the face with a camel-hair brush two or three times a day. With the judicious use of the above remedies and applications, I may probably nearly remove and cure the disease; but troublesome redness, with often great vascularity, still remains in spite of treatment, and it is to this stage of the affection my remarks tend. To those small indurations and superficial venous enlargements and plexuses, I generally apply the acid nitrate of mercury with care by means of a spun-glass brush. One or two applications, with the immediate use of bibulous or blotting paper afterwards, will usually suffice to effect complete obliteration without a scar; and I find, after a few days, the morbid states of the capillaries have undergone a complete change for the better. Nevertheless, we may still have some of the larger capillaries, which keep up the localised redness and induration, to contend against; then the mercurial acid treatment does not hold good. I therefore divide each engorged vesicle with the point of the lancet; and, should the hæmorrhage be much, I apply a small ring of silver or steel, which I have fixed into a handle at right angles, and about the eighth of an inch in diameter, over the two divided ends of the vessel. I then insert a minute grain of nitrate of silver, which at once stops the bleeding and obliterates the engorged conspicuous vesicle, leaving only a small black discoloured spot, which may be easily removed by the application of a solution of iodide of potassium.

The above treatment may appear somewhat heroic for so common an eruption. I can only state that I have made use of it in a large number of cases, and find it to be the most successful mode of cure.

GOITRE AND THE HÆMORRHAGIC TENDENCY.

By R. BRUCE LOW, M.D.,

Medical Officer of Health, Helmsley Rural Sanitary District.

IN the course of my reading, I have been unable to find any mention of the fact that goitrous persons are peculiarly subject to hæmorrhages. As I reside in a district where goitre abounds, I have had opportunities of remarking the frequency of flooding among the women, and more especially among the women with goitres. During the last few months, I have collected notes of one hundred and eighty-three cases of goitre. Of these, there were ninety women who had borne children, and, out of these, thirty-one were habitual flooders; besides these, a considerable number of others showed a hæmorrhagic tendency, especially at their menstrual periods. There appears to be a great predisposition to flooding in the district, so much so, that the medical men of the neighbourhood are united in their opinion that more care and attention are requisite after labour is over here than in other districts. Even with the greatest care, flooding sets in. For example: in a case where a goitrous woman was confined safely and the placenta removed, I grasped the contracted uterus through the abdominal wall, and held it firm for more than an hour; but, feeling exhausted, I transferred my post to an attendant, with the effect of seeing the woman flood and faint exactly four minutes after I left hold of the uterus. In cases, again, where labour is over before the arrival of the medical man, flooding is very apt to go on to a very serious extent. The tendency to flood is not entirely confined to goitrous women in the district, but these are usually the worst cases.

The etiology of goitre is still far from being clear; but it is now accepted that the disease is endemic in certain well-defined geological districts, and also that it may be produced by some bad hygienic conditions, *e.g.*, dampness, overcrowding, and bad ventilation. The water-supply of the Helmsley district is pure and good, but, owing to its situation on the oolitic lime-stone formation, the water contains carbonate and sulphate of lime and, in smaller quantity, magnesia. The permanent hardness is not great, the water being used for washing and other domestic purposes. It contains no trace of iron. The district is extensively wooded, and is intersected by numerous small valleys, which are constantly filled with damp foggy emanations. The dwellings are small, badly constructed, ill-ventilated, and generally overcrowded. Many of them are in bad repair. The people, for the most part, are engaged in agricultural pursuits. Intermarriage has hitherto prevailed to a great extent, the isolated position of the district having prevented the inhabitants from mixing with those of other neighbourhoods.

Many medical men have called attention to the evils which arise from the continued use of water impregnated with lime and magnesian salts. Dr. Murray, in an able paper in the *BRITISH MEDICAL JOURNAL* for