not only in infancy, but even in our student days, early impressions are imprinted indelibly on our memory. I had the good fortune to be one of the first set of dressers for Mr. Hilton when he was appointed one of the surgeons to Guy's Hospital, about the year 1845. No one at that time was more ready or more able to apply correct anatomical knowledge to surgical practice than he. It was his delight, as his *chef d'œuvre* on *Rest and Pain* will fully testify. At this time he had under treatment a young woman with necrosis of the lower part of the ulna, without, as was then thought, any present disease in the carpal joints. Mr. Hilton said, "How would it do to take away the diseased bone, and not destroy the integrity of the carpus, by carefully leaving intact the triangular fibro-cartilage which intervenes between the end of the ulna and the cuneiform bone?" This was done, and a perfect recovery resulted. The fact thus presented to my mind was never forgotten. I referred to it from time to time in my anatomical demonstrations, as a possible operation. It was not until quite recently, however, that I had a precisely similar case to treat, one in which, as you will observe, I have removed the lower inch of the ulna without disturbing any of the carpal joints. Recovery has followed, and now, after nearly fifteen months, you may see for yourselves what an useful hand the patient has. But yet there is danger to be averted, as possibly only the after-history of such a surgical case could foreshadow. The two ulnar muscles, the flexor and the extensor, unrestrained by the resistance of the shortened ulna, are now beginning to show a tendency to draw the hand towards the internal side of the fore-arm, and in spite of all I can do, I fear I must admit that, without the use of some mechanical apparatus, the hand will become less serviceable, by permanent fixed adduction.

The lessons to be learnt from the after-history of these cases will, I hope, be as interesting as I believe they might be made instructive; and I trust that the present condition of these patients will convince you that, in many instances of surgical disease and surgical treatment, careful study of the after-history of each individual case will prove an endless source of accumulated experience.

AN ADDRESS

DELIVERED AT THE OPENING OF

THE SECTION OF OBSTETRIC MEDICINE,

At the Annual Meeting of the British Medical Association,
in Manchester, August 1877.

By W. O. PRIESTLEY, M.D., F.R.C.P.,
President of the Section.

THE HISTORY OF OBSTETRIC MEDICINE IN MANCHESTER.

GENTLEMEN,—When I was honoured with an invitation to accept the Presidency of the Obstetrical Section at the present meeting of the British Medical Association, it struck me that it would be interesting to look back on the part which Manchester has from time to time taken in the advancement of obstetric science, and to recall some of the great names who have contributed to its literature.

Manchester, indeed, has had its school of obstetric medicine as well as its school of politics. It has made its mark both in the present and preceding century, and it seems to me a happy coincidence that the Address in Obstetric Medicine, which is given once only in three years, should be delivered in Manchester, a town so much identified with the progress of the science and art of midwifery, and the Association may be congratulated that the address is to be delivered by so competent an authority as Dr. Barnes.

So long ago as 1773, Mr. Charles White, a pupil of William Hunter and Surgeon to the Manchester Infirmary, wrote a treatise on the *Management of Pregnant and Lying-in Women*, which went through three editions; and, in 1784, he issued a work on the *Nature and Cause of Swelling of the Lower Extremities which Sometimes Happens in Lying-in Women*. This work also appeared in three editions, and was remarkable as one of the first scientific attempts to unravel the pathology of the disease called "phlegmasia dolens" by a careful study of the anatomical changes taking place in the limbs. Mr. White gave a very accurate description of the affection, and adopted the view that it depends on "obstruction, detention, and accumulation of lymph in the limb".

In 1800, Dr. Hull published an extended treatise on the same subject; and, from his introductory remarks, it may be gathered that he first bestowed the term "phlegmasia dolens" on the inflammatory swell-

ing of the legs observed in puerperal women, and assigned it its place in the nosology of Cullen. He expresses a fear lest some of his readers should be puzzled by the new name. He regarded the disease as an inflammatory affection, producing suddenly a considerable effusion of serum and lymph into the cellular membrane of the limb. Whatever may be thought of the pathology of these two authors in view of the more recent researches of Dr. Robert Lee, Dr. Mackenzie, Professor Virchow, and others, there can be no doubt that both Mr. White and Dr. Hull made most important contributions towards elucidating the nature of phlegmasia dolens, and deserve great credit.

Probably no single locality in the United Kingdom has contributed so much as Manchester and its vicinity to the discussion and elucidation of the subject of Cæsarean section. In 1769, Mr. Wood here performed the operation upon a woman whose pelvis became greatly distorted by mollities ossium after she had previously borne children naturally. This case led to a discussion, in which Mr. Simmonds, Mr. Ogden, and Mr. Tomlinson of Manchester, with Dr. Sims of London, took part; and out of this controversy came the excellent treatise by Dr. Hull entitled *A Defence of the Cæsarean Operation*, etc. In this work, Dr. Hull entered most intelligently into the nature of the cases requiring the operation, the best method and the proper time for performing it, the sources of danger, and the subsequent treatment of the patient. This book besides gave much information on the subject of pelvic distortions. If, with the light of later experience, exception be taken to some of Dr. Hull's positions, his book may be said to have formed the groundwork of information on Cæsarean section for British practitioners, and to this day it is regarded as containing sound opinions and good practice.

The promulgation of Dr. Hull's views was unfortunately mixed up with a most rancorous controversy between him and Mr. W. Simmonds, in which Mr. Simmonds charged Dr. Hull with being "little acquainted with the decencies common among authors, and ignorant of the language and manners of a gentleman". The contention in bitter personality inevitably recalled the celebrated combat between Susannah and Dr. Slop, graphically described by Sterne in *Tristram Shandy*, and its occurrence is much to be regretted.

In 1801, Dr. Hull translated and published M. Baudelocque's two memoirs on the Cæsarean operation, thus giving English practitioners further information on the subject; and the history of the operation is still further identified with Manchester through the much respected name of Dr. Radford, who delivered the first obstetric address before the Provincial (now named British) Medical Association at Manchester in 1854. Dr. Radford then selected Cæsarean section for the subject of his address, and subsequently published a more extended memoir with an appendix of cases. From this memoir, it appears that, up to a certain period, the greatest number of cases, relatively speaking, in which Cæsarean section had been performed in Great Britain and Ireland had occurred in this city and neighbouring districts. Of fifty-five cases in England and Wales collected by Dr. Radford up to 1865, no fewer than twenty-five had occurred in Lancashire. The occurrence of this large proportion of Cæsarean operations in Lancashire was, no doubt, in a great measure, due to the pernicious influences surrounding the occupation of young girls and women in factories. Cogent evidence of this is to be found in the fact, which I have on good authority, that, since the introduction of the Factory Act, which prohibits girls being employed in factories before a certain age, and limits the hours of work both for women and children, such deformities of the pelvis as necessitate the Cæsarean section have become much rarer, and Lancashire is losing its unhappy pre-eminence in this respect. This, one of the indirect ways in which the Factory Act has ameliorated the condition, and lessened danger and suffering for poor working women in parturition, deserves to be recorded, and must furnish a source of gratification to Lord Shaftesbury and others who were concerned in passing this beneficent measure.

Looking still onward, we learn that to Mr. Kinder Wood of Manchester is probably due the first suggestion that the placenta should be detached from the os and cervix uteri in certain embarrassing cases of placenta prævia, where the severity of the hæmorrhage places the patient in peril, and immediate delivery is difficult or impossible. Sir J. Y. Simpson developed this question more fully later, and impressed upon it, as he did on all subjects he handled, the imprint of his genius. Sir James evidently was not aware, when he made his first communication on this subject, that Mr. Kinder Wood had before him adopted this plan of treatment, or had recognised the fact that complete separation of the placenta would arrest hæmorrhage in cases of placental presentation. Becoming informed later, he made a full acknowledgment, and expressed himself as having "great and sincere pleasure in rendering posthumous justice to the memory of a man of distinguished professional attainments".

And here I may remark that some misapprehension still exists concerning Sir James Simpson's teachings as to the treatment of placenta prævia. It is common enough to hear Sir James accredited with recommending separation of the placenta as a general plan of treatment for cases of unavoidable hæmorrhage. Those who have carefully studied his writings are not likely to fall into such an error; but it may be well once more to point out that Sir James Simpson only proposed the separation of the after-birth before delivery in cases where the hæmorrhage had been so great that the patient could not bear the shock of immediate delivery, or where perilous flooding was associated with such an undilated or undilatable state of the passages as to render speedy delivery impracticable.

These doctrines, it is well known, have been still further modified more recently by the obstetric orator at the present meeting—Dr. Barnes.

Tracing further the history of obstetric medicine in Manchester, I approach delicate ground, inasmuch as many of the men identified with its progress are still living, and their names are familiar to all of us. Of the venerable Dr. Radford, whose name I have already mentioned, and whose first medical diplomas date as far back as 1817, everybody must speak with admiration and esteem. A long life, combined with ardent love of his profession, have enabled him to make many substantial contributions to the literature of obstetrics, and to form a very fine collection of instruments. This collection now belongs to the Radford Museum in Manchester, and, by kind favour, was lent for a brief period to the Obstetrical Society of London for exhibition at its *conversazione* last year. Among Dr. Radford's more important writings, I may mention his *Essays on Various Subjects connected with Midwifery*; his *Essays on Cesarean Section and Deformities of the Pelvis*; *Cases of Torsion, Doubling, and Expulsion of the Fetus in Shoulder-presentations*; *Cases of Laceration of the Uterus*, etc.

Dr. John Robertson, whose first diploma dates as far back as Dr. Radford's, has also, during his life-time, rendered signal service to his profession. His writings bear the stamp of a genius which has been recognised by many eminent authorities, and his practical suggestions have greatly helped to further a more scientific and precise use of midwifery instruments.

Dr. Charles Clay, besides entering the domains of archæology and general science, has brought renown to the Manchester School by the diversity and extent of his contributions to obstetrics. Without going into the question of priority, I may state that Dr. Clay's name is indisputably associated with the earliest successful cases of ovariectomy in this country, and that he materially aided, by his own work and writings, to establish the propriety of an operation which has led to such wonderful results in the hands of Mr. Spencer Wells, Dr. Thomas Keith, and others. Further, Dr. James Whitehead, both by his writings and practice, has attained more than a local celebrity as an accomplished gynaecologist; and Dr. Renaud, who, I understand, does not engage in obstetric practice, yet has added important material to its literature. I have often heard my late friend Dr. Montgomery of Dublin speak of the value of his researches on the corpus luteum, and, in his admirable work on *The Signs and Symptoms of Pregnancy*, Dr. Montgomery says of Dr. Renaud's account of the corpus luteum that, "as far as it goes, it is decidedly one of the best and most accurate".

I may add that, within the last day or two, Dr. Lloyd Roberts has informed me a Mr. Ward, a surgeon in Manchester, was the first to point out that infantile leucorrhœa was an idiopathic affection, and not necessarily produced as the result of carnal intercourse. The importance of this discovery in a medico-legal sense was illustrated by a case occurring at the time. In 1791, Mr. Ward was asked to examine a little girl four years old who was supposed to have been violated. He gave evidence to the effect that there were proofs of such violation, and a man was committed for the capital offence. Immediately afterwards, Mr. Ward saw other cases of infantile leucorrhœa in hospital, which a careful examination convinced him were due to natural causes. He was courageous enough to report this to the authorities, with an avowal that he might have been mistaken as to the evidence he had given in the criminal case. This was explained to the judge of assize at Lancaster, and the man was acquitted.

This brief recital does not at all profess to complete the history of obstetric medicine in Manchester; it merely gives a *résumé* of some of its more salient points up to a given period. In any fuller exposition, one could scarcely omit to mention the names of the present honorary secretaries of this Section. They are the worthy representatives of gynaecology in Manchester at this moment, both as teachers in the medical schools and as pioneers in practice.

Since the last meeting of the British Medical Association in Manchester, considerable strides have been made in perfecting the science

and art of midwifery, and in investigating and treating the diseases of women. New light has been thrown on the treatment of difficult and anomalous labours; fresh expedients have been suggested to meet special emergencies. The subjects of uterine hæmorrhage, the use of midwifery instruments, the diversities in the method of turning, and the pathology of puerperal ailments, have each received fresh illustration in this country by such men as Simpson, Matthews Duncan, Barnes, Braxton Hicks, and others, and by a host of fellow-workers abroad. In the treatment of diseases of women, the advance has even been more marked. The wonderful results which have recently attended the operation for ovarian tumours are the marvel of our time, and indicate a courageous and skilful battle with disease and death, crowned with a success which twenty years ago was scarcely hoped for.

The impetus given to the cultivation of uterine pathology not many years since by the writings of Dr. Henry Bennet in London and by Dr. Simpson in Edinburgh may be said almost to have created the modern school of gynaecology in this country. The various implements introduced into practice, more especially by Simpson, have made diagnosis much more precise and accurate than before. With these improvements, the danger is lest the physician be merged too much into the mere mechanist, and lest he should look on all derangements of the reproductive organs as having too exclusively a local origin to be remedied with undue frequency by local expedients.

As we are avowedly met here for mutual discussion and criticism, I trust I may be pardoned if I indicate in passing one or two pitfalls into which gynaecologists are prone to stumble in the present phase of our art.

In the first place, out of the mechanical improvements in diagnosis and treatment, there seems to have grown too great a proclivity to trust less to the information afforded by the educated fingers and more than is desirable to instruments. These mechanical aids are often employed as a mere matter of routine, without regard to the pathological nature of the uterine disturbance, in the vague hope that they may assist to find out something the nature of which is not well defined beforehand, and apparently with the idea that full justice is not meted out to the patient if any are dispensed with. These remarks apply both to the speculum and sound, with other instruments, but especially to the uterine sound. The uterine sound is a great addition to our resources, but there are many uterine affections in which it affords no information whatever, and many in which it is positively injurious. My experience convinces me that it is employed with unnecessary frequency both as an instrument of diagnosis and for restoring the displaced uterus when a finger in the vagina or rectum would equally well answer the purpose. A sound cannot be introduced into an inflamed and tender womb without stirring up more mischief, and it is often forgotten that a preliminary investigation on this point is as needful and desirable before having recourse to the sound as it is to exclude the chances of pregnancy. The result is the infliction of a large amount of avoidable suffering; sometimes it may be an aggravation of the disease for which the patient seeks relief, and an amount of discredit attaching to the practitioner which most men would be glad to escape.

It is of great moment, now that we possess instruments of acknowledged value, to learn the exact limits within which they render actual service, and when they had better be avoided.

Again, as certain remedies have had their fashion for the day, and then have sunk into unmerited oblivion, so, in defiance of sound principles, a single idea in uterine pathology has been raised into a position of undue prominence and then given place to something else.

At one time, ovarian pathology was in the ascendant; at another time, inflammation of the os and cervix uteri was regarded as the chief ailment from which women suffered, and lately these have given place to the theory that mechanical displacements of the uterus are the root of all evil, and patients taking their cue from the doctors at once jump to the conclusion that all pelvic discomforts come from uterine dislocation. Thus the world is afflicted suddenly with what seems an epidemic of flexions and versions of the womb, and a large amount of time and ingenuity are expended in the invention of pessaries which might be more profitably employed in other directions. Need I point out how important it is that, in all our scientific progress, we should relegate each special subject of study to its proper place in a sound system of pathology, and attach to each morbid condition its true importance? Both in diagnosis and in attempts at cure, while we should have courage enough, when necessary, to face both difficulties and dangers, it comports best with scientific practice to measure as exactly as possible the amount of interference to the needs of the case. All beyond this is not only superfluous; it is apt to be mischievous.

Turning to another subject, I am anxious to draw the attention of members to the subject of transfusion, which has been chosen by our

Manchester *confères* as a topic for discussion in this Section of our meeting. The operation of transfusion has had a checkered career, and, after being forgotten or neglected for long periods, is again attracting much attention.

From a sketch of the history by the late Mr. Pettigrew and others, we learn that the earliest experiments in transfusion were made in France about 1658 by Hansheau. Lower performed the operation in this country in 1665. M. Denis, satirically described as a physician "plus occupé des jeux de hazard, que des jeux de la machine animale", subjected a man to the experiment about the same time. Biva and Manfredi in Italy, and Sinnibaldus in Flanders, repeated the experiment only a few years later. The first four volumes of the *Philosophical Transactions*, which cover the years from 1665 to 1704, contain records of several, but all unsuccessful, cases.

After this time, the practice seems to have fallen into abeyance, although Dr. Harwood of Cambridge published a thesis on the subject in 1785, and endeavoured to rouse attention to its importance. It was not until 1824, when Dr. Blundell of Guy's Hospital reopened the question, that it gained distinct vitality. Dr. Blundell showed, by a series of experiments, that blood may be transfused with success and safety from one animal to another, provided they are the same species; but, if from one animal to another of different species, the result is fatal. MM. Prevost and Dumas in France, and Bischoff in Germany, believed they had afterwards verified Dr. Blundell's conclusions; but Dr. Brown-Séquard has shown that there were fallacies in these experiments, and that, under certain conditions, the blood of other animals may be injected into the human body without danger, while he holds with Dieffenbach that defibrinated blood answers as well as fibrinated blood.

Dr. Blundell performed the operation of transfusion eleven times with his own hand; in ten of these cases there was pressing danger from loss of blood, and in only four of these did he succeed in restoring the patient.

The sparing results attained by Dr. Blundell did not encourage the adoption of the practice by medical men generally. The operation was regarded as delicate, difficult, and uncertain of success, and it languished, although from time to time new instruments were devised and fresh suggestions were made for perfecting the process. In later days, Dr. Hamilton of Ayr, Dr. Graily Hewitt, and Dr. Aveling have each suggested modifications of apparatus. Dr. Aveling's method of immediate transfusion was at once so simple and practical that its introduction gave a new impetus to the operation in this country.

In limine, it may be said that, as increasing attention is paid to the subject, and apparatus becomes more perfected, greater promise is afforded and favourable results are gradually growing. The late Professor Martin of Berlin recorded fifty-seven cases, in which forty-three were completely successful; Dr. Higginson of Liverpool fifteen cases, in which ten were successful; and Dr. McDonnell of Dublin, de Belina of Paris, with others, have published interesting cases.

It will be in the recollection of many members that, at the meeting of the British Medical Association in London, there was a discussion on transfusion, and a lively debate took place between the advocates of *mediate* and the supporters of *immediate* injection.

Just now new life has been imparted to the subject by the appearance of M. Roussel, demonstrating the utility and simplicity of his method for direct transfusion. This subject is obviously a study of deep and increasing interest, as the introduction of the practice in a perfect and simple form may be the means of saving many from the very jaws of death. I commend it to your careful consideration, and, at the appropriate time, to your earnest and dispassionate discussion.

WORTLEY UNION.—This is nominally a rural district; but the bulk of the population are miners or metal-workers inhabiting very large villages, which are much too crowded, and devoid of proper water-supply and means of drainage. Bronchitis and pneumonia caused the largest number of deaths; whilst the mortality from phthisis was below, and that from zymotic diseases was above, the average. The population in 1876 was estimated at 14,880, and the number of deaths 300. The death-rate in 1874 was 20.0; in 1875, 21.7; and in 1876, only 18.6 per 1,000 population. The death-rate under five years of age was 9.4, and above five only 8.93; so that more than 50 per cent. of the deaths occurred amongst children under five, which certainly does not indicate good or even ordinary care of the young. This excessive mortality of children is a common feature in all mining districts, and is usually attributed to carelessness, want of proper food during infancy, and neglect as regards proper clothing and medical attendance. The nuisances abated were 189, but notices were served in thirty cases which were not removed on December 31st.

AN ADDRESS

DELIVERED AT THE OPENING OF

THE SECTION OF PUBLIC MEDICINE,

*At the Annual Meeting of the British Medical Association,
in Manchester, August 1877.*

BY F. DE CHAUMONT, M.D.,

Professor of Military Hygiene, Army Medical School, Netley;
President of the Section.

SCIENTIFIC RESEARCH IN RELATION TO SANITARY PROGRESS.

GENTLEMEN,—It was with feelings of great gratification that I accepted the flattering invitation of the Council to take the chair in this Section, but it was not without some diffidence that I did so. Tempting as it may be to assume that it is a personal honour, I think I ought the rather to take it as a compliment to the institution I represent, the more so as the chair I hold is one that is inseparably connected with the loved and honoured name of Parkes. To him, and to many others of the medical officers connected with the public services of the Crown, a great debt of gratitude is due, both for the work they actually did and for the start they gave to the great principles of sanitation. The opportunities furnished by our troops and seamen for observations on an extended scale have been of great value, and have, I venture to hope, not been neglected; whilst the vast field of observation in our Indian empire has been productive of a great mass of sanitary knowledge. I may refer to the works of two eminent observers—Dr. Timothy Lewis, of the British Service, and Dr. D. Douglas Cunningham, of the Indian Medical Service—who have been employed for a number of years as special commissioners, primarily for the investigation of the causes of cholera, but secondarily for various other inquiries, which have proved of marked importance. Their appointment to this work is greatly to the credit of the Indian Government, and the productive character of their labours has been most encouraging. Among these, I may cite their valuable contributions to the history of cholera, in the course of which they have cleared away much that was of an untrustworthy and misleading character; their researches on the nature of the soil, air, and the ground-water fluctuations of Calcutta; the observations on the suspended matter of the atmospheric air; the discovery of the *Filaria sanguinis hominis*; and their more recent researches on the Madura foot, the Delhi ulcer, and upon Indian leprosy. It is not, of course, my object to praise one class of medical men as sanitarians above another; and, indeed, it would be unbecoming in me to do so; but my desire is twofold: first, to point out the advantages to the community of special scientific inquiries on an independent basis; and, second, to vindicate to some extent the position of our profession in sanitary work.

With regard to the first point—the encouragement of special scientific research—this touches the much vexed question of the "endowment of research", about which so much has been said and written in late years. To a certain extent, the principle has been recognised by the Government in the admirable series of inquiries first carried on under the enlightened direction of Mr. John Simon; but, excellent as these were, and as those still are which are continued under his successor, they are far too limited in scope for the requirements of the age; and it would have been a movement particularly appropriate for a Government whose motto was, "*Sanitas sanitatum, omnia sanitas*," to extend such inquiries in a much wider circle, and so encourage investigations which can seldom be adequately conducted by means of private resources. In private life, those who have the means sufficient have rarely the knowledge or the inclination proper for such researches, whilst "chill penury" too often "represses the noble rage" of those who could most effectually carry them out. Such men as Cavendish are rare, of whom his French biographer said, "*Il était le plus riche de tous les savans, et le plus savant de tous les riches*". But it does seem a strange anomaly that almost unlimited sums can be obtained for experiments on "Woolwich Infants", torpedoes, and the like—in short, anything that will most effectually destroy the human family on the largest scale; whereas there is untold difficulty in getting the most modest amount if the experiment be merely for the purpose of *saving* life. Of course I know it will be said that this is a trite complaint—one that has been made over and over again; also that upon the efficiency of our torpedoes and big guns depends our existence as a nation. Well, I can only answer that, if the complaint is trite, it is all the greater reason that the evil should be redressed; and that, if our exist-