treated, the majority without operation, and the total mortality was 80.5 per cent. Prior to the introduction of the Rammstedt operation in 1918, the operative mortality at this hospital was 100 per cent., not one single case having recovered after operation. In the following year, when the Rammstedt operation was adopted, it fell to 41 per cent. after surgical treatment, whilst in 1923 for 55 collected cases the operative mortality was 29 per cent.

During the past twelve months we have operated on 16 cases, with 5 deaths. This gives a death rate of approximately 30 per cent. Death occurred in 2 cases within twenty-four hours of operation, and was due to post-operative collapse in weakly and debilitated infants. In 3 cases it occurred four, seven, and twelve days later, and appeared to be due to gastro-intestinal disturbances and inability of the infant to assimilate its nourishment.

In cases with a long history it is not unusual to find the intestines small and shrunken when examined *post mortem*, and it appears that death is due to their atrophy from disuse and their failure to recover their digestive function. The average age in these cases was 7 weeks, and the duration of symptoms twenty-one days. The condition most commonly appears in the third week of life.

Whilst fully realizing that the figures here cited are too small to allow deductions to be drawn, they serve none the less to illustrate the average results of surgical treatment of the disease at the present time.

CONCLUSION.

We would conclude by again urging that, once the condition has been recognized, surgical aid should be sought at the earliest possible moment, for in operation lies the quickest and most certain chance of success.

After surgical treatment the infant usually shows rapid improvement and within a few weeks is frequently able to be discharged as cured. Medical treatment, on the other hand, may require months of careful attention, and the ultimate result is often uncertain. If recovery takes place after operation it appears to be permanent; in fact, the majority of those who recovered after operation in infancy are in later life a good deal above the normal weight for age.

A NEW TUBERCULIN.*

BY

ALEXANDER LUNDIE, B.Sc., M.B.Ed.,

D.C.M.S. (PATHOLOGY) MINISTRY OF PENSIONS, W. MIDLAND REGION.

ORGANIC analysis is not likely to yield any single substance capable of producing antibodies to the tubercle bacillus, and if it is carried too far it is not even likely to yield any mixture of substances capable of antibody formation. The most efficient antigen is likely to be a suspension of entire or of pulverized bacilli in normal saline. There are technical difficulties in the way of obtaining ideal products, and these difficulties are aggravated when the antigen is required for immunization experiments or for therapeutic use.

Besredka's antigen is prepared at a high temperature, and most tuberculins are made from bacilli killed by heat. The analytical processes employed to determine the composition of tubercle bacilli mostly involve the use of heat, to kill the organisms if for no other purpose. Heat is required for extracting the acid-fast material from the organisms. I know of no simple solvent that can extract this substance without the prolonged action of heat. A number of observers have noted that hydrochloric acid added to a simple fat solvent is able materially to assist the process of extraction.

If any chemical substance could be found to dissociate this acid-fast substance from its union with other constituents of the bacillus, at a comparatively low temperature and without loss of time, it would make the protein constituents of the bacillus more easily accessible to simplesolvents, and would yield them up in a state more nearly resembling the parent substance. Phosphorus trichloride seems to have

this action, but it also reacts with the protein constituents of the bacilli. Very definite considerations led to the use of this reagent for the purpose of preparing a tuberculous antigen directly from living organisms. It is not always possible to foretell exactly what will happen when a new reagent is tested. A number of reagents were previously tried, with more or less futile or even alarming results. The process adopted is offensive, and must be conducted with due precautions for safety. Phosphorus trichloride gives a colour reaction with all bacilli and with other organic substances containing protein. The colour is purplish-brown. Enough water must be present to act catalytically; too much will interfere with the reaction, and there will be no reaction with absolutely anhydrous materials. Under proper conditions violent effervescence takes place, and a voluminous evolution of irritating fumes occurs, which cause coughing and headache, which may last for several days.

Although the original mixture is liquid, the end of the reaction is indicated by its becoming a firm pasty mass, which can be liquefied by the addition of water. All the bacilli can be precipitated along with the phosphorized fats by the addition of acetone. This precipitate is the basis of the tuberculin about to be described. Its antigenic action is feeble, but is much enhanced by cholesterin.

Sound guinea-pigs bear relatively large doses of tuberculin, but tuberculous animals are readily killed in thirty-six to forty-eight hours by comparatively small doses. A healthy animal bore a large dose of this new preparation with impunity. A seriously diseased animal also bore the same dose, and even appeared to be better twelve hours afterwards. Twenty-four hours later a second dose, even larger than the first, was given, and borne equally well. Forty-eight hours after the first dose a third dose was given, and next day the animal was markedly better in health. Clearly this substance had not the lethal property of old tuberculin towards tuberculous animals. Its therapeutic action on the same animal was then tried. A weekly dose was thereafter administered until all the enlarged glands became impalpable, while the animal actually became the healthiest and heaviest in the establishment. When it was at last killed no trace of the diseased glands could be found, but there were a few scars of healed tubercles in the spleen, and an area of perisplenitis. An emulsion of the spleen injected into another guinea-pig produced no disease.

The various strains of human and bovine tubercle bacillus differ in virulence, so a culture of bovine bacilli of well established virulence was obtained from the National Collection of Type Cultures for the purpose of the next experiments, which were performed on rabbits.

The rabbit never survives an infection of bovine tuberculosis, no matter how contracted. If the bacilli be inoculated subcutaneously into one of the hind limbs an insignificant glandular enlargement soon arises. After the lapse of a period, depending upon the mass of the bacilli inoculated, general symptoms supervene. General miliary tuberculosis sets in and rapidly kills the animal. On *post-mortem* examination none of the lesions shows any evidence of circumscription. They tend everywhere to spread diffusely.

Five young rabbits were each inoculated intraperitoneally with 5,000 million virulent bovine tubercle bacilli. One was kept as a control without subsequent treatment. In six days its temperature had risen two degrees, and in eleven days four degrees. Its weight varied, reaching a maximum in eighteen days; it died in thirty days, weighing exactly the same as at the beginning.

The second rabbit was treated with the new tuberculin. It gained weight for twenty-one days. It was accidentally injured, and died on the twenty-second day.

The third rabbit became ill five days after inoculation. On the eleventh day its temperature was 105.3° F. Treatment was then commenced. It gained weight for thirty-one days, but the temperature remained high, and it died on the fortieth day.

The fourth rabbit became ill one day earlier than the others. Treatment was commenced on the tenth day, and it died on the thirty-fifth day.

The fifth rabbit became ill on the sixth day. Treatment was begun on the eleventh day, but in this case human, not bovine, tuberculin was used. All the other rabbits were treated with bovine tuberculin. This animal trebled its weight in fifty-five days. The local infection soon became sharply delimited from the surrounding healthy tissues and began to heal. The animal was killed at that stage on account of my holiday arrangements. *Post mortem* the lungs showed large, well defined tubercles in process of healing.

* A paper read before the Birmingham Branch of the British Medical Association, November 20th, 1924.

A pair of larger rabbits was infected by the subcutaneous instead of by the parenteral route. Both were decidedly out of sorts in seventeen days. Treatment was commenced on the twenty-sixth day. The first had already begun to decline seriously eight days before, owing to the infection having been intentionally accelerated by massage of the infected glands. Tuberculin of human origin was administered to both animals, but the first died thirty-five days after infection, without having made the slightest sign of rallying.

The second rabbit of this pair was seriously ill within sixteen days of infection, and treatment was commenced ten days later, and continued to the fifty-sixth day after the original infection. Only seven doses of tuberculin were given altogether. This animal increased in weight long after treatment was stopped, for nearly five and a half months. Eventually it was seized with nasal catarrh and signs of pneumonia. Nasal catarrh had never been observed in any of the other animals; it was probably a superadded infection by one of the pasteurilla group of organisms, and on this supposition the animal was promptly killed. *Post mortem* large healing tubercles were found in both lungs as well as fresh haemorrhagic patches. The site of infection was completely healed, but there was a large subscapular tuberculous gland, circumscribed by healthy tissue, though otherwise certainly capable of rekindling the infection.

This case and the last one of the other series were both decidedly encouraging, because rabbits at the best are difficult to manage under experimental conditions. Many die from accidental causes, and it says much for the care and skill of my assistants that the death rate of both rabbits and guinea-pigs is the lowest that I ever heard of. All these animals were very severely infected before treatment commenced, and they were all treated with very excessive doses of tuberculin. Not one reacted fatally to this excessive dosage.

Man has a much greater natural resistance to tuberculosis than the guinea-pig or rabbit, and his treatment is accordingly more satisfactory. Still, I have little fear about the results of my next animal experiments, if I think it necessary to resume them. As soon as it was thoroughly ascertained that the use of this tuberculin was perfectly safe for animals, I took advantage of the courtesy of Mr. Gemmill, and later of Mr. Naughton Dunn, who enabled me to treat cases of lupus vulgaris and of surgical tuberculosis. These cases responded very favourably.

Lupus erythematosus has responded to treatment even more rapidly than lupus vulgaris. But I consider an apparent cure not to be beyond danger of relapse, and would advise further treatment for eighteen months. I am indebted to Mr. Yates and Mr. Heath for cases of this disease.

I am indebted to Dr. Taylor of Syston, Leicester, for the first reports on the treatment of pulmonary tuberculosis with this tuberculin. I deliberately avoided treating this disease until I had studied the effect of treating the other forms of tuberculosis. Encouraged by Dr. Taylor's reports on advanced chronic cases, I undertook the treatment of a very advanced case of phthisis. I am indebted to Dr. Castellain for an introduction to this case, and to Dr. Garbutt for permission to attend and give the necessary treatment.

There were physical signs in both lungs and large cavities in the upper lobe of the right lung. The prospects were as hopeless as could be desired for the trial of a new method of treatment, and in that opinion we all concurred. The patient was aged 23, and was herself a good example of the "spes phthisica." She was quite sure she was recovering. She had three doses of the bovine tuberculin, followed by a course of autogenous tuberculin, beginning with a dose of a millionth of a gram. All anorexia rapidly ceased, the appetite became excellent, and sleep was restored; sweats ceased, sputum was reduced from about two pints a day to about a quarter of a pint. The pulse became regular and much slower (144 at first, afterwards 88), respiration was calm, tubercle bacilli were comparatively scanty in the sputum, and the physical signs greatly improved. The temperature became more equable, but was slower in coming under control than any other symptom. Flesh was put on quite obviously.

Later the patient had a complete pneumothorax of the right side. The temperature rose to nearly 103°, but it afterwards returned to normal, and she made some progress. Nineteen days after the occurrence of the accident the lung apparently collapsed again suddenly and death quickly ensued. She had been examined regularly all the time, and the lung was found to be expanding.

The principle I have enunciated is probably applicable to the treatment of leprosy and certain other diseases. The nastin treatment of leprosy by Professor Deycke of Constantinople is akin to it, but nastin is very toxic.

I am indebted to the Director-General of Medical Services, Ministry of Pensions, for permission to publish this paper.

PLEURAL REFLEX ASSOCIATED WITH FRACTURE OF THE TWELFTH RIB.

BY

GLADYS M. WAUCHOPE, M.D., M.R.C.S.,
HOVE.

THE following case seems worthy of record as a striking and unusual complication of fractured rib.

An unmarried woman, aged 41, fell from a stool on the night of July 15th, 1924, striking her back against a chest of drawers. There was an abrasion of the skin extending vertically upwards from a point below the level of the last rib about four inches from the right border of the spinal column. The twelfth rib was fractured about midway between its junction with the vertebra and the abrasion.

Strapping was applied, and the patient remained in bed, experiencing, for the first few days, considerable discomfort and pain even on slight movement. Four days after the accident she had an attack of hiccup lasting for some hours; both before and after this she had an occasional single hiccup. Progress was otherwise uneventful, and on July 28th the patient got out of bed for the first time. The next morning, when out of bed, she stooped for something and was suddenly conscious of a feeling of faintness, breathlessness, and palpitation, accompanied by severe pain in the upper part of the sternum. She felt very ill and thought that she was going to die. When seen an hour later she was pale, anxious, and dyspnoeic. The pulse (136 a minute) was feeble, and small, but regular. The heart was not enlarged. Harsh, rapid, and shallow breath sounds were heard all over the lungs. There were no areas of diminished resonance and no adventitious sounds. The temperature was 98° F. Adrenaline and cardiac stimulants were administered and the breathing improved; the general condition, however, remained serious during the day. In the evening the pulse rate was still 136, and the temperature had risen to 100.8° F. There were no physical signs in the chest and no pain at the site of fracture.

After this improvement was steady. The pain in the upper part of the sternum was continuous until the third day; it was not increased by breathing or movement. The temperature fell gradually, and was normal on July 31st (fourth day). On August 2nd the pulse rate was 72, the respiration rate 20, and the general condition good. The patient was allowed to get up. From about August 1st onwards, when all acute symptoms had subsided, involuntary sighs occurred from time to time, seeming to the patient to arise from the mid-sternal region. They continued two or three times a day for about a week, ceasing gradually. There was no further interruption to convalescence.

It seems probable that the sudden movement of stooping irritated the nerves of the pleura, either by tearing an adhesion or by friction, and so gave rise to a pleural reflex of the vasomotor type.

A consideration of the anatomy of the part shows that the pleura and diaphragm are in close proximity to the twelfth rib at the site of fracture, the external arcuate ligament of the diaphragm being attached to the tip and lower margin of the twelfth rib and to the transverse process of the first lumbar vertebra. The lung is above, reaching only to the eleventh rib, while the pleura extends to below the twelfth. In the normal subject the kidney would be liable to injury through the pleura and diaphragm, but in this patient it was movable and well below the level of the rib.

Pleural reflex or pleuro-pulmonary reflex has usually been recorded in connexion with operative thoracotomy. Stivelman¹ concluded that in over 75 per cent. of cases there was injury to the visceral pleura and underlying lung tissue. Capps,² however, has shown experimentally that the reflex can be occasioned by irritation of the pleura alone, the healthy pleura being much more tolerant than inflamed pleura. It is probable that, in the case reported, the pleura had been injured. The patient experienced more pain and discomfort than usually attend simple fractured rib, while the hiccup during the first week points to involvement of the diaphragm and therefore of the diaphragmatic pleura. The inflammatory reaction may well have persisted till the thirteenth day, when the cardio-respiratory symptoms occurred.

The sighing which arose during convalescence calls for some comment. Physiologists explain sighing as a deep inspiration following, and compensating for, a temporary cessation of breathing.³ Various causes which depress the respiratory centre may give rise to sighing—for example, cerebral anaemia and the excessive use of tobacco. It also occurs when the mind is deeply engaged, so that people unconsciously sigh at their work.

In the case under review a cause for suppression of respiration might be found in the injured pleura and