In a number of other cases (about six in all) treated with raw gland by the mouth the same results were obtained. One case (a woman) is worth referring to, as she was treated with both raw gland and insulin, and a comparison of the results obtained seems to indicate that she did better on the former than on the latter.

This patient commenced gland treatment in the spring of 1924 and did very well. In June of that year I described my gland treatment to one of the honorary physicians on the staff of one of the Sheffield hospitals, and on my suggestion he admitted her to his wards so that gland treatment might be carried out under test conditions. Unfortunately, before she was seen by the physician in question an over-zealous house-physician gave her an injection of insulin, and this treatment she has carried on at home since her discharge from hospital.

Comparing her condition now (on insulin) with what her condition was in the spring of last year (on gland) I find that she has lost weight, does not feel so well, and some sugar is to be found in the urine.

These facts encourage me to think that raw fresh gland by the mouth is a substitute for insulin, and that it is a very much better form of treatment, as it is perfectly safe and free from the grave risks that may attend even a small dose of insulin; furthermore, it is less irksome to the patient and less troublesome to the doctor in charge—for all who have had much experience of insulin will agree, I think, that insulin treatment is both troublesome and irksome and fraught with considerable risk to the patient.

It may be objected that the eating of raw gland is a loathsome thing; but patients on gland treatment inform me that it forms quite a delicious meal, especially when taken with lettuce—reminding us in this respect of phthisical patients on raw or very much underdone meat, who tell us how difficult they find it to return to cooked chops and steak, which seem tasteless to them after being used for so long to the more palatable and tasty fare of raw meat.

THE PREVENTIVE MEDICINE OF THE EAR.*

BY

SOMERVILLE HASTINGS, M.S.LOND., F.R.C.S.ENG.,
SURGEON TO THE EAR, NOSE, AND THROAT DEPARTMENT,
MIDDLESEX HOSPITAL.

It is stated that a learned lecturer on surgery used to preface his lecture on diseases of the ear somewhat as follows:

"There are two kinds of deafness: One, the curable, due to wax, which we treat ourselves, and the other, the incurable, due to some other cause, which we send on to the specialist."

There is a good deal of truth in this statement, and it is certainly of value to inquire to what extent deafness, not due to obstruction in the external ear, can be influenced by treatment. That acute middle-ear deafness can be cured, sometimes with complete return to normal hearing, is within the experience of everyone; but can any real improvement in hearing result when the deafness is due to the chronic disease of the middle or internal ear?

Let us take the middle-ear cases first. They divide themselves naturally into the suppurative and non-suppurative. Where there is chronic suppuration in the middle ear† this can in most cases be cleared up by treatment, but the perforation does not heal and the hearing but rarely improves, and not a few patients do not really thank one for drying up their ear, as they can hear better when it is moist. In middle-ear deafness without perforation of the drum, of the chronic catarrhal type and its many modifications, the outlook is, perhaps, a little more hopeful. In recent cases treatment of a septic focus in the nose, nasopharynx, or pharynx, or catheterization of the Eustachian tubes, may result in benefit to hearing, but the improvement is but rarely marked, while the results of treatment for internal-ear deafness are, I think, even more hopeless.

This does not mean, however, that we can do nothing for the sufferer, for though chronic deafness is very often

a progressive disease, it is also in many cases a disease that can be arrested. If we can induce a suppurating ear to dry up and cease to discharge and remain in this satisfactory condition, although no improvement in hearing may take place, any increase in deafness generally occurs very slowly. In cases of middle-ear deafness, also, without perforation or suppuration, much more can be done to arrest the disease than is frequently thought, though not to cure it. Some authorities are accustomed to label all cases of middle-ear deafness in which other members of the family have suffered in the same way, and where low pitched tuning-forks cannot be heard by air conduction, as oto-sclerosis, and, having made up their minds that oto-sclerosis is a progressive disease uninfluenced by treatment, to do nothing further for the patient. I do not for a moment deny the existence of the pathological changes described by Fraser, Gray, and others in connexion with cases of middle-ear deafness with the above symptoms, but I personally can recall only one case of middle-ear deafness in which the nose, nasopharynx, and ear drums appeared perfectly healthy and in which a clear history of previous trouble in these regions could not be obtained when carefully inquired for. A slowly progressive inflammatory process in the middle ear, started and maintained by some infection from the nose or nasopharynx, may apparently go on for years without giving rise to any appreciable deafness, and only when the region of the ossicular joints or the secondary membranes are involved may deafness come on. I feel, therefore, very strongly that even in cases where considerable deafness exists it is our duty to treat any septic focus that may be found, in the hope, not of appreciably improving the hearing, but of preventing further deterioration. Especially is the removal of nasal obstruction or sepsis likely to be of value in cases in which the hearing varies from day to day, or where immediate improvement in hearing follows the passage of a Eustachian catheter. I know quite well that there are some cases that nothing will touch, and have learned especially to fear patients with atrophic membranes, who hear better in a noise. A large proportion of these cases give a history of earache in childhood, and though, when seen, all infection of the nose and nasopharynx may have disappeared, I believe that a good many of them are the result of inflammatory trouble during the early years of life, and that in these cases, although the causative infection in the nose or nasopharynx has disappeared, the infective process still continues in the ears. If this type of case is excluded, my experience leads me to the conclusion that in a large proportion of cases of slowly advancing middle-ear deafness the disease may be arrested or its progress much retarded by dealing with a neighbouring septic focus. I can recall only one case in which I have had reason to regret operation for nasal obstruction in the case of middle-ear disease, but I can remember many in whom the history of deafness, both before and after operation, has made me wish I had operated sooner. In the much fewer cases of internal-ear deafness, where a cause can be assigned, it should, of course, if possible, be removed. I have thought that the progress of those cases due to sclerotic changes in the vessels has been retarded by the adoption of a purin-free diet and flushing out the system by drinking quantities of hot fluids, but it is difficult to be sure of this.

THE PREVENTION OF DEAFNESS.

If the foregoing conclusions be accepted it must of necessity follow that since little or nothing can be done by way of treatment for deafness, except to arrest the progress of the disease, the right thing is to get hold of cases early and deal with the sepsis or other condition producing the trouble; or, better still, to deal with the cause even before the effect is produced. So that the whole art of medicine, as far as the ear is concerned, is preventive—by either dealing with the hygiene of the mouth, ear, nose, or throat, before any changes in the ears occur, or, if this is not possible, directly there is the slightest indication of the commencement of mischief in the ear.

It is most important to remember that, with the exception of tumours, practically all diseases of the ear are due to

* Abridged from a lecture given before the Society of Medical Officers of Health.

† Dr. Kerr Love states that about 1 per cent. of the school children of Glasgow have chronic suppuration of the ear. (*Diseases of the Ear in School Children*, 1919.)

trouble elsewhere, and that in by far the greater number of cases the disease starts in the middle ear. Further, it must not be forgotten that in practically all cases of chronic middle-ear trouble, and in most acute cases, the ear disease is an extension of some chronic sepsis in an adjacent region, and even when acute otitis follows one of the exanthemata, that a septic focus is nearly always present. Hence, in middle-ear suppuration following scarlet fever, some authorities recommend the removal of adenoids as soon as the patient is well on the road to recovery from the disease. To operate for mastoid disease without searching for a septic focus before discharging the patient is as foolish as to open an appendix abscess without searching for the appendix.

The results of careful attention to general hygiene, and especially to that of the mouth, nose, and throat, in preventing deafness is very striking, and it is noticeable how large a proportion of the working classes over the age of 30, who for the most part have had the misfortune to lack this in the past, have some impairment of hearing. Sir George Newman, in his annual report for 1923 as Chief Medical Officer of the Board of Education, tells us that the percentage of children found on routine inspection in London with defective hearing has during the last ten years been halved, and that in Cambridge and Exeter the results have been even more striking; but he hastens to add that while this satisfactory result has been obtained in connexion with children of school age, the proportion of children going to school for the first time at the age of 5 or thereabouts with ear disease has hardly changed at all. This clearly points, as do many other facts, to the necessity of dealing with the beginnings of disease in children below school age, and that it is imperative that the commencement of the disease should be recognized and the first preventive measures taken in the infant welfare centres or the nursery schools. Mastoid disease calling for operation is by no means rare in babies of less than 12 months, and I am of opinion that otitis media is much more common in infants than is generally supposed. I have seen babies with red inflamed drums apparently quite well and happy and but little disturbed by the condition. When an infant or young child persistently rubs his ear, or frequently rubs the back of his head on his pillow, it should be taken as an indication of some ear disease and a careful examination carried out; if adenoids are present they should be removed. The removal of adenoids even in a baby a few days old is a very simple matter, if the proper sized curette is made use of, and it is remembered that, since it is impossible to insert the finger into the nasopharynx of a babe, the curette must serve for both diagnosis and treatment. No anaesthetic is needed, and most babies are quite prepared to enjoy a meal within a few minutes of the little operation, and appear little, if at all, inconvenienced by it. In children who are old enough to tell us when they have pain, two complaints of earache should be sufficient to determine a careful examination for adenoids,* and if they are present even in small quantity they should be removed. Whether the tonsils, unless obviously diseased, should be removed at the same time is a moot point, and all I would like to say here is that on several occasions in which I have removed adenoids only, the subsequent history has led me to regret that I did not remove the tonsils at the same time.

But we ought to be able to prevent ear disease without waiting for deafness, pain, or other ear symptoms. Babies who cannot take the breast or bottle properly usually have adenoids. These should be looked for and removed. Every child also with a running nose needs immediate treatment, for not only may this condition directly encourage the growth of adenoids, but also, possibly without the existence of adenoids at all, a chronic infection of the ear may result. In addition to general hygiene and a sufficiency of vitamin-containing food and fats, there are four points especially worth remembering in the prevention and cure of chronic rhinitis and its ear complications in children.

1. In visiting a number of nursery schools last year I was especially struck by the difference between the children in the open-air and indoor schools. Anything from 10 to

40 per cent. of the children shut in rooms, even though some windows were open, had running noses, whereas in the open-air schools I saw but one or two. It is not always understood that open-air schools need not be very cold even in winter if an efficient system of heating is installed.

2. As soon as children are old enough they must be taught to blow their noses at frequent intervals. They should not pinch the nose with a handkerchief, but should hold the handkerchief a few inches below the nose and blow into this, at the same time blocking one or other nostril by a finger pressed on to the side of the nose. It is wonderful how early a child will learn this if carefully taught.

3. A little ointment composed of 1 to 3 grains of menthol and 2 to 8 minims of oil of eucalyptus to the ounce of vaseline, applied to the interior of the nose at bedtime, is often a very considerable help in keeping the nose clear and preventing discharge.

4. Constant nose-breathing must be insisted upon. It is but little good to operate for tonsils and adenoids for the prevention or treatment of middle-ear deafness unless at the same time the child is taught to breathe constantly through the nose. I have seen cases in which tonsils and adenoids have been completely removed, but in which the habit of mouth-breathing still remained, and in which deafness steadily progressed until this habit was got rid of. To cure children of mouth-breathing deep breathing exercises are very helpful. In these the mouth is kept constantly closed, both during inspiration and expiration, and breathing is carried on entirely through the nose. Children are still taught to breathe in through the nose and out through the mouth by some teachers. This is exceedingly harmful, for the following reasons: In the first place, it is quite clear that no one can breathe in and out through the nose unless the nose is fairly clear—that is, free from excessive secretion—but it is quite easy to breathe in through the nose and out through the mouth with the nose full of mucus, as each inspiration draws this material further back until it reaches the pharynx and is swallowed by the child. Inspiration is followed by expiration, after which there is a very short period of rest. At the conclusion of expiration with the mouth open the child finds itself in a position in which a definite muscular effort (that is, shutting the mouth) is required before starting the next inspiration, if it is to be a correct one. There is a strong tendency for this muscular effort to be deferred, with the result that the child starts the new inspiration, as he finished the previous expiration, with the mouth open. It is essential, therefore, that all breathing exercises should be carried out with the mouth closed the whole time. They should be repeated frequently rather than carried out for a longer period at less frequent intervals. At night a piece of 1-inch rubber adhesive plaster about 2 inches long can be placed vertically between the upper and lower lips, or a piece of thin rubber sheeting, such as dentists use, 6 or 8 inches long by 2 or 2½ inches wide, can be placed over the lips and tied tightly at the back of the neck by tapes at the four corners. In many cases, however, the cure of mouth-breathing cannot be carried out except by constant supervision of the child by someone whose business it is to remind him to close his mouth whenever it is seen open.

PRACTICAL POINTS ON PELVIC PAIN.

BY

A. E. CHISHOLM, F.R.C.S. Ed.,

ASSISTANT, OBSTETRICAL AND GYNAECOLOGICAL DEPARTMENT, ROYAL INFIRMARY, DUNDEE.

At the December meeting of the Edinburgh Obstetrical Society Professor Ranken Lyle of Newcastle read a paper on pelvic pain in its relation to neurasthenia, which provoked certain thoughts in me which I here seek to set in order.

I gather that Professor Lyle finds many patients who complain of pelvic pain, and who may or may not be found to have some demonstrable pelvic abnormality—for example, a uterine displacement—but whose symptoms he ascribes to neurasthenia and not to anything abnormal found in the

* The examination of the nasopharynx of several hundred boys, aged from 9 to 16, by the mirror, has shown that some quantity of adenoids was present or had been removed in about 80 per cent.