examination; the latter only reveals the gross size of the uterus, which may be masked by the presence of a fibroid or an unduly thick wall.

We have used the method to determine the patency of a tube after salpingostomy for tubal mole (impossible by the other methods if the opposite tube is patent), and also in a case which proved to be an instance of a double uterus. The value of the method as an aid to diagnosis in other difficult cases is obvious.

We believe also that the demonstration of the passage of particles from the uterine cavity to the fimbriated extremity of the Fallopian tube throws some light on the origin of ascending infections of the genital tract, and perhaps also on that of endometrioma of the ovary and the deposit of fragments of exfoliating endometrium on the pelvic peritoneum and surface of the ovary.

In conclusion, we wish to express our thanks to Dr. Eden and Dr. Cuthbert Lockyer for permission to investigate cases in the hospital.

THE HEART IN CHRONIC TOXAEMIA.

BY

H. CLULOW NIXON, M.D. EDIN.,

PHYSICIAN, ROYAL MINERAL WATER HOSPITAL, BATH.

DURING the examination of patients suffering from infective arthritis and fibrositis, an attempt has been made to estimate the degree of involvement of the heart and arterial system in this form of chronic toxæmia. These types of cases can very frequently be traced to a local focus of infection, which most usually arises in or around the teeth, tonsils, or nasal accessory sinuses, but may later involve the colon, gall bladder, etc. Less frequently the source of infection is in the uro-genital tracts.

This paper is concerned principally with those streptococcal infections associated with cases classified as "rheumatic," or more properly "infective." No attempt has been made to include hearts poisoned—first, by acute diseases such as rheumatic fever, typhoid fever, dysentery, or malaria; secondly, by endogenous means, as, for example, excessive thyroid action or chronic kidney disease; thirdly, by an exogenous poison such as alcohol; or, lastly, by syphilis. Toxins may influence any property of heart muscle, altering its rhythm, excitability, conductivity, contractility, or tonicity. They act chiefly on those tissues which govern rhythm through stimulation of the accelerator nerves, causing sinus tachycardia, and also on the tonicity of the muscle; to a lesser extent they influence the excitability, and to a still smaller the conductivity and contractility. They may directly affect the tissue of the heart by their local action, or organisms in other parts of the body may affect the heart by their toxins. From the fact that they are set free within the body in varying quantities their effect on the heart will also vary in degree from time to time. Apparently in streptococcal infection of low virility extending over many years, the heart does not become profoundly poisoned as in Graves's disease or chronic nephritis, where myocardial degeneration is common.

That the heart has partaken of the general toxæmia may be established by physical examination.

1. *Palpation*.—Estimate first the position of the apex beat. In many cases it is found in the nipple line or outside it. Secondly, note its character. It is often diffused and rather feeble and slapping; if it is the latter, it has to be differentiated from mitral stenosis, where the left ventricle is unfilled, or from myocardial degeneration, where the left ventricle is weak. There is no left ventricular hypertrophy, except when aortic incompetence, increased blood pressure, or chronic Bright's disease exists as a complication.

2. *Percussion*.—By careful percussion the left border of the heart can be found to have moved outwards, measuring more than $3\frac{1}{2}$ in. from the mid-sternal line. Increase in transverse diameter may therefore amount to $1\frac{1}{2}$ in. to 2 in., denoting myocardial weakness. The poisoned muscle fibres having lost their tonicity, there is an increase of pressure from a larger flow of blood into the heart during diastole, and an incomplete emptying is the result; in this way dilatation of the left ventricle takes place.

3. *Auscultation*.—Particular attention is to be paid—

(a) To the character of the first sound at the apex. It may be muffled and distant, or it may be short and sharp, or actually it

may be loud. If it is short and sharp, it must be remembered that the muscle may be inflamed, fatigued, or degenerated.

(b) To the absence or presence of a murmur. In the early stages no murmur is audible; later, when dilatation takes place, a mitral systolic murmur can be heard—that is, a murmur of relative incompetence, due to stretching of the circular muscular fibres around the auriculo-ventricular valves. This murmur is soft, diminuendo in character, and is not conducted beyond the mid-axillary line. It is louder after inspiration and on lying down, disappearing on full expiration, cessation of respiration, and with exercise.

The toxic heart is a large heart with relative increase in the oblique diameter, and its long axis is horizontal. It is soft and atonic, with the transverse diameter measuring more when lying than standing. On examination by x rays it is seen to move up and down in the manner of a concertina, and its size, determined by percussion, can be verified. It is a heart of low efficiency. On standing up after resting, the normal difference of rate of 7 to 15 beats a minute is markedly increased. On exercise, the heart unduly accelerates, and returns slowly to normal; often several minutes elapse before it has regained its original rate; this should be accomplished in two minutes. During exercise the blood pressure sometimes fails to rise with the pulse rate, or even may actually fall.

SYMPTOMS.

Certain general symptoms which are associated with any form of toxæmia may be present—for example, loss of weight and of the feeling of well-being. The skin is often yellow, and may present a papular rash. There is often some mental depression and irritability of temper. Vasomotor and nervous instability is prominent, and hence tremulousness, precordial pain, palpitation, breathlessness, giddiness, exhaustion, faintness, cold feet and hands, numbness, local sweating and flushing, dermatographism, or myotonic irritability may appear. Morning anorexia, nausea associated with gastro-intestinal disturbance (for example, offensive stools and flatulence), intestinal stasis and chronic constipation, headache and derangement of the endocrine glands, and sleeplessness are sometimes present. Night symptoms are frequent.

Taking particularly those symptoms most often present the commonest are:

1. *Palpitation*.—Tachycardia often makes its appearance at an early stage of the infection, and its severity bears some relation to the degree of toxæmia. An increase of 10 to 50 beats a minute represents the usual extent of the heart's acceleration. Quite frequently the symptom passes unnoticed by the patient. It is a sinus tachycardia.

2. *Breathlessness* on even slight exertion, such as going upstairs or climbing a moderate incline, is a common complaint.

3. *Exhaustion* is a troublesome symptom which causes patients more discomfort than increased effort.

4. *Precordial Pain*.—Pain is referred to an area situated usually below the left nipple, and is commonest in women. It must be looked upon as a vasomotor neurosis, or secondary angina caused by the circulation of toxins which increase the excitability of the nervous system and sensitivity of the reflex arc, so that mild stimuli are sufficient to produce an exaggerated impression. The pain is more usual when the patient is resting and in the evening. The attack may be precipitated by emotion or other reflex cause. Hyperalgesic areas of skin and referred pain follow the usual paths as in primary angina.

The Condition of the Arterial System.—Although admittedly in many cases systolic blood pressure is found considerably raised, yet more frequently sphygmomanometric readings are distinctly lower than normal. In 43 cases observed the following were the results:

Systolic Blood Pressure.				Cases.				Cases.			
				Cases.							
Between 180 and 190	...	1		Between 130 and 140	...	9		Between 120 and 130	...	10	
" 170 " 180	...	4		" 120 " 130	...	10		" 110 " 120	...	8	
" 160 " 170	...	4		" 110 " 120	...	8		" 100 " 110	...	6	
" 150 " 160	...	0									
" 140 " 150	...	1									

William Russell has drawn attention to the fact that, irrespective of the pulse wave, arteries contract and relax, due to the fibres of the muscular coat being placed transversely. This variation in size indicates the degree of hypermyotonia and hypomyotonia, and is determined by

the nervous mechanism of the arterial wall, and the blood composition or blood content acting directly on the vessel wall. In the earliest stage of toxæmia the arterial system partakes of the general atony, and there is hypomyotonia; but as the blood impurity, being absorbed from the alimentary tract, continues to circulate, it acts as an irritant or stimulant to the arterial wall, causing hypermyotonia. It is to be noted, however, that apparently considerable time elapses before permanent hypermyotonia, with changes in the muscular coat, appears, for on taking the blood pressure in these cases it is found to be subnormal, though the artery is hypertonic to the touch. On palpating the brachial artery definite local thickenings are felt in the arterial wall, small portions being affected. This occurred in 20 per cent. of cases examined.

Relation to Cardiac Irregularities.—The variety of toxic heart now under consideration does not reveal much of particular interest. Sinus arrhythmia is often present, as is generally the case in tachycardia, but it has been noted where this is absent. Isolated premature contractions, both ventricular and auricular, are of common occurrence—if anything, more frequently the latter; this may possibly be due to the action of the toxins on the thinner, less vascular muscle of the auricle. Auricular fibrillation is rare. I can only record one case which could be definitely traced to severe pyorrhœa. Willius states that streptococcic infections are one of the causes of auricular flutter, resulting in local injury to the heart muscle. It was not met with in the present series. Sino-auricular block, more common in infections of an acute nature, has not been seen, neither has pulsus alternans, which Mackenzie noticed in temporary cardiac exhaustion. Minor degrees of partial heart-block have been observed.

Electro-cardiographic changes are of considerable importance and are an indication of the damage to the heart muscle.

1. Very short rest time or diastole (τ - p).
2. Diminution in value of the waves—that is, the amplitude of their excursions is lessened, and the α and τ waves are flattened. (The α wave shows the state of the ventricular muscle with regard to nutrition, and the τ wave the strength of contraction.)
3. Left-sided preponderance.
4. Inversion of τ (lead iii).
5. Complete inversion of lead iii is a sign of severe toxæmia.
6. Lengthening of the α - τ interval.

Willius states that the normal limit of the α - τ interval is between 0.24 and 0.28 sec. Accepting this, then conductivity is definitely delayed in certain instances. It is only reasonable to suppose that impulse transmission is as liable to modification as any other function of heart muscle in toxæmic processes, where all the other functions may be involved. Lengthening of the α - τ interval has not been noticed, but this may be explained by the fact that the auriculo-ventricular node is specialized tissue, and therefore behaves in an independent manner. It is particularly in lead iii that the signs of poisoned heart appear; α in this lead is chiefly a left ventricular event. Why should there be left-sided preponderance?

There is no sign of left ventricular hypertrophy in uncomplicated cases. It is argued that the arterial hypermyotonia, already mentioned, is responsible for this by creating increased resistance to the blood flow. The τ wave negativity is due to changes in the cardiac function affecting contraction preponderance resulting from functional myocardial fatigue, thereby altering electropotential.

CONCLUSIONS.

1. There appears to be a definite connexion between left-sided preponderance and a hypertonic state of the arterial wall. When by treatment the toxins are removed, and can no longer act as irritants to the wall of the artery, left-sided preponderance disappears.
2. Coincidentally τ wave negativity in lead iii depends on the cardiac atony, and it tends to assume its normal deflection as the toxæmia becomes less pronounced, the heart regaining its power of sustained partial contraction, in which stretching is resisted.
3. From the foregoing it would seem that the generally accepted view that inversion of τ in lead iii when it appears is a normal event is incorrect.

4. The heart responds quickly to treatment directed to the removal of the focal infection and to eliminative methods such as are carried out in spa treatment, even though the poisoning has been of long duration.

5. The importance of adequate treatment to the alimentary canal, particularly intestinal, cannot be overestimated. Lack of space prevents the publication of illustrative cases.

BIBLIOGRAPHY.

- Mackenzie: *Principles of Diagnosis and Treatment in Heart Affections*.
 Lewis: *Soldier's Heart and Effort Syndrome*.
 Cowan and Ritchie: *Diseases of the Heart*.
 Willius: *Clinical Electrocardiography*.
 S. Calvin Smith: *Heart Records*.
 Price: *Diseases of the Heart*.
 Brookbank: *Diagnosis and Treatment of Heart Disease*.
 Strickland Goodall: *Lectures*.
 Russell: *Arterio-sclerosis*, BRITISH MEDICAL JOURNAL, May 17th, 1924
 Pardee: *Clinical Aspects of the Electrocardiogram*.

SEVERE DERMATITIS FOLLOWING ULTRA-VIOLET LIGHT.

BY

HENRY MACCORMAC, C.B.E., M.D., F.R.C.P.,

AND

H. MORELAND MCCREA, O.B.E., M.D.,
LONDON.

THE effects of the exposure of the whole body to ultra-violet light are well known, and have now an accepted place in the treatment of disease, particularly lupus vulgaris. Ultra-violet light may also be used for its general tonic effect, for it has been observed to produce a feeling of well-being in those who have undergone a course of treatment with one of the forms of apparatus now on the market. It is not, however, so generally known that overexposure may be followed by undesirable, and sometimes even serious, results, as the case recorded below very well shows.

The patient whose case we relate was an unusually healthy man aged 72 years. He informed us that he was accustomed to take in the evening what he termed an artificial sun bath. The lamp used for this purpose was placed above the bed, and as a rule an exposure of about ten minutes allowed for a single application. On October 10th, 1924, while thus engaged, he happened to fall asleep, with the consequence that the lamp continued to play over him for one hour and ten minutes. The immediate effect was a severe burning of the skin similar to that seen after ordinary sunburn, the front of the body and legs becoming at first bright red and swollen, subsequently desquamating. We were surprised to find that the back, a region that was not, so far as we can ascertain, directly influenced by the light, also became affected in a similar but slighter fashion.

Then an interval of about seven days elapsed, when further and more serious effects developed. The body, both back and front, was at this stage markedly pigmented, the right leg was swollen with in places raw areas exposing the papillae, and a similar but less pronounced condition was observed over the left leg. On October 22nd a general erythema of the trunk and extremities was noted, with urticaria, and the patient was very much disturbed by the severity of the itching, which seriously interfered with rest and sleep. Three days later, although in the interval the oedema had diminished considerably, there was an extension of the eruption to the face, which became puffy and wept in places.

So far the general condition had given no cause for anxiety, but on October 27th a marked change occurred, possibly as the result of the absorption of toxic substances from the damaged skin. There was restlessness, disturbance, and little and fitful sleep; the cardiac rhythm became disordered, extra-systoles developing. The patient continued to be gravely ill for some four days, then rallied and appeared to be out of immediate danger.

Although this satisfactory state of affairs can be recorded as regards the general condition, the integument continued to be extensively affected, the ears weeping, the face peeling freely, while the whole scalp exhibited a branny desquamation. The back was also desquamating slightly, and the skin there was still considerably infiltrated: the nails were striated and rough. The appearance of the skin of the