or it has been the presence of toxins in the exudate acting upon leucocytes therein and converting them into fibroblasts.

In my visit to Harvard University a little over a year ago I was extremely interested to learn from Dr. Drinker of some experiments he had been carrying out in regard to pulmonary fibrosis. It had occurred to Drinker that since fibrotic changes in the lungs are the result of inhaled dust having penetrated the alveolar wall and induced irritation of the interstitial tissue, possibly a similar chain of events might take place if particles of insoluble dust were injected into veins not far distant from the heart. Accordingly Drinker injected into the veins of animals extremely minute particles of manganese silicate, and he found that structural alterations were established in the lungs. Since my conversation with Dr. Drinker I have carried out a series of experiments on the Harvard lines, but instead of using manganese silicate I employed finely powdered bauxite, an aluminium ore. This was conveyed into the auricular veins of rabbits, and after a series of injections conducted over a few months we found in the microscopical sections of the lungs, kindly prepared for me by Mr. John Secker of the Physiology Department, large phagocytic cells lying within the alveoli and laden with dust particles, similar cells surrounded by mucus passing outwards by the small bronchi, and the same type of cell in the spleen. Until the series of experiments which I am at present carrying out is concluded, it is somewhat difficult to explain the presence and escape of the phagocytic cells alluded to. They are not leucocytes in the ordinary acceptance of the term. When an injection of finely divided particulate material is made into the veins of an animal not far from the heart the particles are carried along in the blood stream, and as the pulmonary capillaries are those through which, after leaving the heart, they would primarily pass, several of the particles are probably taken up by the endothelial cells. From the size and shape of the phagocytes found in the pulmonary interstitia and alveoli, also in the minute bronchi, the probability is that they are endothelial cells which have broken off from the pulmonary capillaries, and that they were fulfilling their function of eliminating foreign material from the blood. Their presence in the splenic pulp is similarly explained.

PROLONGED MENINGOCOCCAEMIA WITH TERMINAL MENINGITIS.

BY

G. LOVELL GULLAND, C.M.G., M.D.,

PRESIDENT OF THE ROYAL COLLEGE OF PHYSICIANS OF EDINBURGH;
PROFESSOR OF MEDICINE AND CLINICAL MEDICINE IN THE
UNIVERSITY OF EDINBURGH;

AND

W. R. LOGAN, M.D., F.R.C.P.E.,

CLINICAL PATHOLOGIST TO THE ROYAL INFIRMARY, EDINBURGH.

THE following record emphasizes the importance of searching for the meningococcus in obscure cases of pyrexia, particularly of an intermittent type.

The patient was a man of 57, an engine fitter, who had been unusually healthy. For thirty-four years he had not lost a day's work; he lived comfortably, and was a teetotaler. About the beginning of May, 1924, he had an attack of so-called "influenza." It commenced abruptly with shivering and fever, but he had no headache or other pain, and no catarrh or pulmonary symptoms of any kind. He had some vomiting, and in a day or two slight diarrhoea came on, which lasted only a few days. Weakness was really the principal symptom. The fever was not continuous, but he had rigors at odd times, apparently almost daily. He was in bed for eight or nine weeks. He was up for about a fortnight before he was admitted to Ward 32, but could hardly walk because of weakness, and he had lost nearly 3 st. in weight. He came into the infirmary on July 8th, 1924.

His only symptoms were weakness and loss of appetite; the appetite returned after a few days in hospital. Physical examination gave entirely negative results, and need not be detailed. All the systems were normal for a man of his age, and remained so throughout. The spleen was not enlarged at any time. For the first two days the temperature remained

normal; on the third day it went up with a slight rigor, but only to 100° F., the next day to 101°, and on that day his white count was found to be 10,400. Throughout the illness this leucocytosis with polymorph increase persisted; the highest figure reached was 29,100, a month later.

From this time onwards there was, in the way of symptoms, nothing but increasing weakness. The temperature followed a most erratic course; on some days, and sometimes for two or three days together, there would be only a slight rise, say to 100° F., on others, coming singly or in small groups, rigors, when the temperature ran up for a short time to, say, 103°. These occurred at any hour of the day or night; between times the temperature was generally subnormal. From August 26th to September 9th the rigors occurred daily, and were more severe; the temperature was higher, reaching 104° or over on most days. The pulse rate varied strictly with the temperature: when it was subnormal the pulse was in the 60's, when it reached 104° it often ran up over 120. In the same way the respiration rate varied from 20 to 32.

Throughout this period everything we could think of was investigated in the hope of getting some light. The chest was x-rayed, with negative results; the urine was examined daily and was always normal. Though the persistent leucocytosis made enteric or malaria unlikely, the blood was subjected to the Widal test for the usual organisms, with negative results, and examined constantly for plasmodia, equally unsuccessfully. The spirochaete of rat-bite fever was sought in vain; the patient denied ever having been bitten by any animal, and it was only about ten days before his death that his wife remembered that about the middle of April he had been badly scratched on the back of the hand by a "wild" cat that lived by rat-catching. This led to several intravenous injections of novarsenobillon, without improvement. The Wassermann test was negative in both blood and cerebro-spinal fluid, and the latter showed no increase of cells and no organisms on culture. The first blood culture, in July, gave a growth of staphylococci and Gram-negative diplococci. As the staphylococci were almost certainly a contamination, it was thought the diplococci might be so also; but the observation was repeated a few days later and they were found alone, and again on August 12th. A blood culture was not made again till September 8th, and then no organism of any kind was found. Cultures were taken from the urethra, but no diplococci were found.

About September 9th meningeal symptoms began to develop—headache, head retraction, drowsiness, and incontinence of urine and faeces. The temperature was more continuously raised, seldom dropping to normal, but never running above 101°; usually it was about 100°. The rigors ceased. The patient gradually became weaker, emaciated very rapidly, grew more drowsy, and died suddenly on September 16th.

The first lumbar puncture to yield positive results was made on September 11th. The fluid was definitely under pressure, was turbid, and microscopically showed many polymorphs and large lymphocytes. In some parts of the films diplococci were found, practically all intracellular, and always in the large lymphocytes, never in the polymorphs. The same diplococcus which had been found in the blood was grown from this fluid. As lumbar puncture gave relief it was repeated daily. On the 12th the cell count was 5,800 per cubic millimetre. In the later punctures the fluid became clearer.

Post mortem a certain amount of bronchitis and hypostatic congestion of lungs were found. The heart was practically normal; very slight atheroma in the aorta; liver normal, except for some cloudy swelling and perhaps fatty degeneration; spleen not enlarged, apparently normal; cortex and medulla of both kidneys slightly narrowed; other organs all normal except the brain. The pia arachnoid showed areas of localized thickening in various parts, chiefly towards the base and over the cerebellum. These were evidently patches of meningitis, and on section microscopic examination confirmed this view. There was a marked exudate, with numerous polymorphs; slight excess of cerebro-spinal fluid, with slight dilatation of the ventricles internally, but the brain substance was normal.

Cultures made *post mortem* from the heart blood, spleen, and liver remained sterile, but the organism was obtained from the exudate at the base of the brain.

The Micro-organism.

The diplococcus, when first obtained from the blood stream, resembled a meningococcus in its cultural characters, save that it grew more readily on ordinary media than do most strains of meningococcus. On plain agar slopes it grew well, even when inoculation was not particularly heavy, and when no accessory substances could have been carried over from the original medium. Growth on glucose ascitic fluid agar (Thomson's) was profuse. After subculture for some time it lost this readiness of growth to some extent, and finally became indistinguishable culturally from type strains of meningococci, so far as readiness of growth on various media was concerned. Morphologically, it was indistinguishable from the meningococcus or gonococcus.

Its fermentation reactions were those of the meningococcus; it produced acid in glucose and maltose, no change in lactose, saccharose, and salicin. The gonococcus does not ferment maltose,

while *Micrococcus catarrhalis* ferments neither maltose nor glucose.

A rabbit and guinea-pig inoculated with cerebro-spinal fluid containing the organism showed no effect, though the guinea-pig died about a fortnight later. No pathological lesions were discovered, and cultures made from its brain and organs did not yield the organism in question.

Guinea-pigs inoculated with the patient's whole blood, with the plasma, and with cultures of the organism yielded similar negative results.

No agglutinating serums for meningococci could be obtained from various sources tried. A commercial polyvalent antimeningococcal serum did not agglutinate the organism even in low dilutions, but as it did not agglutinate the type strains of meningococcus I, II, III, and IV either, this observation was obviously not of much value.

A beginning was therefore made in the immunization of two rabbits with the diplococcus from the patient. The organism was first found in the patient's cerebro-spinal fluid on September 11th. It was present in films of the fluid only in small numbers, but growth was obtained, although with difficulty. As obtained from the spinal fluid, it did not grow nearly so profusely as it had when obtained from the blood stream. It behaved now, in short, as a meningococcus, and there was now no hesitation in labelling it as such. The patient died before the agglutinating serum from the rabbits was ready.

Four type strains of meningococcus (Lister Institute Collection) were tested with the serum along with the organism under examination (the "Murray coccus"). The emulsions were made in saline from growths on Thomson's medium, and the macroscopic method in a water-bath at 55° C. was used. Readings were made at intervals up to four hours. Both serum and saline controls were used. The test was not carried beyond 1 in 64 with Type III meningococcus.

Identification of Type of Meningococcus.

	Murray coccus serum: Dilutions, one in—									Controls.
	16	32	64	128	256	512	1,024	2,048	4,096	
Murray diplo-coccus	+++	+++	+++	++	++	++	+	+	Trace	—
Meningococcus Type I	±	Trace	—	—	—	—	—	—	—	—
Meningococcus Type II	+++	+++	++	++	+	±	±	Trace	—	—
Meningococcus Type III	++	±	Trace	—	—	—	—	—	—	—
Meningococcus Type IV	++	±	±	Trace	—	—	—	—	—	—

+++ represents complete clearing with sedimentation, and the other signs correspondingly smaller degrees of agglutination.

The organism could therefore be definitely labelled meningococcus Type II of Gordon's types.

Dock¹ has recently analysed sixty-eight reported cases of meningococcaemia, occurring in different parts of the world. In the case which he studied personally the organism was not agglutinated by the patient's serum or by polyvalent antimeningococcus serum at the second or third subculture, but at a later date was agglutinable in low dilutions both by antimeningococcus and antigenococcus serums. It was at first thought to be a gonococcus, but ultimately determined to be a meningococcus. The intermittent character of the fever, often with a definite malaria-like periodicity, was a salient feature of these cases.

Rolleston² also drew attention in his Lumleian Lectures on cerebro-spinal fever to these cases of meningococcaemia, which "may precede, follow, or occur without meningeal infection." He quotes some interesting observations by Netter and Dopfer. Dopfer's observation is that whereas before the war 96 per cent. of the cases of meningococcal meningitis were infected with the meningococcus Type A (Gordon's I and III types) a change has taken place, so that now the parameningococcus or Type B (Gordon's II and IV) is responsible for an equal, if not a larger, number of the cases. Netter³ comments that this observation may be correlated with an increased incidence of meningococcaemia, and meningococcal manifestations in the skin, joints, and eyes.

The organism in our case, a meningococcus of Type II (British classification) or Type B (French classification) certainly showed no particular affinity for the cerebro-spinal system, as it was more than four months before the

spinal fluid became infected. It is interesting that the organism disappeared from the blood stream before this terminal meningeal infection took place.

We shall not here embark on a discussion of the mutability or fixity to type of these bacteria beyond noting that the organism as obtained from the blood stream was a hardier, coarser-growing type than that obtained later from the cerebro-spinal fluid. Such a change might, of course, be merely a temporary or fluctuating variation due to environmental conditions, particularly as the hardier qualities were quickly lost on subculture.

The case emphasizes the necessity for bearing in mind the possibility of a change in the nature of meningococcal infections.

REFERENCES.

- ¹ Dock, William: Intermittent Fever of Seven Months' Duration due to Meningococcaemia, *Journ. Amer. Med. Assoc.*, vol. lxxxiii, No. 1, July 5th, 1924.
- ² Rolleston, Sir Humphry: Lumleian Lectures on Cerebro-spinal Fever, *Lancet*, April 5th and 12th, 1919.
- ³ Netter: *Bull. et Mém. Soc. Méd. des Hôp. de Paris*, 3e série, vol. xli, 883, 1917.

THE CHEMOTHERAPY OF RECENT EXPERIMENTAL WOUND INFECTIONS:

WITH SPECIAL REFERENCE TO THE ACTION OF
ACRIFLAVINE.*

BY

C. H. BROWNING AND R. GULBRANSEN.

(From the Pathological Department of the University and the Western Infirmary, Glasgow.)

THE question whether localized infections can or cannot be treated effectively by means of antiseptics applied to the infected area appears still to be a subject for argument. Animal experiment might be expected to yield a decisive answer. But a difficulty, as most workers have found (for example, see Watson Cheyne, Bassett-Smith, and Edmunds⁶), is the tendency for infections in small animals to become generalized and so to escape beyond the range of action of the antiseptic. In the case of mice inoculated intraperitoneally with organisms so highly invasive as virulent pneumococci, we² have shown that it is possible to effect cure sometimes by means of intraperitoneal injections of various antiseptics; but this line of work has not been pursued owing to the difficulty in controlling the virulence of the infection. Recently, the infection of experimental wounds in guinea-pigs with a virulent culture of *B. diphtheriae* has been recommended as a test object for investigating the therapeutic properties of antiseptics (Braun,¹ Feiler⁴). In the first place, there is very little tendency for the infection to become generalized; and secondly, when a highly virulent culture is employed the results are decisive, since the unmodified infection is rapidly fatal.

METHODS.

Guinea-pigs of about 200 to 250 grams weight were used. An area of the skin of the abdomen was shaved. When the animal was under the influence of a general anaesthetic two parallel linear cuts, each measuring about 1.75 cm. and at a distance apart of 0.8 cm., were made through the skin in the direction of the long axis of the body, care being taken not to incise the deeper tissues. Then, by means of a stout wire bent into a loop, the strip of skin between the cuts and also the skin for about 1 cm. on either side of the incisions was raised from the deeper structures, so that pockets were produced; there was, of course, very little bleeding from the wound. About a quarter of a well grown forty-eight hours' agar slope culture of *B. diphtheriae* was then rubbed into the wound and the pockets. The animals were left for varying periods and were then treated. Treatment consisted in placing the animal on its back and pouring slowly into the wound 1 c.cm. of a watery solution of the substance the action of which was to be investigated, the wound and pockets being meanwhile swabbed out with a small

* Work done with the support of the Medical Research Council.