content of blood serum or plasma. It has numerous advantages over other tests which have subsequently been proposed: (a) its extreme delicacy; (b) is not given by other yellow substances which may colour the plasma; (c) gives important qualitative distinctions between certain forms of icterus; (d) can now be used for a satisfactory quantitative estimation of bilirubin.

2. The qualitative reaction may enable a positive diagnosis of obstructive or of haemolytic icterus to be made. Pernicious anaemia, except in the stage of rapid temporary improvement, can be distinguished from secondary anaemias. In some diseases (for example, catarrhal jaundice, subacute liver atrophy, cardiac failure) the reaction may pass through all the stages from completely delayed to prompt direct, as the disease progresses. As a rule, however, in the commonest type of jaundice (the toxic and infective hepatic group) no information of any diagnostic or prognostic value is given to the clinician.

3. The test is of great practical value for the detection of "latent" jaundice.

4. Various alterations in the original technique have improved and simplified its use in clinical medicine.

BIBLIOGRAPHY.

- Adler, E., and Strauss, L.: *Klin. Woch.*, 1922, xli, 2285.
 Andrewes, C. H.: *Lancet*, 1924, i, 590.
 Idem: *Quart. Journ. Med.*, 1924, xviii, 19.
 Broun, G. O., Ames, O., Warren, S., and Peabody, F. W.: *Journ. Clin. Invest.*, 1925, i, 295.
 Brulé, M., Garban, H., and Weissmann, C.: *Presse Médicale*, 1922, xxx, 986.
 Feigl, J., and Querner, E.: *Zeit. f. d. gesamte exp. Med.*, 1919, ix, 153.
 Fishberg, A. M.: *Journ. Amer. Med. Assoc.*, 1923, lxxx, 1516.
 Fouchet, A.: *Journ. de Pharmacie et de Chimie*, 1918, xvii, 44.
 Gerrard, W. I.: *BRITISH MEDICAL JOURNAL*, 1924, ii, 224.
 Hijmans van den Bergh, A. A.: *Der Gallenfarbstoff im Blute*, Leiden, 1918.
 Idem: *BRITISH MEDICAL JOURNAL*, 1924, ii, 498.
 Hughes, T. A.: *Indian Journ. Med. Research*, 1924, xii, 403.
 Lephne, G.: *Deut. Arch. f. klin. Med.*, 1920, cxxxii, 96.
 Idem, *ibid.*, 1921, cxxxv, 79.
 McNee, J. W.: *BRITISH MEDICAL JOURNAL*, 1922, i, 716.
 Idem: *Quart. Journ. Med.*, 1923, xvi, 390.
 Meulengracht, E.: *Deut. Arch. f. klin. Med.*, 1921, cxxxvii, 38.
 Rosenthal, F., and Holzer, P.: *Ibid.*, 1921, cxxxv, 257.
 Schamberg, J. F., and Brown, H.: *Journ. Amer. Med. Assoc.*, 1924, lxxxii, 1911.
 Schiff, E., and Eliasberg, H.: *Klin. Woch.*, 1922, xxxviii, 1891.
 Strauss, L., and Buerkmann, W.: *Ibid.*, 1922, xxviii, 1407.
 Thannhauser, J. S., and Andersen, E.: *Deut. Arch. f. klin. Med.*, 1921, cxxxvii, 179.

RADIOLOGY OF THE GALL BLADDER BY GRAHAM'S METHOD.*

BY

HENRY COHEN, M.D.,

BRIGHT MEMORIAL RESEARCH FELLOW; HONORARY ASSISTANT PHYSICIAN,
LIVERPOOL ROYAL INFIRMARY;

AND

R. E. ROBERTS, M.D., B.Sc., D.M.R.E.,

HONORARY RADIOLOGIST, LIVERPOOL ROYAL INFIRMARY.

It is generally admitted that the routine examination by x rays of the gall-bladder area for the presence of stones is subject to serious limitations as regards diagnostic value, for even by means of good radiographs and skilful interpretation their existence can only be clearly demonstrated in, at the outside, 35 to 40 per cent. of the cases where they are present. Though an x-ray finding if positive is of definite diagnostic value, a negative finding does not exclude the presence of gall stones and must be ignored.

In the interpretation of shadows found in the gall-bladder area there are two types sufficiently characteristic to be practically diagnostic of gall stones. These are the annular or ring-like shadow (Fig. 1) and the cluster of shadows resembling a bunch of grapes (Fig. 2). The finding of these shadows in a radiograph of the right upper abdomen is practically certain evidence of the presence of gall stones; especially if, by radiographs taken in different positions of the body, they are shown to be situated nearer the front of the abdomen than the back. Other atypical shadows may be found in the region of the gall bladder which call for more serious thought and skill in interpretation before they can be taken as evidence of gall stones.

A large number of gall stones, however, are not sufficiently opaque to x rays to throw any shadow at all when they are in the body. Some, indeed, are so translucent that even when the gall bladder containing them is removed and radiographed they are no more opaque than the bile or the walls of the gall bladder by which they are surrounded; in

fact, they may be even less opaque, and may show in the radiograph of the free gall bladder as translucent areas in the shadow of the more opaque gall bladder and bile.

In 1913 Holland and Williams investigated the factors underlying this varying degree of opacity of gall stones. They found that the opacity depended on the proportion of the pigment and other salts of calcium. Those containing a large proportion of cholesterol were the most translucent, while those containing a large proportion of pigment or other salts of calcium were the most opaque. The calcium salts are often deposited at the periphery of the stone and so give rise to the characteristic annular shadow.

Apart from the direct x-ray evidence of gall stones, the existence of a pathological condition of the gall bladder may often be suspected from abnormalities of contour, position, or function, of the stomach or duodenum, found during the examination of the gastro-intestinal tract by the opaque meal; or by a process of elimination, by means of x rays, of gastric, duodenal, and appendicular lesions, and of urinary stones.

GRAHAM'S METHOD OF EXAMINATION.

The method of radiologically investigating the gall bladder, observations on which we wish to record in this paper, is attributable to Graham of St. Louis, U.S.A. It is based on the use of a salt opaque to x rays which, when injected into the blood stream, is excreted by the liver, enters the gall bladder, and is there concentrated, thus rendering the gall bladder under normal conditions opaque to x rays.

Rowntree has shown that certain dyes—for example, phenol-tetrachlorophthalein—when injected into the blood stream, are excreted almost entirely by the liver. The rate of disappearance of this dye from the blood

stream has been used as a test of liver function. Graham and Cole showed that tetrabrom-phenolphthalein salts—of both calcium and sodium—were excreted by the liver almost entirely and were opaque to x rays. The iodine compound, although more opaque, was, in their opinion, too toxic for general use. More recently Whitaker and Milliken, after a careful clinical and

FIG. 1.—Two gall stones, showing opaque periphery and nucleus, the intermediate portion being more translucent. Annular type of shadow.

* Read at a meeting of the Liverpool Medical Institution.

FIG. 2.—Gall bladder containing a large number of small opaque gall stones. Cluster type of shadow.

FIG. 3.—Collection of many small opaque stones in the fundus of the gall bladder causing a hemispherical shadow to the outer side of the duodenum. This was an accidental discovery during an opaque meal examination of the stomach and duodenum in the erect position.

FIG. 4.—Normal gall bladder seen eight hours after intravenous injection of sodium salt of tetrabrom-phenolphthalein (Graham's method).

FIG. 5.—Same case as Fig. 4 twenty-four hours after injection. Shadow of gall bladder is smaller but more opaque than in Fig. 4.

experimental comparison of the bromine and iodine salts, state their preference for the latter. In our investigations the sodium salt of tetrabrom-phenolphthalein has been used throughout. It was obtained as a crystalline powder from the Mallinckrodt Chemical Works of St. Louis.

Method followed in this Inquiry.

Our procedure was as follows:

1. All cases were first examined radiologically by the ordinary direct method for gall stones. In every case of our series which was operated on, except one, where a doubtful and non-diagnostic shadow was seen in the region of the gall bladder, the result of this examination was entirely negative.

2. The cases were then examined radiologically by Graham's method of intravenous injection of the dye. The technique adopted was as follows:

No special preliminary preparation was made except that during the four hours preceding the injection no food or fluid was allowed.

Five grams of the salt were dissolved in 40 c.cm. of distilled water and sterilized by boiling for twenty to thirty minutes. This solution was very slowly injected intravenously in two doses, the second being given half an hour after the first. In view of the danger of necrosis which might follow extravasation of the dye into the subcutaneous tissues, the needle was inserted separately into the median basilic vein and blood was allowed to flow freely from it before attaching the syringe. It is most important that the rate of injection should be slow if untoward symptoms are to be avoided. The injection of the dye was followed by a few cubic centimetres of sterile normal saline solution to avoid leakage of the salt when the needle was removed.

During the first day no food other than a glass of milk at lunch and in the evening was allowed. During the second day ordinary fish diet was given. For the forty-eight hours following the injection 40 grains of sodium bicarbonate were given every three hours during the day.

Two and a half hours after the injection (that is, one and a half hours before the first radiographic examination) 0.5 c.cm. of pituitrin was given hypodermically, followed three-quarters of an hour later by a simple enema.

Radiographs of the gall-bladder area were taken four, eight, twenty-four, and thirty-six hours after the injection.

Reaction following Injection.

The injections occasionally gave rise to unpleasant symptoms of vasomotor shock—such as pains in the back, giddiness, nausea, vomiting, cyanosis, coldness, or "queer feelings"—associated with a lowering of the blood pressure; a preliminary rise was sometimes observed.

We have encountered no serious symptoms in our cases; only a slight reaction occurred in most of them; serious symptoms can be avoided by slow injection. If necessary a hypodermic injection of 10 minims of a 1 in 1,000 solu-

tion of adrenaline hydrochloride may be given; we have never had to use this; occasionally we have given brandy after the injection.

The X-ray Appearances of the Normal Gall Bladder after the Injection.

The shadow of the gall bladder containing the dye, which has been excreted by the liver, can be made out four to five hours after the injection has been given.

1. The shadow increases in *intensity*, owing to the progressive concentration of the dye, up to anything between eight and twenty-four hours. The shadow is homogeneous, oval or pyriform in shape, and of even contour. After this the shadow becomes less intense as the dye becomes diluted and passes out.

2. The gall bladder shadow also varies in *size*, being larger at the earlier examinations (five to eight hours), and then becomes gradually smaller. This variation in size is important evidence of normal distensibility and contractility of the gall bladder.

Physiological Requirements.

Before the shadow of the gall bladder containing the dye can be produced three requirements must be fulfilled—namely:

(1) *The dye must be excreted by the liver.* This will not take place if there be, for example, gross cirrhosis or carcinoma of the liver.

(2) *The dye must be able to pass into the gall bladder.* This will not take place if there be obstruction to the cystic duct or common bile duct, such as might be caused by gall stones, adhesions, papilloma, thickened bile, catarrh, etc.; or, if the gall bladder be filled with gall stones or growth.

(3) *The dye must become concentrated in the gall bladder,* by absorption from the gall-bladder mucosa, contraction of the gall bladder, and retention of the dye for a sufficient length of time. This will not take place if there be chronic cholecystitis or if there be a fistula connecting the gall bladder directly with the gastro-intestinal tract and so allowing the immediate or early escape of the dye from the gall bladder.

The Abnormal Gall Bladder.

If there is any obstruction due to gall stones, adhesions, thickened bile, etc., we may find either that the gall bladder fails to fill at all with the dye and so gives no x-ray shadow; or that it fills scantily, giving a persistently faint shadow; or that there is delayed filling.

Another evidence of pathology is marked delay of emptying, or an unvarying size of the shadow, due to loss of elasticity of the gall-bladder walls. Or there may be a

TABLE I.—Showing Results of Investigation by Graham's Method of Ten Cases who were subsequently Operated on.

Case.	X-ray Findings.								Deduction.	Operative Findings.
	Gall-bladder Shadow after Injection.									
	Intensity.				Size.					
	4 hrs. 2	8 hrs. 4	24 hrs. 2	35 hrs. —	4 hrs. 4	8 hrs. 3	24 hrs. 1	35 hrs. —		
W. D.									Normal gall bladder	Normal gall bladder (gastric ulcer).
M. J.	1	3	4	—	4	4	3	—	Normal gall bladder	Normal gall bladder (pathological appendix).
J. O.	—	—	—	—	—	—	—	—	Pathological gall bladder	? Normal gall bladder. ? Slender adhesion at neck of gall bladder.
R. C.	—	—	—	—	—	—	—	—	Pathological gall bladder	Small pathological gall bladder. Stone impacted in cystic duct.
M. C.	—	—	—	—	—	—	—	—	Pathological gall bladder	Gall bladder full of stones. Cystic duct completely occluded.
P. W.	—	—	—	—	—	—	—	—	Pathological gall bladder	Many stones in gall bladder. Gall bladder adherent to duodenum with fistulous connexion.
M. W.	—	—	—	—	—	—	—	—	Pathological gall bladder	Small thickened gall bladder with one large stone almost filling it.
M. M.	1	1	—	—	4	4	—	—	Doubtful	One small stone in gall bladder. No evidence of obstruction of cystic duct.
M. F.	—	—	—	—	—	—	—	—	Pathological gall bladder	Healed duodenal ulcer. Dense adhesions around neck of gall bladder.
H. H.	2	3	3	3	3	2	3	2	Pathological gall bladder	Pancreatitis and subacute cholecystitis. S-shaped twist in neck of gall bladder.

Note.—In this and the following table (Table II) the numbers 1, 2, 3, 4, have been employed to denote different degrees of intensity and size of the gall-bladder shadow. As regards intensity, 4 represents the maximum intensity of the gall-bladder shadow seen after injection. As regards size, 4 represents the size of the shadow of the normal full gall bladder.

TABLE II.—Showing Results of Investigation by Graham's Method of Four Cases who were not Operated on.

Case.	X-ray Findings.								Deduction.	Remarks.
	Gall-bladder Shadow after Injection.									
	Intensity.				Size.					
	4 hrs. 1	8 hrs. 3	24 hrs. 3	36 hrs. 1	4 hrs. 4	8 hrs. 4	24 hrs. 3	36 hrs. —		
D. J.	—	—	—	—	—	—	—	—	Normal gall bladder	Normal individual with no gall-bladder symptoms.
A. H.	—	—	—	—	—	—	—	—	Pathological gall bladder	Clinically and biochemically—obstructive jaundice.
R. F.	—	—	—	—	—	—	—	—	Pathological gall bladder	Clinically and biochemically—carcinoma head of pancreas.
I. W.	—	—	—	—	—	—	—	—	Pathological gall bladder	Two gall stones shown by ordinary x-ray examination.

deformity of contour of the shadow; or mottling, or central defects (indicating stones in the gall bladder itself, but not causing obstruction, or papilloma).

RESULTS OF INVESTIGATION.

Of the series of cases in which the gall bladders were investigated by Graham's method ten were subsequently operated on and four were not operated on. Our radiological findings and deductions together with the operative or other findings are shown in Tables I and II.

In only one of the ten patients operated on were the surgeon's findings at variance with our radiological deductions. In this case (J. O.) the evidence obtained by an examination of the gall bladder by Graham's method (on two occasions) indicated a pathological condition of the gall bladder. Yet at operation no gross evidence of pathology was obtained. The only abnormality found was a slender adhesion over the common duct; when this had been divided the gall bladder, previously full, was seen to empty. It is doubtful whether this could be sufficient to prevent the dye from passing into and being concentrated in the gall bladder. We trust that a further examination of the gall bladder by Graham's method after the patient has recovered sufficiently from the effects of the operation may throw further light on the question.

CONCLUSIONS AS TO THE VALUE OF GRAHAM'S METHOD.

In estimating the value of this method of radiologically examining the gall bladder three facts must be borne in mind:

1. That only a maximum of 35 to 40 per cent. of gall stones show evidence of their presence by the ordinary method of direct radiography.
2. That a negative finding by the ordinary method does not exclude the presence of gall stones.
3. That "gall stones are incidental and not essential to a cholecystitis which may necessitate surgery."

It is therefore in those cases which by the ordinary direct method of radiography fail to show the presence of gall stones that this new method is of particular value. By no other means short of operation can we ascertain whether this negative finding signifies a normal condition of the gall bladder or not. In other words, it should be used as an adjunct rather than as an alternative to the ordinary method of examination.

A striking feature of the cases shown in Table I is the fact that of all seven cases where a pathological condition of the gall bladder was found at operation, only one case gave the slightest evidence of this before injection, whilst all gave evidence of a pathological condition when examined by Graham's method. The two cases where a definitely normal gall bladder was found at operation both gave definite indication of a normal gall bladder by Graham's method.

The single remaining case where a pathological gall bladder was deduced from the findings by Graham's method, and where at operation no gross evidence of a pathological condition was found, as already stated, needs further investigation.

However, the successes encountered in all the other cases are, in our opinion, a striking testimony to the value of this method as a diagnostic aid in the examination of the gall bladder.

We wish to express our thanks to Dr. Leggate for his valuable assistance, and to those physicians and surgeons of the Liverpool Royal Infirmary who have allowed us to investigate their cases.

BIBLIOGRAPHY.

1. Graham and Cole (1924): *Journ. Amer. Med. Assoc.*, lxxxii, 613.
2. Graham, Cole, and Copher (1924): *Ibid.*, lxxxii, 1777.
3. Graham, Cole, and Copher: *Annals of Surgery*, September, 1924.
4. Carman and Councillor: *Amer. Journ. Roent.*, xii, 403, November, 1924.
5. Whitaker and Milliken: *Surg., Gynecol. and Obstet.*, January, 1925.
6. Graham, Cole, and Copher (1925): *Journ. Amer. Med. Assoc.*, lxxxiv, 14.

LEPTOSPIRA IN LONDON WATERS.

BY

EDWARD HINDLE, M.A.CANTAB., Ph.D.,
F.R.S.E.,

MILNER RESEARCH FELLOW, LONDON SCHOOL OF HYGIENE AND
TROPICAL MEDICINE.

THE occurrence in the British Isles of sporadic cases of spirochaetal jaundice, although apparently few in number, shows that under certain conditions the disease may be acquired in this country, and therefore any observations throwing light on the origin of these outbreaks may help to eradicate the infection. The following notes on the presence of a leptospira, morphologically identical with *Leptospira icterohaemorrhagiae*, in London tap water, Thames river water, and water from several other sources in the neighbourhood, are published in the hope that they will draw attention to these interesting organisms, which under certain conditions may assume grave importance from the point of view of public health.

Attention was first directed to this subject by the discovery of leptospira in one of Professor Leiper's cultures at the London School of Hygiene and Tropical Medicine, which had been prepared to show the ciliated embryos of the broad tapeworm, *Dibothriocephalus latus*. The faeces of a patient infected with this worm, and containing large numbers of ova, had been mixed with ordinary London tap water in a Petri dish, then incubated at 30°C. for sixteen days, and subsequently kept at room temperature for about one week. When examined by dark-ground illumination, the water was found to be swarming with leptospira, there being as many as thirty to forty organisms in each field of the microscope. Under the impression that the leptospira might have had a human origin, the faeces of a number of patients in the Seamen's Hospital, and also from other sources, were mixed with tap water in Petri dishes and the mixtures incubated at 25°C. In every case without exception large numbers of leptospira appeared in the water after an incubation period of ten to fourteen days. Subsequently cultures were prepared by mixing faeces with boiled tap water and incubating, when it was found that these organisms never appeared, thus showing that they were present in the water and not in the faeces. Finally, cultures were made by mixing tap water with faeces, both human and animal, which had been sterilized by boiling, and in every case leptospira appeared after incubating at 25°C. for not less than ten days.

From these experiments it was evident that leptospira is present in London tap water, and an attempt was made to determine in what numbers. Although, subsequently, leptospira was found in the slime on the inside of a dripping tap in the laboratory, the direct examination of water under the microscope gave negative results, even