the stronger snake may, and probably will, engulf his opponent as well as the rat. It has happened that when a boa has engulfed a companion, the keepers have succeeded in withdrawing it by steady traction on their tails. Snakes have not only recovered after such an occurrence, but lived together in amity.

Snake-swallowing by snakes was well known to the ancient Egyptians. When Moses and Aaron met the magicians in Pharaoh's court they knew the trick of making serpents become stiff rods, and when cast upon the ground Aaron's rod swallowed up the Egyptians' rods (Exod. vii, 12).

Meditations on the psychology of the swallowed suggest that the animal world may be divided into swallowers and the swallowed. A whale swallowed Jonah, but it is wiser for a man to swallow a whale than to swallow a lie!

BIBLIOGRAPHY.

- Boulenger, E. G.: *Reptiles and Batrachians*.
 Buckland, F.: *Natural History of British Fishes*, 1880, 160.
 Darwin, C.: *Voyage of the "Beagle"*, 1845, 14.
 Elijah (1 Kings xvii, 23).
 Exodus vii, 12.
 Ditmars, R. L. (1910): *Reptiles of the World*.
 Gadow, H.: *Amphibia and Reptiles*, 1900, 253.
 Gunther, A. C. L. G.: *The Study of Fishes*, 1880, 311.
 Hart, F. R.: *The New England Whale Fisheries*, 1924.
 Hedges, F. A. M.: *Daily Mail*, March 25th, 1925, p. 6.
 Jonah, Chapters i and ii.
 Kingsley, Mary H.: *Travels in West Africa*, 1901, 320.
 Lockwood, S.: *American Naturalist*, 1870-71, iv, 257.
 Matthew xii, 40.
 Parker, N.: *Trans. R. Irish Acad.*, 1892, xxx, 201.
 Procter, J. B.: *Unrecorded Characters seen in Living Snakes*, *Proc. Zoological Society, London*, 1924.
 Regan, C. T.: *Ann. and Mag. Nat. Hist.*, 1925, Ser. 9, xv, 53.
 Roscoe, J.: *The Soul of Central Africa*, 1922, 315.
 Turner, L. M.: *Contributions to the Natural History of Alaska*, 1886.
 Turner, Sir William: *Journ. of Anat. and Phys.*, 1883, xix.

DIFFICULTIES IN THE DIAGNOSIS OF LEUKAEMIA.

BY

G. LOVELL GULLAND, C.M.G., M.D.,

PRESIDENT OF THE ROYAL COLLEGE OF PHYSICIANS OF EDINBURGH; PROFESSOR OF MEDICINE AND CLINICAL MEDICINE IN THE UNIVERSITY OF EDINBURGH.

THE nomenclature of the leukaemias is still in a fluid state, and the subject has been discussed elsewhere (Gulland and Goodall, *The Blood*, third edition, 1925, p. 212). It seems desirable to keep the terminology as simple and yet as precise as possible, so we have adopted the terms "non-granular" and "granular" as the main subdivisions—the former including two acute forms, the myeloblastic and lymphatic, and the chronic lymphatic form, the latter the acute and chronic forms of myelocytæmia.

The acute myeloblastic and lymphatic forms are probably the most common, but many of them are certainly not diagnosed, and in any case their duration is short. The chronic forms last for years, are practically always recognized sooner or later, and bulk therefore more largely in the mind of the practitioner. I shall accordingly deal first with them.

CHRONIC GRANULAR LEUKAEMIA.

In the chronic granular leukaemias the disease is so insidious in onset that the first thing to attract attention, if the patient comes under observation before the anaemia is marked, is usually the enlarged spleen. It is surprising how often there seems to be difficulty in determining whether it is the spleen which is enlarged or some neighbouring organ. I remember a febrile leukaemia in a woman, many years ago, which was sent into a surgical ward as a cellulitis of the abdominal wall. The surgeon was clear that this was a wrong diagnosis, but thought the spleen was a tumour of one of the pelvic organs, and transferred the case to a gynaecologist, who was certain that the tumour was not pelvic, but refused to determine its nature. Of course a blood examination settled the matter. Enlarged spleens in leukaemia may and do wander all over the abdomen, and might be taken for any organ in turn, according to their temporary position. Comparatively small enlargements are often taken for tumours of the splenic flexure, and several times patients have come to me who had been x rayed after barium meals and enemas. The true nature of the mass was only suspected when the passage was found normal. It often takes much more than a momentary examination to settle whether a given enlargement is spleen or kidney. Neither organ, when enlarged, invariably retains its typical shape; the kidney may become very superficial, and the colon which crosses it may be loaded, or empty and spastically contracted, and not easily made out, and there are many cases of kidney swellings with blocked ureters in which the urine gives no help. Some cases have required air enemata, ureteral catheterization, full blood examination, etc., before their nature was definitely determined.

To-day I saw a man whose right kidney was removed some years ago for calculi associated with constant pain and hæmorrhage and the formation of a large tumour. At operation the kidney was found to be cystic in addition, and it was suspected that the left kidney was in the

same state. It has steadily enlarged and the diagnosis is undoubtedly correct. Since my last examination the kidney had increased considerably in size, and so altered in shape that it looked at first as if a splenic enlargement was superadded to that of the kidney. The converse of this case was supplied some time ago by a very anaemic woman with great digestive disturbance, who came to me with the diagnosis of cancer of the stomach. Her left kidney had been removed for pyonephrosis many years before, and the spleen, which was only moderately enlarged, did not come forward as usual, but, in the recumbent position, lay well back in the left flank and under the ribs, and had to be looked for. She had advanced granular leukaemia.

The blood in these cases when they first come under observation is almost invariably so typical that no error is possible if the attendant examines it or causes it to be examined. The initial count, however, even with a markedly enlarged spleen, may vary greatly—from 500,000 or more to 42,000, 35,000, and 22,000, which are the lowest I have seen. The blood in these cases with low counts was nevertheless quite typical. The only likelihood of error is the presence of a febrile complication, which sometimes causes a drop in the white count to normal or subnormal figures, or the rare possibility of a spontaneous remission. But even then abnormal cells will almost always be present which will give the necessary indication, though they are sometimes few in number.

The effect of radiation on the blood is by no means uniform and is difficult to forecast. The rule, of course, is a marked diminution in total numbers with a special drop in the immature granular cells, but there may be no effect at all. Generally speaking, this refractory state is more common in cases that have previously been irradiated, but it does occur in cases treated thus for the first time as soon as the disease was diagnosed; and, on the other hand, some cases will respond repeatedly to radiation until the course is cut short by some complication. It is important to remember that the effect of radiation persists for months sometimes, and that there may be a progressive diminution in the number of the leucocytes. For this reason one usually stops radiation when the white count has reached 20,000, as there will probably be a further drop when treatment has ceased. The case mentioned above with an initial count of 22,000 was treated by x rays to the long bones for special reasons. This was stopped when the count fell to 12,000 three months ago, and the count is now about 5,000 with practically normal proportions. In the last films no immature cells of any kind could be found; the usual excess of basophils had disappeared, and the film looked like an ordinary mild secondary anaemia with a slight leucopenia. After prolonged search one normoblast was found. If this film had been examined by someone unacquainted with the history of the case, diagnosis would have been impossible.

Radiation may apparently, however, sometimes produce

a condition closely resembling myelocytæmia. Some years ago, in November, I saw a lady whose breast had been removed for cancer a year before. She had been treated by x rays from that time till August. She was anaemic—of secondary type—there were enlarged glands above the inner ends of both clavicles, the liver was down to the umbilicus and apparently nodular, and the spleen was palpable. Generalized carcinomatosis was the probable diagnosis, but the examination of the blood introduced a doubt. The white cells were 12,800, there were numerous neutrophil myelocytes and some myeloblasts, and, further, there were nearly 7,000 nucleated red cells to the cubic millimetre, about half of which were normoblasts, the rest megaloblasts. A second count, a week later, gave 9,400 whites with the same picture as before and the same number and proportions of nucleated reds. The blood picture was that of an acute myelocytæmia; the eosinophil and basophil excess of the chronic type was absent. The alternative diagnosis was widespread metastatic cancer in the bone marrow, which may produce a blood picture resembling leukaemia, and which was obviously more likely in this case. The lady returned to London and died in Westminster Hospital about six weeks later. Dr. Hildred Carlill kindly wrote to me about the subsequent history. A fortnight after I saw her the white count was 8,000—neutrophil polymorphs 59 per cent., lymphocytes of all forms 40 per cent., eosinophils 1 per cent., no myelocytes, and no nucleated reds. The spleen had returned to normal in size. A later count was almost exactly similar. At the *post-mortem* examination cancer was found in the liver and ovaries, but not in the bones examined, and indeed, in view of the later counts, it was not to be expected. One is almost driven to the conclusion that my findings represented an erythroblastic and leucoblastic crisis resulting from the prolonged x-ray treatment.

CHRONIC LYMPHATIC LEUKAEMIA.

Chronic non-granular or lymphatic leukaemia is much rarer than the granular form, and less dangerous to life on the whole. I have seen cases, ultimately fatal, in which the first glandular enlargement—which had never disappeared—had been noticed eighteen and twelve years earlier respectively. Both these patients had been in fair health until a year or two before the end. In such cases as these the process almost certainly starts in the lymphatic glands and gradually involves one group after another; the final anaemia does not set in until the bone marrow becomes completely involved. There is no reason why the process should not start in any bit of lymphatic tissue in the body, but a start in such extraglandular lymphatic tissue as the follicles in the bronchi or the intestine would mean an almost immediate spread to glands, and an apparent glandular onset. In cases where the process starts outside the bone marrow, the chronicity or acuteness of the individual case seems to depend entirely on the period when the marrow becomes involved, and on the rate at which the marrow space is filled by lymphocytes. When this occurs the area available for erythroblast proliferation is encroached on and an anaemia which is practically aplastic in character always results.

But this anaemia is rarely so severe as in the acute myeloblastic and lymphatic forms, in which the starting point is in the marrow itself. Nor is the tendency to haemorrhage so great on the whole as in the chronic myelocytæmias. The chronic lymphatic cases are much more apt to die from the pressure effects of the glands, which are so much more uniformly enlarged in this form than in any other, from the tendency to thrombosis of veins, and from the interference with function due to the infiltration of vital organs. As internal glands may sometimes be the first affected, pressure effects may occur early, and may cause puzzling clinical pictures until the blood is examined. An interesting example of this difficulty presented itself about eighteen months ago.

A lady who had complained of shortness of breath for about six weeks developed pleurisy with effusion without rise of temperature, about a fortnight before I saw her. The effusion had to be tapped several times before it ultimately settled down. Her doctor, who had gone over all the usual possible causes without result, examined her blood and found a great excess of lymphocytes. My first count

was: red cells 4,700,000, haemoglobin 75 per cent., white cells 55,800. Of the whites, 91 per cent. were lymphocytes, 9 per cent. polymorphs, and there were, in addition, many degenerated lymphocytes; there were no myeloblasts. There was no enlargement of liver or spleen; one enlarged gland could be felt above the inner end of the right clavicle, none elsewhere, but the veins in front of the chest were unduly prominent. The single gland was evidently an offshoot from a mediastinal mass. During the next six months while she was under observation the white cells fluctuated between 35,000 and 85,000; the proportions remained much as in the original films. The left lung was slightly dull all over, with diminished expansion. This was evidently a pleurisy resulting from the presence of a mass of leukaemic glands in the mediastinum. The patient is now well symptomatically, except for some weakness and shortness of breath on exertion; no other glands have appeared.

As these cases advance the white count usually rises slowly, and may reach a very high figure. In March of this year a man was seen who had noticed some enlarged glands at the back of the neck in 1918; other groups were gradually involved, and at his first visit all the external glands were greatly enlarged, and the spleen reached the umbilicus. A diagnosis of lymphadenoma had been made, but the blood had never been examined until he gave up his work elsewhere and settled in Edinburgh. My first count gave: red cells 2,680,000, haemoglobin 50 per cent. I had drawn the blood up to the figure 1 in the pipette, and in the counting chamber the white cells were so closely packed together that the ruled lines could not be seen. When the count was repeated with the blood drawn up to 1/10 it gave the astonishing figure of 2,224,000, which, so far as I have been able to find, is the highest count of white cells recorded in any condition. The films bore out this count; the white cells seemed quite as numerous as the red, and were practically all small lymphocytes. The patient was treated with arsenic. Six weeks later the glands had gone down to about a quarter of their former size—very often a bad omen in these cases unless it is accompanied by a general improvement in health. He had thrombosis of the veins of the left leg, and generalized oedema with pleural effusion; he ultimately died suddenly. On this occasion the red count and haemoglobin were rather better than before, the white cells had fallen to 1,132,000, and were again practically all small lymphocytes. One neutrophil polymorph was seen in counting 500 cells. Though he had never been robust he had been able to carry on office work until a few months before his death.

ACUTE MYELOBLASTIC AND LYMPHATIC LEUKAEMIAS.

Every physician is familiar with the hopelessness and tragic rapidity of the acute myeloblastic and lymphatic leukaemias. It has happened to me to see a patient who was well until a certain Saturday, to make the diagnosis on the following Tuesday, when he had a count of 10,000 white cells, and to see him die on the Friday with a count of 20,000. Recently the opportunity presented itself of observing throughout a case very nearly as rapid. The first recognizable symptom was epistaxis on the 20th of the month; there was haemorrhage later into the skin and from the kidney and bowel, and the termination was profuse haematemesis on the 29th. The red count was at first 4,800,000, haemoglobin 65 per cent. It showed a steady fall, and on the 29th was 2,400,000, haemoglobin 42 per cent. This is typical of these cases. The rule is a gradual drop in red cells with a rising colour index—in this case from 0.6 to 0.87. If the case lasts long enough it is not uncommon to get a colour index above unity, and to find megaloblasts as well as normoblasts in the blood. Another point of importance is the progressive diminution in the number of blood plates. The white count ran: 9,300, 6,400, 10,400, 15,700. At first the polymorphs were 30 per cent., and at the end 10 per cent.; at first also there were some neutrophil myelocytes and some eosinophils and basophils. The percentage of non-granular cells was about 60 in the first films, of which 48 were myeloblasts. At the end the non-granular percentage had risen to 88, of which 65 were myeloblasts.

On analysing my cases it is evident that the acuteness of

the disease is nearly proportional to the percentage of myeloblasts and to the rate at which the percentage increases. Cases in which the true lymphocyte proportion remains high last much longer. Most of the cases of acute leukaemia are easy to diagnose, because though the leucocyte increase is small, there is usually an increase, and, except for the foregoing consideration, it matters little whether the excess is lymphocyte, myeloblast, myelocyte, or promyelocyte, or a mixture. The important point is the diminution of polymorphs and the great excess of cells with a round nucleus. With an ordinary Jenner stain all the forms except the myelocytes look very much alike, and special staining methods have to be used to differentiate them. But there is a group of cases which are very doubtful. Two recent examples will illustrate it.

The first was a lady of 52 who had suffered from melancholia for ten months. For a fortnight the temperature had been high. No physical signs of any kind were present to account for this; the Widal reaction was negative to typhoid and the paratyphoid fevers, and no enteric bacilli were found in the stools. There was no enlargement of external glands, the splenic dullness reached the border of the ribs, but the spleen could not be felt definitely. The white count was only 1,900. Not a single polymorph was seen; about 35 per cent. of the whites were myeloblasts, the remainder lymphocytes. She died two days later.

The second was a man of 50 who was first seen because of an irregular pneumonia which had lasted for about a week. The white cells were 16,300, with only 70 per cent. of polymorphs, and no glycogen reaction. The comparatively low polymorph percentage made me suspicious that there was some condition behind the pneumonia. A week later the lung had cleared up, and the white count had fallen to 7,600, polymorphs 54 per cent., lymphocytes 40 per cent., eosinophils 3 per cent., myelocytes 2 per cent., myeloblasts 1 per cent., and a few normoblasts; fairly numerous blood plates. A week later he was definitely anaemic: red cells 3,250,000, haemoglobin 50 per cent., white cells 6,600—polymorphs 22 per cent., lymphocytes 66 per cent., myeloblasts 8 per cent., myelocytes 2 per cent., eosinophils 2 per cent., a few normoblasts; blood plates few and large. Three weeks later (intermediate counts are omitted) the red cells numbered 1,000,000, haemoglobin 18 per cent., white cells 5,900—polymorphs 18 per cent., lymphocytes 66 per cent., myeloblasts 12 per cent., myelocytes 4 per cent., with numerous normoblasts; there were no megaloblasts. There was no enlargement of spleen or glands throughout and no haemorrhages—only progressive weakness as the main symptom.

How are we to class these cases in the absence of *post-mortem* evidence? In spite of the low white counts I can only regard them as acute leukaemias—the first probably mainly myeloblastic, the second mainly lymphatic. The differential count must be the criterion. The whole question, however, depends on the significance of myeloblasts occurring in the circulating blood—a subject too large for discussion here. It is to be hoped that some better method of differentiating myeloblasts in films than either Jenner-Giemsa or the oxydase reaction will be discovered, and that some haematologist will undertake the embryological studies necessary to clear up our views of the phylogenetic relationship of these cells.

In connexion with the type of case just described, it will be realized that difficulty will sometimes arise in the diagnosis from pernicious anaemia, and this for two reasons—first, the not infrequent occurrence of a colour index of unity or above it in myeloblastaemia, and secondly, the high non-granular percentage in the differential count of pernicious anaemias. The first difficulty only arises if we are to regard cases of the type quoted as leukaemias, for increase of the white count is the rule in myeloblastaemias, but is very rare in pernicious anaemia, and when it does occur is always a polymorph leucocytosis, and not a non-granular increase. The second difficulty may be illustrated by the following recent case. A man of 65 who showed all the characteristic signs and symptoms of pernicious anaemia, had a red count of 1,500,000, haemoglobin 35 per cent., white cells 3,500; of the latter 84 per cent. were lymphocytes, the remainder polymorphs. There were no myeloblasts. At first sight such a percentage makes one think of an acute lymphatic leukaemia, especially if the whites have not been counted, but the actual number of lymphocytes in the cubic millimetre is not really greatly increased. If we assume that the normal number of white cells per cubic millimetre is 7,000, of which polymorphs are 70 and lymphocytes 30 per cent., the actual number of polymorphs in the cubic millimetre will be 4,900, and of lymphocytes 2,100, while in the case quoted the lymphocytes are 2,940—not an overwhelming increase.

We are far too apt to use the percentages of the differential count; it would be wiser and would avoid misconceptions if we always quoted the absolute numbers of the different cells in the cubic millimetre. One safeguard in this bit of diagnosis is that myeloblasts are rare in pernicious anaemia, though they do occasionally appear. I have never seen them in sufficient number to cause error.

The difficulty of diagnosis between lymphatic leukaemia and glandular fever may be a very real one. Most cases of the latter disease are of short duration and do not present much difficulty, but in a recent case the temperature lasted for eight weeks and the enlargement of the glands nearly as long. The patient was a woman of 22; the enlarged glands were widely distributed on both sides of the neck but not elsewhere; there was slight enlargement of the spleen. The red side of the blood was a mild chlorosis; the red cells dropped slightly as the disease progressed. The first white count taken by her doctor, about a fortnight after the onset, was 16,000—polymorphs 14 per cent., lymphocytes 86 per cent. My first count, a few days later, was 12,500—polymorphs 26 per cent., lymphocytes 74 per cent.; but in addition there were a great many degenerated lymphocytes with smeared nuclei; practically all the lymphocytes were "large." No myeloblasts were seen. Thereafter the count gradually dropped as the temperature fell and the glands diminished in size, and it has now for some weeks run about 3,000, with such proportions as polymorphs 44 per cent., lymphocytes (almost all small) 54 per cent., eosinophils and basophils 1 per cent. each—that is to say, the blood picture is now that of a chlorosis with the polymorph leucopenia so common in that condition. Before I saw the patient the doctor had given the diagnosis as lymphatic leukaemia—a very natural conclusion on the evidence. I have heard of glandular fevers lasting a longer time, but have never seen one.

"A SARCOMA OF THE LEUCOCYTES."

No more need be said about the essential nature of leukaemia than that I regard it as a sarcoma of the leucocytes. The assumption that the disease is a sarcomatosis, however, only removes the difficulty a stage further back, and the problem is bound up with the question of what influence it is which in health keeps the normal number of leucocytes in the blood within such narrow limits. We know that the action of parasites, using the term in its widest sense to include bacterial and other organisms, can powerfully alter the count in disease in both positive and negative directions, that the proteins of food and probably other proteins and many other substances have definite actions. Does the interaction of an infinite variety of substances keep the balance in health, and, if so, does it act directly on the bone marrow and the blood, or through the medium of some other organ? These questions require a definite answer before we shall progress much further in the understanding of leucocyte problems.

It is very striking when one examines the marrow of a myeloblastaemic patient who has died with a count of, say, 20,000, to find all the tissue one can examine packed with myeloblasts, to the practical exclusion of every other type of cell. The wonder is, not that the leucocyte count is increased, but that it is not vastly higher. One cannot resist the feeling that the excess of cells in the blood is there rather as an overflow than in response to any active call upon the marrow cells. Obviously the condition is widely present in the marrow before there is any evidence in the blood of its existence. A leucocytosis responding to an organismal infection occurs practically at once, and the marrow gets under way immediately, but never goes beyond the amount of hypertrophy required in each special instance. A leukaemia progresses in the marrow more or less steadily, in the acute cases progressively till death occurs. We do not know whether the spontaneous remissions which sometimes occur in the blood picture of the chronic leukaemias correspond to a real alteration in the condition of the marrow; if they do they are not more frequent than the retrogressive changes that occur in some tumours. It is well to keep an open mind, but until further research throws fresh light on the disease the neoplastic theory of its causation is the most logical and satisfying explanation.