DIABETIC TREATMENT SIMPLIFIED.

BY

FRANK C. EVE, B.A., M.D.CAMB.,

F.R.C.P.LOND.,

PHYSICIAN, ROYAL INFIRMARY, AND TO CHILDREN'S HOSPITAL, HULL.

Few will deny, least of all general practitioners, that the treatment of diabetes has now become very complicated, and, in my opinion, there is a real danger of the main issues becoming fogged in a cloud of bewildering details. The main issues are: (1) the quantity of sugar excreted day by day; (2) the presence or absence of ketones; (3) treatment by regulating the quantity and quality of the diet, or by giving insulin if dietetics prove inadequate; (4) the blood sugar (in special cases).

The ground can be cleared by distinguishing clinically between cases of simple glycosuria, which are common and usually unimportant, and stubborn diabetes, which is uncommon and serious. This paper deals with genuine diabetics who are living at home.

Measurement of the Daily Excretion of Sugar.

Obviously it is a cardinal necessity that both doctor and patient should have daily knowledge of this over prolonged periods. Without this knowledge treatment cannot be either intelligent or effective. Yet, in private practice, even at the present day, this indispensable minimum of knowledge is

CHART 1.—Diabetes and acute phthisis in a woman, aged 21. Showing the effect of insulin in pinching the curves together when diet was powerless to do so. A, 20 units of insulin injected; B, 15 units of insulin injected; C, 10 units of insulin injected; D, strict diabetic diet. The patient's appetite was poor from August 29th till September 14th, after which it improved, needing more insulin.

lacking, or rather is available to doctor and patient only at rare or occasional intervals. The daily variations in sugar excretion—which I know to be very wide—are hidden: only the occasional testing days are illuminated.

This deficiency, during the last four years, I have remedied by a very simple method whereby the patient himself is able to measure and record on a chart the daily variations in the excretion of his sugar. I have yet to meet the private patient who was not able and willing to do this, usually with enthusiasm. The method is described at the end of this paper.

Now what does this daily chart of sugar excretion enable us to do?

1. The doctor can see at a glance whether in his absence the patient has been going on well and has been obeying instructions. In the green-pea season a patient who was too fond of them was detected by his chart. A paint manufacturer's sugar was seen to increase owing to worry when the price of oil jumped. A small boy in hospital crept from his cot at 3 a.m. and devoured a box of chocolates. Chart 3 reveals this and his other falls from grace. Further, the patient himself can see when he makes mistakes. He takes pride in producing a chart without such blemishes, and hence is more careful in his diet. Also he is rewarded visibly for his starvation days if they are imposed (Chart 4).

2. If insulin is being given in private practice one has to leave a little sugar (1 or 2 grains per ounce) in the urine as a safeguard against hypoglycaemia. By the help of the daily chart (see Chart 1) this can easily be done by adjusting the amount of insulin or bread. Without a daily chart the steering must be very wild—and dangerous, I would think.

3. The daily chart enables the practitioner to send calories to the devil, where they belong. Calories are well enough in a research laboratory, but human patients are not engineering propositions. Each man varies in his efficiency,

both digestive and metabolic; he varies in respect of the calories he has to spend in exercise or in keeping up his body heat in variance with his house, his clothes, and the weather. Then there are the varying hosts of bacilli which we feed in our intestinal canal and which form so large a proportion of our faeces.

For reasons of this kind it is, I fancy, a pseudo-scientific presumptuousness to pretend to be able to work out by arithmetic a cast-iron table of a man's food requirements. On the other hand, Nature does give a man an appetite varying roughly with his daily needs (bulimia is quite exceptional in diabetes and at any rate laughs at calories). And if restrictions of the quality of his diet are shown by the sugar chart to be inadequate, the patient's quantity of food must be restricted, especially if he weighs too much.

What I do is to put the diabetic gradually on to a diet free from carbohydrate. If sugar persists and no ketones appear, diminish the quantity. If ketones appear, 1 to 3 ounces of bread (weighed) is added, and enough insulin injected to bring the sugar almost to zero. Or weekly "starvation days" on greens, bouillon, and coffee may obviate the necessity for insulin.

I do not claim that the daily chart of sugar excretion is a perfect method, but I have found it reliable, adequate, and most helpful. A daily chart of the blood sugar would in some ways be more perfect. But this fluctuates violently from hour to hour; it yields no summative measurement as the urine does; and, at any rate, daily blood sugar tests are impracticable in private practice.

4. Another point of interest (and perhaps of novelty)

CHART 2.—Simple glycosuria easily cured by diet; no ketones. Diet makes the two curves coincide.

shown by the daily charts is that occasionally the urine, when fermented, has the same specific gravity as water. At the beginning of the little boy's chart (3) this occurred for two consecutive days, and also in Chart 4. Weak sugar and water does the same when fermented. If this be verified by more accurate methods it would mean that the kidneys secreted no other solids than sugar on those brief occasions, although the patient's health does not seem affected.

Insulin.

The main practical points are:

1. Insulin is something which somehow enables a diabetic to burn up sugar like a normal person, but does not enable him to store it as glycogen.

2. Insulin can in this way cause the blood sugar to sink to a dangerously low level (hypoglycaemia). This danger can be guarded against if the patient always carries about with him eight large lumps of sugar (1 or 2 ounces of barley sugar—Maclean) and promptly eats them all if he feels queer after an injection of insulin. Two grams of glucose are supposed to keep busy one unit of insulin; from which I reckon that eight average lumps of sugar would act as an antidote to 15 units of insulin.

3. Insulin is given two or three times a day, just before a chief meal, which if possible should contain 1 ounce of bread; 10 to 50 units daily is the usual range of dose; more is seldom useful.

4. The patient can be taught to inject it into the pinched-up skin of the thigh, under weekly supervision. A trained nurse is better, but a doctor cannot be at hand three times a day before meals.

5. Insulin has to be acid; in a few subjects this causes painful lumps. It is said that neutralization in the syringe by sterile sodium bicarbonate solution of proper strength prevents this irritation and does not impair the action of the insulin.

6. As to blood tests with insulin, I have rarely found them helpful (see "Coma in diabetes"), for it is unattainable

in private practice to keep the blood sugar normal. You dare not even keep the urine quite free of sugar in insulin cases.

7. The risk of giving insulin in unsuitable cases will be practically eliminated without blood tests (a) if you first try to cure all cases by diet alone; (b) if ketones are present, and abundant sugar, insulin is surely needed; (c) leaky kidneys (renal glycosuria) are a real but uncommon danger. These cases may be recognized by their uncanny failure to improve with diet, by their good health, by the absence of ketones, and by the absence of glycosuria three hours after a good starchy meal. A blood sugar will decide the matter by showing a low renal threshold for sugar. So also, I imagine, would the absence of rhythmic oscillation in a weekly starvation chart (as Chart 4); but

venously nowadays). We must certainly give insulin freely; but the terrible question is, how much? Either timidity or rashness may kill. If you can have a blood sugar test every hour or so, then you are on sure ground. But this must not entail delay. Failing this, it seems to me you must guard your big doses of insulin with glucose. This will ensure against hypoglycaemia, and by its combustion it should destroy the poisonous ketones. However, some authorities are opposed strongly to injecting sugar, though their reasons seem weak to me. Agreement is not yet reached, but if you can inject sugar you do not require blood tests.

There is no time in coma to trifle with half-measures. Hence intravenous injections are needed, starting with 20 to 40 units of insulin, 40 to 80 grams of glucose in

CHART 3.—Diabetic boy aged 8. The shaded area between the curves is the quantity of sugar in the urine day by day. Sugar reappears after Christmas indiscretions.

I have seen no such case in the last four years. To insist that we should test the blood of every glycosuric patient before giving insulin so as to avoid this fairly obvious bugbear seems to me absurd.

8. I do not recollect seeing a single ill effect attributable to insulin, while the benefits are notorious. But if in an emergency you are nervous of causing hypoglycaemia you can guard the insulin with sugar, as above.

Ketones.

Ketones, such as acetone and diacetic acid, are half-oxidized fats. Their presence in the urine is revealed by the nitro-prusside test or the ferric chloride test; I prefer the latter, as the former is too sensitive and becomes a scaremonger.

If the addition of two drops of liq. ferri perchlor. fort. to the urine produces a dark-brown colour then ketones are present in dangerous amount, and must be got rid of, lest coma supervene. Ketones are far more pregnant with danger than sugar, and are often produced by too strict a dietary. They indicate that the fats in the system are not being properly burnt up. Now how can these poisonous half-oxidized ketones be thoroughly and harmlessly burnt up? By giving more carbohydrate food, for the fats are combusted in the furnace of brightly burning carbohydrates. But the special weakness of the diabetic is that he cannot burn up his carbohydrates properly. Hence, to get rid of ketones, it is often needful not only to give extra carbohydrate, but also to give insulin, which has the mysterious property of enabling the diabetic to burn up sugar like an ordinary person. So that if ketones appear, watch for coma, ignore the sugar in the urine, and treat the much greater danger by giving alkalis freely and more bread. And if you find that large doses of alkali fail to make the urine alkaline, then the peril of coma is menacing indeed. Ten grains of sodium bicarbonate should alkalize a normal urine; so try your ketosis patient with 20 grains. An hour afterwards boil and cool the urine in a test tube and test with blue litmus paper.

Coma in Diabetes.

Here I can only touch upon the main principles. Since the coma is probably due to poisoning by the "acid" bodies formed by the imperfect combustion of fats, we are told to give alkalis freely by mouth or rectum (not intra-

800 to 1,600 c.cm. of water. Then you must feel your way by results, but 80 to 100 units or even more may be needed in the first day. Croton oil, strychnine, and camphor will also probably be indicated. I have seen nothing in medicine more dramatic than the swift escape from the very jaws of death into the enjoyment of apparent health which usually follows this treatment.

Persistent vomiting, with coma, is a very fatal complication. Although albuminuria may be slight, I have wondered if the real poison was uraemic in nature; but I have not yet had a chance of measuring the blood urea. Certainly, after long periods of diabetic overwork and irritation, the kidneys frequently show signs of wear and injury. Thus, numerous casts, albumin, and diminished sugar are often found in the urine in coma.

Exercise.

Exercise in moderation is important in diabetes to help the combustion of sugar to carbonic acid which takes place in the muscles. Gardening is an excellent hobby for diabetic patients, and often makes them look the picture of bronzed and ruddy health. Bed should be prescribed only during starvation or illness.

Method of Domestic Estimation of Urinary Sugar.

The apparatus required is: (1) A urinometer with a long stem indicating clearly specific gravities 1000 to 1050. This, and its glass cylinder, should have a case for travelling. (2) An engineering student's exercise book of squared paper. Its left-hand margin is also graduated 1000 to 1050.

(3) A thermos flask, pint size. (4) A pennyworth of yeast, fresh twice a week.

Method.

Every night the specific gravity of the urine is taken when warm. The thermos flask, which has been warmed and cleansed with hot water, is filled with the warm urine; a piece of yeast the size of a small wainut is added and shaken up. The flask is corked loosely (or with cotton-wool) and wrapped in blanket. Next morning the specific gravity is again taken. It will be less, because all the sugar has been converted by the yeast into carbonic acid gas (mainly). Let us suppose the gravities before and after fermentation were 1040 and 1020. A red dot and a black dot will be placed at their proper heights in the first column of the squared page. Next day another pair of dots will appear in the next column. Thus will be spun out an upper and a lower curve, one red and the other black. The space between these curves represents the excretion of sugar in grains per ounce. The object of treatment is to pinch these two curves together. When they coincide there is no sugar.

CHART 4.—Showing effect on sugar excretion of Sundays on greens only. The two curves approximate at the week-ends (no sugar). From a case of diabetes with myxoedema.

and it is well to mark these days by a red dot within a circle.

If the morning urine alone is recorded the results will be more constant, more flattering, and less useful. The night specimen is best for a patient who does not always urinate at home. A twenty-four hours specimen is not much better; it is a nuisance to warm, and is only available for the home-keeping patient.

Ketone Test.—Every morning the patient watches two drops of liquor ferri perchlor. trickle down the near side of the urine glass. If it changes from yellow to dark brown the doctor must be informed that day.

THE SCHICK TEST AND ACTIVE IMMUNIZATION AGAINST DIPHTHERIA.

BY

JAMES P. WATT, M.A., M.D., D.P.H.,

CONSULTING MEDICAL OFFICER OF HEALTH FOR ABERDEENSHIRE.

The burgh of Fraserburgh has, according to the Registrar-General's estimate for 1923, a population of 10,367. The births in that year numbered 257, and the average for the preceding five years was 287.

Incidence of Diphtheria in the Burgh.

Since 1900 there has been in Fraserburgh, as in other towns in the kingdom, an increasing incidence of diphtheria, as the following total notifications indicate:

1900-04	...	...	...	...	...	48
1905-09	...	...	...	...	...	127
1910-14	...	...	...	...	...	164
1915-19	...	...	...	...	...	61
1920-23	...	...	...	...	...	401

In view of this increased incidence and the cost of hospital isolation, the Public Health Committee, on my recommendation, decided to try the newer measures of prevention by toxin-antitoxin inoculation, and accordingly, on March 25th, 1924, decided to open a series of clinics, free of charge, for the inoculation of children under 5. The clinics were to be held once a week in the Town House, and to be conducted by the local medical practitioners in rotation, who had cordially agreed to give their assistance.

In recommending that children under 5 should be dealt with first, I was influenced by the fact that at this age reactions either do not occur or, as Dr. Park says, are "slight or negligible." The susceptibility at this age to diphtheria (70 to 80 per cent.), and the fact that half of the diphtheria deaths are among children under 5, were also reasons for trying to get as many of them as possible inoculated. It seemed likely also that the absence or slight nature of the reactions in young children would lead parents more readily to give their consent for the inoculation of the older children when the schools were dealt with, and this proved to be the case.

The first step in the way of propaganda was the distribution through the schools of a leaflet to all parents or guardians having children attending school, explaining the purpose of the clinic and the value of toxin-antitoxin inoculation, while the same leaflet was shown in large posters in shop windows and other places throughout the town.

The following is a copy of the leaflet.

DIPHTHERIA PREVENTION.

To Parents or Guardians:

Diphtheria is a most dangerous disease, and is particularly fatal to young children. Of a hundred deaths from diphtheria, 80 are under 10 years of age, the larger proportion being under 5.

Your children need not have diphtheria. It can be prevented.

Some people are able to resist the germs of diphtheria, others are not.

A simple method—painless and harmless—has been discovered to find this out. It is called the Schick test.

If this shows that your child does not resist diphtheria germs, it can be inoculated and protected against diphtheria.

As almost all children over 6 months and under 5 years of age are liable to contract the disease, it is not necessary to apply the Schick test to them. They should all be inoculated and protected. The inoculations are followed by no constitutional reactions and very little local irritation.

This method has been used for several years, and so far almost no one who has been so protected has got diphtheria.

Diphtheria can be stamped out and many lives saved every year if mothers will have this done and protect their children.

The Town Council of Fraserburgh have resolved as a first step to give parents an opportunity of having all children under 5 years of age and over 6 months protected free of charge. They have accordingly arranged to open a clinic in the Town House to which children under 5 may be brought. The clinic will commence on Wednesday, the 11th June, and will continue for six weeks, a meeting being held each Wednesday at 2.30 in the Town House. This clinic will be conducted by Dr. Beddie.

Subsequent clinics will be conducted by the other doctors in rotation.

To obtain complete protection each child will have to be brought to the clinic three times, at a week apart.

Articles giving a more complete explanation of the Schick test and toxin-antitoxin inoculation were published in the Aberdeen and Fraserburgh newspapers, and we were greatly indebted to the press for the strong support they gave us in the work. I carried out also, as far as time permitted, a personal visitation of the homes, more especially in the poorer parts of the town where diphtheria had been most prevalent.

Clinics.—The first clinic was opened on June 11th, but there was no attendance on the first three occasions. To the fourth clinic, however (July 2nd), 5 children were brought, and to the fifth 10. To the next three clinics 4, 3, and 0 came forward—the total attendances being 19, 17, and 6. To the eight clinics, therefore (June 11th to July 30th), only 22 children had been brought. This was largely due to the fact that, owing to absence on holiday in July, I was unable to continue the visitation of the homes which I had begun in June. On my return this visitation was energetically resumed, and the numbers coming to the clinics began to increase. On August 6th, 15 fresh children, and on August 13th 41, came, making the total at this latter clinic 57. With the consent of the Public Health Committee a second weekly clinic was opened on August 18th, and as the numbers continued to grow a third clinic was commenced on September 5th. These three clinics were carried on throughout September, October, and November, while one of them was continued into December. Apart from school children, there was a total attendance of 596 children under 5, 92 per cent. of whom were inoculated thrice, while of the remaining 47, 25 had two inoculations and 22 had one; 780 children from the schools were also inoculated at these clinics.

Schools.—On September 4th, by which date 183 children had attended the clinic, the town council resolved unanimously to take measures to deal with the schools, and authorized me, with the assent of the education authority, to ask for the consent of parents to inoculate those under 7 without Schick testing, to use the Schick test for those over 7 and inoculate those found Schick-positive, and to obtain such assistance as might be necessary. I did not think it desirable to apply the Schick test to children under 7, and the experience of applying the Schick test in the schools has confirmed me in this view. The head masters and mistresses of the schools were approached, and they most kindly promised their hearty co-operation. With their assistance consent forms were distributed to all parents and guardians; the forms were of two kinds, according as the children were under or over 7. Class sheets were provided, and these were very kindly filled in by the teachers of each class from the consent forms returned. No propaganda was employed further in connexion with the schools, other than had already been carried out for the children under 5. There are five schools in Fraserburgh, of which three are attended by children under 7—namely, the Infant, the North, and St. Peter's Episcopal. St. Peter's and the North School have children also up to 14, and the Infant School to 11. The Central is attended by children from 8½ to 14 years; the Academy pupils are over 12 years.