2. Excluding certain conditions, the reaction is of value in the diagnosis of malignant disease.

REFERENCES.

- ¹ Murray: Eighth Scientific Report of the Imperial Cancer Research Fund, 1923, p. 75. ² League of Nations Health Organization, Report of Technical Laboratory Conference, C.S.M.5, 1924, III, p. 102. ³ Sampietro and Tesa: *Annal. d'Igiene Speriment.*, 1908, xviii, 657. ⁴ Weinberg and Mello: *Bull. de l'Assoc. française pour l'étude du Cancer*, 1910. ⁵ Von Dungern: *Muench. med. Woch.*, 1912, lix, p. 65. ⁶ Pfeiffer and Finster: *Wien. klin. Woch.*, 1909, xxviii, 989. ⁷ Weil: *Journ. Amer. Med. Assoc.*, 1908, 12, p. 158. ⁸ Crile: *Ibid.*, 1908, 12. ⁹ Elsberg: *Ibid.*, 1909, 13, 1036. ¹⁰ Marcus: *Berl. klin. Woch.*, 1908, 689. ¹¹ Brieger and Trebing: *Ibid.*, 1908, 1041. ¹² Ascoli and Izar: *Muench. med. Woch.*, 1910, 403. ¹³ Abderhalden: *Die Schutzfermente des tierischen Organismus*, Berlin, Julius Springer, 1912. ¹⁴ Leitch: *BRITISH MEDICAL JOURNAL*, 1914, ii, pp. 151 and 330. ¹⁵ Weichardt: *Med. Klin.*, 1909, 9, p. 328. ¹⁶ Freund and Kaminer: *Wien. klin. Woch.*, 1910, xxviii, 1921. ¹⁷ Shaw-Mackenzie: *Lancet*, 1922, ii, 759. ¹⁸ Kahn: *Klin. Woch.*, 1924, xxi, 920. ¹⁹ Botelho: *ref. Wilbouchewitch, C. R. Soc. de Biol.*, 1922, lxxviii, 1339. ²⁰ Fry: *Bull. de l'Assoc. française pour l'étude du Cancer*, 1925, 14, p. 52. ²¹ Hoffgaard: *Muench. med. Woch.*, 1924, lxxi, 231. ²² Weil and Braun: *Wien. klin. Woch.*, 1908, xxi, 650. ²³ Roffo: *Les Néoplasmes*, 1924, iii, 18. ²⁴ Dreyer and Ward: Medical Research Council, Special Report Series No. 73, p. 72. ²⁵ Browning and Mackenzie: *Recent Methods in the Diagnosis and Treatment of Syphilis*, Constable, 1924, second edition. ²⁶ Loeper and Tonnet: *C. R. Soc. de Biol.*, 1920, lxxviii, 1139. ²⁷ Wegievko: *Ibid.*, 1924, xci, 717. ²⁸ Kennaway: *Quart. Journ. Med.*, 1924, xvii, 302. ²⁹ Luden: *Canadian Med. Assoc. Journ.*, 1922, xii, 147. ³⁰ Reeves: *Lancet*, 1924, i, 726. ³¹ Falcitore: *Giorn. di Clin. Med.*, 1923, No. 4. ³² Volta and Benedetti: *Il Policlinico*, 1924, iv, Med. Session, p. 189. ³³ Holker, J.: *Journ. Path. and Bact.*, xxv, 1922, p. 281. ³⁴ Bauer, E.: *Zeit. f. Krebsforsch.*, 1925, vi, 20.

NASAL, AURAL, AND OTHER FOCAL SEPSIS AS A CAUSE OF NEURASTHENIA AND INSANITY.*

BY

PATRICK WATSON-WILLIAMS, M.D.LOND.,

CONSULTING SURGEON FOR DISEASES OF THE EAR, NOSE, AND THROAT,
BRISTOL ROYAL INFIRMARY.

THE purport of this communication is, first, to invite more widespread attention to sepsis as often the essential cause of functional psychoses, ranging from slight blunting of intellectual activity, through the more definite forms of abnormal taciturnity, inability to concentrate, unsociability, and the varying syndrome of psychic abnormality grouped under the terms "neurasthenia" and "hypochondriasis," to the graver manifestations of unsoundness of mind accompanied by suicidal impulse, delusional insanity, mania, and ultimately dementia praecox; and, secondly, to lay stress on the import of the fact that the sepsis is most frequently what is conveniently termed "focal," and "latent" or "cryptogenic."

Many years ago I held a post of resident clinical assistant at Winsor Green Asylum, where there were about 600 inmates, and to which the acute cases occurring in and around Birmingham were mostly sent. One could not fail to be impressed with some of the acute maniacal cases which were apparently cut short by a blue pill and active treatment of the gastro-intestinal tract. It is now generally accepted that general paralysis of the insane is a legacy of syphilis, and alcohol and toxæmia of gastro-intestinal origin are recognized as causes of functional psychosis of toxic origin. The number of cases of insanity due to apical dental sepsis and cured by extraction of offending teeth is growing rapidly. It is because it is not as yet widely known that focal sepsis of the nasal sinus, tonsil, and ear may cause functional psychosis and insanity that I desire to emphasize the importance of not overlooking such sources of mental trouble, without, however, contending that therein lie the most frequent source of sepsis causing mental aberration. Nor do I touch here on a subject I have elsewhere discussed²—the question of focal sepsis as a cause of mental delinquency and crime—although a fruitful subject if approached with judgement as a matter requiring careful deliberation. On defective delinquency and functional psychosis the splendid work carried on at the State of New Jersey's mental hospital, as described by Dr. Henry Cotton,³ is worthy of most careful attention.

In 1907 in the Long Fox Lecture¹ I stated that—

"All patients suffering from purulent sinusitis are liable to suffer from general toxæmic symptoms, toxic products reaching the blood either from direct absorption from the implicated sinuses or from

the gastro-intestinal tract, but lymphatic absorption of toxic matters, in the presence of channels of communication with the intracranial venous sinuses, must be held to account for the profound mental depression and difficulty in thinking clearly, almost amounting to slow cerebration, that is very frequently present in fronto-ethmoidal or spheeno-ethmoidal sinus suppuration, and sometimes a less marked degree, even in simple maxillary sinus disease. Many patients have come with drawn, sad expression and sallow complexion, expressing their weariness of life and a profound melancholia which was quite foreign to their natural state of mind—symptoms which have completely disappeared with the removal of their sinus disease; sometimes it has been difficult to recognize in the round-faced cheerful individual one who shortly before treatment was a haggard melancholic."

Up to sixteen years ago I had never met with a case where definite mental aberration preceded and was relieved by accessory sinus treatment, though many of my patients were the subjects of a psychosis and several had confessed to strong suicidal impulses. Shortly after this I encountered examples of definite insanity. The most remarkable in my experience occurred in 1911. It will be more useful to relate this one case at some length than cite many cases, some of which I have already published.¹⁰

A healthy-looking, well nourished man, aged 29, keen and good at outdoor games, shooting, and so forth, had a severe attack of influenza in 1901. In 1903 he had his first definite appendicitis. Appendicectomy was performed in 1907. This was followed by insomnia, for which he went on a voyage to Madeira.

Recovery seemed complete, and till early 1911 he was very bright, full of fun, and keen on politics and his work. (? Acquired relative immunization.) In February, 1911, he had influenza with bad headaches and general malaise, but not badly, so did not stay in bed, and went on till insomnia compelled him to give up work and go away for five weeks. He returned thin, miserable, and depressed. He tried to work for the next two months, his doctor advising this "as it would take his mind off himself"; but he found it impossible to continue. He was sent on a voyage, but "seemed in a mental mist all the time" and terribly depressed. (Later, he told me, he took a pistol with him intending to shoot himself.)

When I saw him in August, 1911, he looked ruddy and well nourished, but had been obliged to give up work, though he still played fairly good golf. But he said his memory was gone, he could not think, and his "mind was a blank." He had post-nasal discharge, which he had been told was of no importance. I found fetid pus in the one antrum explored. He travelled to Bristol, wandering up and down the train corridor the greater part of the night, and could not sit still. He declared he was "devil driven," and that "it was a sin to have an operation, because he was dead." Mr. Stock gave an anaesthetic on September 23rd, 1911, and I opened both maxillary antra, which contained fetid pus, and from the frontal sinuses pus was washed out. Lavage of frontal sinuses and antra was continued daily, but his delusions became more insistent. He refused all food because "he was dead." The matron of the home said she feared to keep him because he was insane, and ought to be under certificates; but, instead of certifying, with his parents' consent I operated again on October 29th, assisted by Mr. A. J. Wright, performing my radical osteoplastic flap-frontal sinus operation. The frontal sinuses were unusually large and the mucosa much diseased. Cultures from the antra and frontal sinuses yielded pure *Staphylococcus aureus* (J. Walker Hall).

For several weeks he refused all food and had to be forcibly fed by male attendants; he tore off his dressing when he got the chance.

He went home on December 3rd, with the operation wounds healed, mentally much improved, but still subject to delusions of having "committed unpardonable sins"; even then at times he refused food and declared he was dead. His delusions continued for some months, but they became progressively less dominant, and less than a year later he was writing sensible letters full of interest in life. Very soon after that he was back at his work; since then he has remained in excellent health and has occupied a responsible position for a period of twelve years.

I would emphasize the following points in this case:

(a) The long history of sepsis following influenza in 1901, with gastro-intestinal infection and appendicitis probably of nasal origin—a by no means infrequent occurrence.

(b) After a period during which the infection was inhibited by auto-immunization a second attack of influenza caused a relapse and the nasal sinus infection caused profound mental disturbance—melancholia with resisted suicidal impulses. Despite a prolonged holiday and a voyage, he could no longer regain the power of work, and he then developed delusions.

(c) He did not lose his delusions for several months after disinfection and detoxication, but after his general health had been restored he regained all his old mental activity and geniality and has continued to occupy a position of responsibility without intermission.

(d) The patient was undoubtedly insane, and apart from the symptoms related, the letters in my possession that he

* This paper and the three papers that follow it were read before the Bath and Bristol Branch of the British Medical Association at the University of Bristol, May 27th, 1925.

wrote prior to operation on his frontal sinuses suffice to demonstrate his delusional insanity. Had he been certified as insane it would have been a matter of great difficulty for him to regain the confidence of his bank directors.

(e) Bacteriological examination of the nasal sinus contents demonstrated pure culture of *Staphylococcus aureus*, so often regarded as of little import. Teeth and tonsils were not involved; his previous appendicectomy had apparently removed the only other involved focus of infection, making his a pure nasal infection case.

Let me refer very briefly to one example at the other extreme—the very mild type of chronic nasal sepsis.

A man, aged 56, complained of deafness with post-nasal catarrh, never severe, but of twenty-seven years' duration. He had been treated off and on during that long period, but latterly had been getting depressed, lessened mental alacrity causing difficulty in carrying on his work; he complained also of general muscular weakness. Exploration of his nasal sinuses revealed no obvious pus, but Dr. Todd, who examined the extracted contents, found, together with polymorphonuclears, streptococcus and Friedländer's pneumobacillus in the right sphenoidal and posterior ethmoidal cells, the other sinuses being sterile. The infected sinuses were opened and drained and he soon felt stronger, increased in weight, and, as he put it, brighter than he had ever done for years past.

It is certainly not intended to suggest that all patients whose tonsils, sinuses, ears, or teeth are mildly infected must be operated on, or that they are the subject of functional psychoses, any more than is the individual who is somewhat melancholic because his liver is sluggish or he is constipated. Few individuals are there who, in the course of common colds or influenzal attacks and so forth, have not been the subject of transient nasal sinusitis; fortunately the great majority of such infections clear up spontaneously by auto-immunization, or with the appropriate measures of ordinary medication; while every medical practitioner knows that rest, feeding up, change of air, and so on are very often all that is required to obtain complete cure.

It is cases of chronic sepsis with mental depression and neurasthenic symptoms which do not yield to ordinary medicinal measures, or that seem to clear up only to recur, that we need to recognize, and to associate the functional psychosis with the physical abnormalities that may be the underlying and removable cause.

It is necessary to emphasize the "latent" or hidden character of the focal sepsis in many patients who nevertheless are the worst sufferers from chronic toxæmia. First, there is often an absence of pain or other symptoms directing attention to the real source of infection; and, secondly, it is significant that patients with chronic copious purulent discharge but seldom suffer from toxæmic symptoms as compared with those with little discharge, and that often non-purulent to the naked eye. Many years ago I suggested "that in the numerous polynuclears in the one case, as contrasted with the paucity of polynuclears in the others, we have an explanation of the remarkable differentiation to which I have drawn attention."³ Subsequent experiences have borne out the truth of this, and in the latent character of apical dental sepsis, which often enough can only be detected by radiograms, we have a further exemplification of the cryptogenic character of some of the most potent sources of focal sepsis.

If it be recognized that owing to the existence of focal sepsis a considerable percentage of patients are miserable, the subject of neurasthenia or of functional psychosis of grave character, rendering their lives more or less useless to themselves or a source of danger to the community, it is hardly necessary to urge the duty of ferreting out these septic foci—which involves regard to the gastro-intestinal tract (for example, for evidence of colitis, enteric ulceration, appendicular infection); to the genito-urinary tract, the teeth, fauces, nasal sinuses, the ear, and so forth.

In such investigations we must often invoke the aid of radiology, of analysis of gastric contents and of the faeces. How many asylums to-day have such essential resources at their disposal? Yet within a few years I believe that with each one will be actively associated a consulting physician, surgeon, dentist, radiologist, laryngologist, and pathologist. Let us cast no stones, for hitherto we as a profession have barely recognized the symptoms which should call for the elimination of focal sepsis as the *fons et origo* of some patients' afflictions in whom the ostensible diseases are

merely the end-results. Of this we may feel assured—sepsis is a far deadlier foe to civilized races than tuberculosis.

REFERENCES.

- ¹ P. Watson-Williams: Long Fox Lecture, 1907, On Suppurative Disease in the Nose and Ear, *Bristol Med.-Chir. Journ.*, 1908.
- ² Idem: *Rhinology*, 1910, Longmans, Green.
- ³ Idem: Operation for Drainage of the Sphenoidal Sinus, *Bristol Med.-Chir. Journ.*, 1916.
- ⁴ Idem: Latent Sinusitis in Relation to Systemic Infections; Introduction to a Discussion at the Royal Society of Medicine, *Journ. of Laryngol., Otol., and Rhinol.*, July, 1919, p. 233.
- ⁵ Henry A. Cotton: The Role of Focal Infections in the Psychoses, *New York Med. Journ.*, March 8th and 15th, 1919.
- ⁶ P. and E. Watson-Williams: Posterior Ethmoidal Cell Exploration, *Journ. of Laryng., Otol., and Rhinol.*, October, 1921.
- ⁷ P. Watson-Williams: Rheumatoid Arthritis due to Infection of the Nasal Accessory Sinuses, *BRITISH MEDICAL JOURNAL*, January 21st, 1922.
- ⁸ Idem: Les Sinusites sans Pus, *Arch. Internat. de Laryngol.*, April, 1922.
- ⁹ Burdette G. Lewis: *The Winning Fight against Mental Disease*, December, 1922.
- ¹⁰ P. Watson-Williams: The Diagnosis of Headaches and other Nervous Manifestations of Nasal Origin, *Lancet*, 1922, i, p. 311.

ON AURAL FOCAL SEPSIS AS A SOURCE OF NEURASTHENIA AND INSANITY.

BY

E. WATSON-WILLIAMS, M.C., CH.M.,

CLINICAL LECTURER IN LARYNGOLOGY AND OTOTOLOGY IN THE UNIVERSITY OF BRISTOL.

It has long been recognized that disease of the ear may lead to mental disturbance more or less severe. Even so simple a matter as removing a plug of wax has been recorded to have cut short an attack of mania. As regards aural suppuration, attention in the past appears to have been directed in this connexion to deafness, tinnitus, pain, vertigo, and intracranial complications. In short, only disease of the organ of hearing, and local complications, seem to have been considered. With such I am not here immediately concerned.

Many instances of the association of one or more of these conditions with mental trouble have been recorded. Tinnitus especially is noteworthy; some writers even go so far as to say that most cases with hallucinations of hearing ought to be classed as suffering from delusions, there being some definite aural lesion to account for their symptoms. Vertigo is well known to induce definite neurasthenia in some cases. We ought to be careful that disinclination to mix in traffic, or a shrinking from work, say, as a stoker, or on a scaffolding, is not incorrectly ascribed to neurasthenia, when really due to laudable prudence. I have had the good fortune to see both this false neurasthenia and a true neurasthenia disappear when suitable treatment has relieved the vertigo. A few cases of intracranial abscess have been published where treatment has relieved a definite psychosis, but too many of the records showing this association are from autopsies, no treatment having been instituted. I may summarize the situation by recalling the plea of Torretta,¹ which, in 1901, he added to a record of a case successfully treated—that every patient suffering from mental change should have the ears examined. In the same year Dr. Bernard Hollander wrote:²

"By these cases one is forcibly taught the lesson that an apparently slight and unimportant ailment, such as a running from the ear, which is apt to be looked upon as an inconvenience rather than a disease, may prove very dangerous."

TOXAEMIA.

Putting aside these local manifestations, I shall omit all cases in which tinnitus, vertigo, or intracranial complications were known to be present, and all cases in which there was not a family history free from mental taint; and shall confine myself to those instances in which surgical treatment of aural sepsis brought relief from the mental symptoms.

Towards the close of last century much attention seems to have been directed to the question whether toxæmia was a factor of importance in the causation of insanity. Among the records of many discussions, perhaps the most interesting is that at the Medical Society of London on May 11th, 1896, opened by Dr. A. M. Hamilton of New York, on "The connexion of autotoxis with certain forms of insanity." The idea that chronic septic absorption could