THE LONDON SKULL.

BY

G. ELLIOT SMITH, M.D., Litt.D., D.Sc., F.R.C.P., F.R.S.,
PROFESSOR OF ANATOMY, UNIVERSITY COLLEGE, LONDON.

It is just a century since Charles Lamb was pensioned and left East India House in Leadenhall Street; and the place where the gentle Elia, in the intervals of his clerical labours, conceived his imperishable essays is now occupied by a vast excavation, 50 feet deep, in which the new building for Lloyd's is being erected. For 14 feet below the present level of the street remains of the former Roman occupation are visible; lower down 12 feet of undisturbed "dirt" rest upon 14 feet of river gravel, below which (that is, 40 feet below the street level) is the blue clay. Last March the ulna of a woolly rhinoceros (*Rh. antiquitatis*) was found in the blue clay, in which a human skull has recently been found. There can therefore be no doubt of the remote antiquity of the original owner of the skull, who was a contemporary of the woolly rhinoceros and the mammoth in what is now the heart of the City of London. But as the Third Terrace of the Thames, in which it was found, is supposed not to have been deposited until after the disappearance of Neanderthal man from Europe, this is presumptive evidence that the fossil is unlikely to belong to the extinct species (*Neanderthalensis*), and that it cannot be earlier than the Aurignacian period. This skull of the Upper Palaeolithic period, however, represents the earliest human remains yet revealed in the City of London.

The fragment (Fig. 1) includes most of the occipital and left parietal bones, and part of the right parietal. The position of the groove for the middle meningeal vein (Fig. 3, B) indicates that the point of bone in front of it reaches to within a millimetre of the coronal suture. The closing of such parts of the sagittal suture as are represented in the fossil and of the lambdoid suture (excepting its lowermost part (Fig. 1) and the endocranial suture in the neighbourhood of the lambda (Fig. 2)) suggests an age of between 40 and 50 years. The exceptional smoothness of the skull and the faintness of the muscular impressions in a person of middle

age indicate the probability that the sex was female. The reasons for claiming that "the lady of Lloyd's" was left-handed are discussed in a later paragraph.

The rough sketch (Fig. 1) indicates the exceptional flat-

ness and distinctive outline of the cranium in the median line. The profile of the cerebral hemisphere (indicated by the broken line) is identical in size and shape with those of the two women of the Neanderthal species; but the flattening of the cerebellum in the latter (see the lines of the Neanderthal cerebellum and occiput at N) in comparison with its fullness in the London fossil mars the completeness of the resemblance to the Neanderthal type.

But the likeness of the cerebral part of the endocranial cast in the London and Neanderthal women involves not only the contour of the outline but also the modelling of the surface. In the endocranial casts of *Pithecanthropus*, the Piltdown skull, and the Neanderthal women (La Quina and Gibraltar crania) there is a prominent boss in the region of the posterior part of the middle temporal convolution, corresponding to the area so intimately concerned in modern man with the acoustic symbolism of speech. In modern man the expansion of the surrounding cortical areas raises them to the level of this precociously developed eminence so that the boss is no longer

apparent. In the cast of the London skull the temporal eminence (Fig. 3, T) is still present; and the parietal area (P) above it almost as flattened as it is in the Neanderthal casts. In fact, there is a striking resemblance

to the configuration of the La Quina cast. The maximal breadth of the intracranial cast is intermediate (136 mm.) between the measurements of the La Quina (130) and the Gibraltar casts (140). It is obviously impossible to measure the cranial capacity from this fragment; but by comparison with the Gibraltar skull (itself imperfect) one can safely say that the cubic content of the London cranium cannot have exceeded 1,200 c.cm. by more than 50 c.cm. at the most.

Although most of the crania of Upper Palaeolithic men are flatter than those of the average modern man their height approaches more nearly to the latter (Fig. 1,

M) than to that of the London skull, which in this respect agrees with the Neanderthal standard. In other words, the London skull more nearly resembles the Neanderthal type than any other of the Upper Palaeolithic specimens does. The likeness is so real as to raise for serious con-

FIG. 1.—Tracing of the median contour of the London skull (thick line) contrasted with the corresponding part of a modern British woman's skull (M). The dotted lines at N reveal the only difference of the contour of a Neanderthal woman's (La Quina) skull, the cerebellum being much flatter and the occiput adapted to the altered form.

FIG. 2.—Drawing by Mr. A. K. Maxwell of the interior of the London skull viewed from in front. L, crista lunata; F, fossa (cort.) striatae; S, the small fossa striatae of the left side.

sideration the possibility whether in Britain a stray member of the Neanderthal species may not have survived after the Mousterian phase of culture had been superseded by the Aurignacian.

Apart from the age of the fossil (that is, its deposition in beds later than that associated with Neanderthal man), the form of the cerebellum and its bony receptacle (Fig. 1), the thinness of the cranium, and the slightly greater fullness of the parietal area suggest that "the left-handed lady of Lloyd's" was an exceptionally primitive member of the species *sapiens* rather than a belated Neanderthaloid. Though the London cranium is no thinner than the La Quina specimen, this is exceptional for a member of the Neanderthal series. Hence it must be given due weight as a reason for excluding the Londoner from the latter species.

Without the front of the cranium or the face it is not possible definitely to exclude the possibility that this skull may belong to the Neanderthal species; but I think the probabilities are against such a conclusion, both on anatomical and geological grounds. This uncertainty does not diminish the interest of the London skull; for the enigma of its affinities emphasizes how closely a primitive type of *Homo sapiens* may approach the Neanderthal species.

Reference has already been made to the probability that "the lady of Lloyd's" was left-handed. Some explanation will be demanded in justification of a claim so confidently made.

Twenty years ago, when investigating the occurrence of the so-called Affenspalte (sulcus lunatus) in the human brain, I called attention to the fact that it was often found as a large semilunar furrow on the left hemisphere (Fig. 4), but rarely on the right. Moreover, I recorded the observation¹ that the cortical territory, for which I coined the term "area striata," was often raised into a prominence behind the sulcus lunatus (as in the Egyptian brain shown in Fig. 4). Corresponding to this prominence there is often a deep depression on the left side of the occipital bone (fossa corticis striatae), surrounded by a ridge (crista lunata) corresponding to the sulcus lunatus. In the winter of 1907-08 Professor F. Wood Jones collaborated with me in the attempt to discover the significance of the occasional reversal of the asymmetry usually found in the cerebral hemispheres.* In his field notes on the skeletons found in the course of the archaeological survey of Nubia he recorded whether the right

or the left humerus was the longer, and on this basis inferred, in the case of each skeleton, whether the individual was right-handed or left-handed. Using these data I discovered that when the deep fossa striatae and the extensive crista lunata were on the right side, and not on the left, it afforded a reliable criterion of left-handedness. The London skull affords an excellent illustration of this

phenomenon (Fig. 2). On the right side (left of the figure) there is a prominent and extensive crista (L) and a deep fossa striatae (F), whereas upon the left hemisphere there is only a very diminutive fossa (S). This is a reversal of the customary arrangement (seen in Fig. 4), and, I believe, affords definite evidence that the earliest known Londoner was left-handed.

In my earlier memoir on the asymmetry of brain and skull (op. cit., 1907) I laid undue emphasis on the arrangement of the venous sinuses, when I suggested that the common deflection of the superior longitudinal sinus into the right lateral sinus was causally related to the slighter backward projection of the right hemisphere. In fact I assumed that this arrangement of the largest venous channel was determined by purely mechanical factors, the asymmetrical condition of the occipital poles allowing more room for the blood to flow freely to the right. This tentative explanation may still have some measure of justification. But it is evident—as the condition revealed in the London fossil indicates—that other factors must play

a part in determining the asymmetry of the cranial sinuses, and that the reversal of their normal asymmetry is not necessarily an indication of left-handedness. Although the deflection of the superior longitudinal sinus into the left lateral sinus is usually associated with left-handedness the correlation is not invariable. In the London skull, for example, the groove for the superior longitudinal sinus splits at the torcular herophili into two lateral sinuses of almost equal size (Fig. 2), as is usual in the apes; but of the two the right is slightly larger than the left. Thus, while the brain conforms to the condition distinctive of left-handedness, the right lateral sinus is larger than the left, as is usual in right-handed persons. When the evidence of

asymmetry is thus conflicting, examination of a series of crania (and the arm bones of the skeletons) reveals that the brain and not the venous sinuses is the real index of right- or left-handedness.

I have to express my gratitude to Mr. Warren R. Dawson for calling my attention to the interesting fossil discussed in these notes, and to the Committee of Lloyd's for its generosity in presenting it to University College.

REFERENCE.

¹ *Anat. Anzeiger*, 1907, Bd. xxx, p. 574.

FIG. 3.—Left side of the London endocranial cast. B, groove for middle meningeal vein; T, temporal boss; P, poorly developed parietal area.

FIG. 4.—Posterior aspect of an Egyptian brain viewed obliquely from the left side to show on the left hemisphere, but not on the right, the large sulcus lunatus and prominent button of area striata behind it (compare Fig. 2).

* The question of the cranial signs of right- and left-handedness were discussed by me more fully in a paper read in the Section of Anatomy at the Annual Meeting of the British Medical Association in 1908. An abstract of the paper was published in the *JOURNAL* of August 29th, 1908 (p. 596).