be so; Barr found a history of giddiness in 14 per cent. of his cases and Habermann in 16 per cent. The fact that I found it in only 10 per cent. may quite well be attributed to my inclusion of such a large proportion of young men in whom the upper tone limit was scarcely if at all affected.

But it must not be assumed from the juxtaposition in the opener's sentence quoted that the lowering of the upper tone limit has anything to do with the incidence of vertigo. The latter must be due to coincident injury to the vestibular apparatus. The experimental work referred to revealed no *post-mortem* change in the end-organs of equilibrium until Yoshii tried detonation instead of electric bells, when he obtained immediate clinical results, the animal showing nystagmus, rhythmic movements, and inco-ordination, as well as *post-mortem* signs of injury. When one thinks of how much the noise of boiler-making resembles that of repeated detonation, especially when the "holder-on" is inside the boiler with riveters hammering on the outside, one can understand the frequent incidence of vertigo.

Lastly, I agree with the opener that the mass of evidence is in favour of the view that the injury to the cochlea is conveyed by air conduction, not by bone conduction. Middle-ear defects have no part in the causation. Indeed, the very opposite is the case; any middle-ear defect at all, by diminishing the impact of the stapes against the oval window, will prevent the onset of inner-ear deafness. This was exemplified in 6 of the 96 ears I examined. In these either otorrhoea or a large perforation was present, and the signs of nerve deafness were absent. In 3 others, however, in which there was a history of otorrhoea of short duration, and in 2 with small perforations, nerve deafness had supervened, doubtless because the sound-transmitting apparatus was practically unaffected. Von Eicken and Hoessle in their experiments also found that the induction of middle-ear suppuration or the removal of the incus prevented the onset of changes in the cochlea in the ear affected. This point, of course, is all-important as regards prevention. Wittmaack's advocacy of a rubber mat was based on the idea that the injury was conveyed through the bone. Meatal plugs of wax or vaselined wool are the only hope, and if workmen could be persuaded to use these regularly there is no doubt that boiler-maker's deafness would be much less prevalent.

GENERAL DISCUSSION.

Mr. E. B. WAGGETT confined his remarks to the subject of chronic otorrhoea in the army, and excluded the chronic sequelae of the influenza epidemic of 1918, of which he had no special knowledge. The subject, he said, must be considered from two distinct points of view: (1) as occurring in a nation in arms for the purpose of a war; (2) as affecting the regular standing army.

1. He considered that the statistics of the great war gave a wholly misleading impression, and he believed that the very heavy figures concerning hospital entries for chronic otorrhoea and aural, nasal, and throat operations were to be explained on psychological rather than surgical grounds. Chronic otorrhoea formed a convenient excuse for the soldier desiring a holiday at the base; moreover, many men who had neglected a mild otorrhoea in civil life very naturally seized the opportunities for treatment afforded by the Army Medical Service. Very few medical officers had any thorough knowledge of aural diseases, and, being unable to distinguish the different causes of chronicity in otorrhoea—namely, reinfection from the meatus, the nasopharynx, and the mastoid cavities respectively—they very properly relied upon the impressive warning of the "slumbering volcano," and sent cases down to the base indiscriminately. At the base such cases must frequently have come under the charge of medical officers influenced by a desire to improve their operative technique. The speaker based these beliefs upon his personal experience of four years' field ambulance service in the front line. Thanks to the co-operation of his brother officers in the division, and of intelligent and cleanly medical orderlies, he had found it necessary to send to the base precisely one case of otorrhoea during the whole period of the war. He considered that the status of a chronic otorrhoeic was,

under the conditions at the front as he had witnessed them, safer than in normal unorganized civil life. Acute cases had without exception cleared up in the field ambulance. Oto-laryngology had now become a compulsory subject for medical qualification. If teachers would concentrate upon the public aspects of the subject, the next great war would be served by medical officers capable of furnishing statistical figures of a totally different magnitude.

2. The regular standing army presented a different problem, which must take account (a) of the definite fact that, for cogent reasons, the Army Medical Department could not supply competent aurists in sufficient numbers; (b) that a regular soldier must at all times be fit for field service in any war area; (c) financial considerations arising out of discharge during the six months following enlistment, etc.; (d) deafness, increasing to the degree of disability, during prolonged service. For these reasons the problem had to be met and dealt with at the time of recruitment. The War Office had employed him to make recommendations on this point. He understood that these had, in the main, been accepted by the medical recruiting authorities, and if embodied in full in the new recruiting regulations they would, he was confident, satisfy aural surgeons. These regulations were not yet issued, and it was possible that the Adjutant-General's office might have found it necessary, on military grounds, to amend his recommendations.

Dr. J. KERR LOVE (Glasgow) said that he would draw a distinction between true middle-ear disease, in which the hearing was sometimes better in a noise, and boiler-maker's deafness, an internal ear condition in which the hearing was not better in a noise. As an example of the former, a friend of his, sitting along with her husband in the back seat of a motor car, heard what was going on in the front seat better than her husband, who had perfect hearing. The speaker asked Mr. Westmacott whether he had seen what he (the speaker) had called the blue membrane of miners, a condition he had described twenty years ago, and which was due to the passing up by the Eustachian tube of coal dust which was deposited on the lining of the middle ear.

Dr. T. B. JOBSON (Guildford) said that in 1915 he had the opportunity of making some observations on the effect of prolonged gunfire on the healthy ear. He examined fifty German soldiers who had recently been exposed to gunfire. Cases with any history of previous ear disease, abnormal membranes, and naso-pharyngeal disease were excluded. He found that about 80 per cent. of these men showed a slight diminution of acuity of hearing both for bone conduction and for aerial conduction. Examination of these cases after an interval of three to six months showed that the hearing had in almost all cases returned to normal. This corroborated Mr. Faulder's remarks on the powers of resistance to gunfire of the normal healthy ear. The number of cases of serious ear injury, exclusive of wounds, due to gunfire which came under observation was small—unexpectedly small.

DIAGNOSTIC EXPLORATION OF THE NASAL SINUSES BY THE AUTHOR'S SUCTION SYRINGE METHOD.

DEMONSTRATION AT THE BATH EAR, NOSE, AND THROAT
HOSPITAL

BY

PATRICK WATSON-WILLIAMS, M.D. LOND.,

Consulting Surgeon for Ear, Nose, and Throat Diseases, Bristol
Royal Infirmary.

BEFORE proceeding to demonstrate the technique on cases requiring investigation, I would point out that the bacteriological investigation of the nasal accessory sinuses affords valuable guidance both for accurate diagnosis and for treatment. Where the discharge of pus is copious the value of the method lies in the means afforded for rapidly determining which sinuses are healthy and which involved. But few conditions illustrate the value of the suction syringe method more than those cases without purulent discharge—for example, patients in whom signs of canalicular optic neuritis point to the sphenoidal sinuses or a posterior ethmoidal cell as being the probable source of infection,

though even careful anterior and posterior rhinoscopy fails to demonstrate escaping discharge; in other words, cases in which the nasal findings by ordinary methods are negative and yet the progressive optic nerve lesion renders accuracy of diagnosis of a source of infection of grave import. Equally does this apply to cases of arthritis in which a nasal source of infection is suspected.

It is, however, in determining and exactly localizing the focus of infection when present in the sphenoidal sinuses or posterior ethmoidal cells that the suction syringe has proved most helpful. Investigations in the *post-mortem* room which I carried out in 1896 with the co-operation of Mr. S. V. Stock led me to introduce a method of puncturing the anterior wall of the sphenoidal sinus and sucking any contained discharge back into the attached sterile syringe, instead of syringing out the sinus by the natural opening. Subsequently I applied the same method to the posterior ethmoidal cells, and thus with the suction syringe one can determine the presence or absence of active infective processes in either the maxillary antrum, sphenoidal sinuses, and anterior or posterior ethmoidal cells on either side.

Let us take, for example, a patient threatened with loss of vision from canalicular neuritis, suspected as nasal in origin. Having explored various sinuses and found them sterile, it is necessary to make certain that no infected sinus has been "missed," because the failure to locate exactly the infected focus may cost the patient his eyesight, if nothing worse. Further, if some of the sinuses have been proved to be infected, it does not follow that they are the chief or sole source, or even the source at all, of a canalicular neuritis, unless one can be certain that the other sinuses, which might cause the neuritis, have also been explored and proved sterile. Briefly, the method employed consists in attaching to a syringe a cannula suitable for entering the suspected sinus and sucking up the discharge, which is then placed in a sterile bottle and submitted to bacteriological investigation. It is usually helpful to inject a small quantity of sterile water from the syringe into the particular sinus and then suck it back with the contained discharge. By such means the bacteriology of the maxillary antra, the ethmoidal cells, and the sphenoidal sinuses can be determined individually. By centrifugalizing and making stained films of the fluids thus obtained the presence of polymorphonuclear leucocytes and often of phagocytosis, as well as the organisms present, may be noted prior to culture of the organisms by usual methods. While similar methods are often applicable to the frontal sinuses, the resort to skiagrams seems to afford all the information desired in these cavities. It is the systematic exploration of doubtful cases that has proved so helpful to the author, because by exploring both antra, the ethmoids, and the sphenoidal sinuses, it so frequently happens that those believed to be infected have proved normal, while those sinuses which were not suspected have often yielded undoubted proof of infected and purulent discharge.

Diagnostic exploration by the suction syringe can be performed under local anaesthesia by cocaine, tutocain, or eucaine, etc., in all but children or highly nervous adults for whom a light general anaesthetic is desirable. With the patient then lying flat on the back, a cotton-wool carrier dipped in 10 to 12 per cent. solution of cocaine hydrochlorate, or of tutocain or any efficient local surface anaesthetic, is carried well up and back into the middle or superior meatus, according to the route chosen, and in the space of three or four minutes the patient is ready for the exploratory procedure.

For the maxillary antrum a stout curved hollow needle is used, with a bore of 2 mm., so that even thick sticky pus can be drawn up into the syringe. While it may be entered through the inferior meatal wall, it can be passed with greater ease through the middle meatal wall in the region corresponding to the processus uncinatus and with no pain (as it traverses only very thin bone or often only mucous membrane), and this is the route always chosen by the author. For the sphenoidal sinus a straight *blunt* trocar and cannula, measuring $4\frac{1}{2}$ inches in length, is used, and the same is used for posterior ethmoidal cells. With the patient supine, the blunt trocar and cannula are made to enter the sphenoidal sinus through the thin

anterior wall—that is, through the sphenoidal turbinate bone, avoiding, if possible, entering through the natural ostium, because it is then difficult to suck out the contents of the sinus.

For the posterior ethmoidal cell exploration the sphenoidal trocar and cannula are most readily entered through the middle meatus. The trocar and cannula are passed into the top of the middle meatus, between the ethmoidal bulla and the middle turbinal (or sometimes through the bulla), and slipped upwards and backwards until the point rests high up on the posterior oblique attachment of the middle turbinal. Just before puncturing the cell wall the direction of the trocar and cannula is rendered less oblique by raising the instrument, and, if necessary, the tip of the nose, and it is now made to take a slightly outward direction. The instrument is then gently pressed through the thin anterior wall of the posterior ethmoidal cell or cells, so as to enter the posterior cell just outside the thicker bone which corresponds with the attachment of the middle turbinate bone. Care should be taken not to fracture the outer wall (into the back of the orbit), nor to direct the point too much upwards through the roof into the cranium. If this route is followed, and the pressure made somewhat outwards and but slightly upwards, there should be no risk of accident in skilled hands. When the point has fully entered the cell, the cannula is held steadily *in situ*, while the trocar is withdrawn and the suction syringe, containing about 3 c.cm. of sterile distilled water, is attached as in exploring the sphenoidal sinus.

A slight movement of the piston to and fro serves to wash the water into the sinus and suck it back—it may not go quite so easily as with the sphenoidal sinus, probably from a difference in the size of the cell's natural ostium, which serves as a bung-hole. As it is essential to use a sterile instrument and sterile syringe for each sinus explored, the author has six syringes, four sphenoidal trocars and cannulas, and two antral curved needles at hand so as to avoid unnecessary delay in sterilizing any of the used instruments during the course of exploration. The whole procedure, apart from cocainization, usually occupies two minutes only, and often less in cases under general anaesthesia.

Having thus entered a sinus cavity, the blunt trocar is withdrawn and the syringe attached to the proximal or projecting end, and the sample of the investigated sinus content sucked into the syringe. If much pus is lying in the sinus, it may appear in the syringe in ropes, or as very bloody mucus, leaving one in no doubt as to the sinus being infected. In other cases the fluid extracted may be perfectly clear and watery, proving sterile on culture. Other samples obtained may be suspicious only, and doubtful until submitted to film and culture examinations, which may either prove negative or definite phagocytosis; abundance of polymorphonuclears and pyogenic cocci may afford conclusive evidence of active infection.

The posterior ethmoidal cells are similarly explored by the blunt trocar and cannula entered either above and internal to the middle turbinated body, or, as I usually find much simpler, by piercing the posterior ethmoidal cell walls beneath the attachment of the middle turbinated body to the outer wall of the nasal passage.

With the maxillary antrum, ethmoidal cells, and sphenoidal sinuses explored thus through the right and left nasal passages, at least six samples have been obtained for investigation by the pathologist, which, if the anatomical arrangement of the sinuses be normal, should yield exact and reliable information as to the normal or infected condition of each sinus. Unfortunately, the sphenoidal sinuses and ethmoidal cells are often most irregularly developed. For instance, one sphenoidal sinus may be very large and extend right across the mid-line, pushing the sinus of the other side, which is correspondingly small, so far outwards that the exploring cannulas passed backwards through the right and left nasal passages enter the one large sinus, the other small one being altogether missed. This is not difficult to discover if, with both cannulas in one sphenoidal sinus, water injected down one will come back through the proximal end of the other. If left in doubt as to whether one has entered the sphenoidal sinus instead of the ethmoidal cell, by again keeping both cannulas *in situ* and injecting

water down one it is quite easy to determine whether the two cannulas lie in the same or in separate cavities. Such precautions are by no means fanciful, and I have met with cases of canalicular optic neuritis where the sinus infection did not correspond with the retinal defects until further careful exploration revealed such irregularities as I have described, and which had very nearly led to the source of the very serious ocular defect being missed and the patient unrelieved.

LITERATURE.

- P. Watson-Williams: The Diagnostic Value of the Suction Syringe in Maxillary Antral Sinusitis, *Journ. of Laryngol.*, 1912.
P. Watson-Williams and E. Watson-Williams: Posterior Ethmoidal Cell Exploration, *Journ. of Laryngol.*, October, 1921.
P. Watson-Williams: Les Sinusites sans Pus, *Arch. Internat. de Laryngol.*, etc., April, 1922.
E. Watson-Williams: The Diagnostic Value of Transillumination of the Maxillary Antra, *Journ. of Laryngol.*, 1912.

A NEW METHOD OF FITTING ARTIFICIAL LEG SOCKETS.

BY

E. MUIRHEAD LITTLE, F.R.C.S.,

CONSULTING SURGEON, ROYAL NATIONAL ORTHOPAEDIC HOSPITAL.

ONE of the greatest difficulties in fitting an artificial limb after amputations in general, and particularly after amputations of the thigh, has lain in the method of attachment of the false limb to the living stump. The ideal must be so to connect these two that the prosthesis shall partake of all movements of the stump in all directions as though it were part of it. Any movement, whether longitudinal or angular, which occurs between the socket of the artificial limb and the stump—what is called by mechanicians “lost motion”—is a source of weakness, gives rise to troubles in the skin of the stump, and is to be eliminated as far as possible.

In the case of amputation of the thigh the attachment has hitherto been made by means of straps or suspenders passing over the shoulders, and by means of belts around the pelvis made of steel, leather, or woven fabrics. When the patient with such an attachment assumes the erect position and rests some or all of his weight upon the limb, the pressure is transmitted almost entirely through the ischium, which bears upon the inner upper edge of the socket. Very little weight is taken on the thigh stump itself in most cases. When, as in taking a step in ordinary walking, the weight is transferred to the sound limb and the artificial limb is swung forward, the socket slides down upon the stump to a greater or less extent, until its movement is stopped by the suspenders or pelvic band. When weight is once more put upon the artificial limb, as the sound one comes forward, the stump descends into the socket until its movement is stopped by contact between the ischium and the top of the socket. When the patient sits down a similar descent of the prosthesis occurs, so that the artificial knee is more prominent than the living one. It has been the constant aim of the limb-maker to reduce this movement to a minimum, but it is nearly always to be detected. Out of thousands of cases of amputation of the thigh which came under my notice at Roehampton and elsewhere, only one was able to dispense with all straps and belts for the attachment of his artificial leg. This was an officer who had a very long stump, and who had found out that he could so jam it into the socket that the prosthesis would not drop off of its own weight. He thrust his stump into it as one might thrust a foot into a riding boot, and walked off without any suspender or belt whatever.

Some two years ago Mr. Ernest Underwood, a pensioner who had suffered amputation through the middle of the

femur, made himself a wooden socket, not shaped *secundum artem*, but actually turned in a lathe with three or four inch-wide annular grooves on the inside fitting the bare stump as closely as possible. Wearing this he was able to give up all suspensory attachments. His stump measures 9 inches in length from the tip of the great trochanter. It has an end posterior scar which is partly adherent.

From past experience I should have been strongly sceptical of the possibility that there could be any practical value in such a method of fitting, but last July Messrs. Charles A. Blatchford and Sons, Ltd., who had for some time been patiently working upon Mr. Underwood's rather crude idea, brought him, and four other amputees wearing sockets fitted on his principle, to see me. There was then no room for doubt that, in their cases at least, the method was a complete success. Since then I have been shown six other cases fitted with equal success, and have had opportunities of watching the progress of all the eleven cases, of which short notes are subjoined.

CASES.

1. D. J. A., aged 32. Amputation, September, 1916, of right thigh, 9½ in. below trochanter. Posterior scar. Seen August 27th and October 6th, 1925.
2. J. F. C., aged 42. Amputation of right arm and right thigh, in 1917, 14½ in. below the trochanter (he is a very tall man). Posterior scar on which he has formerly borne weight. Seen July 27th, September 1st, and October 6th, 1925.
3. S. D., aged 30. Amputation (in Germany) August, 1918, right thigh 10½ in. Posterior scar. Seen August 27th and October 16th, 1925.
4. H. G., aged 26. Amputation October, 1917, 12¼ in. End scar, adherent. Seen August 27th and October 6th, 1925.
5. E. G. G., aged 33. Amputation August, 1916, left thigh, 9½ in. Extensive posterior and internal scars. Seen August 27th and October 16th, 1925.
6. A. L. J., aged 28. Amputation February, 1918, left thigh, 13½ in. Posterior scar. Seen July 27th and October 16th, 1925. This patient is a record breaker for walking in an artificial limb. He walked the mile in 11 minutes 8 seconds, wearing a limb with shoulder control. In this limb he has done it in 9½ minutes, and 4 miles in 45 minutes.
7. J. J. Amputation September, 1916, right thigh, 12 in. Posterior scar. Seen August 27th and October 6th, 1925. A localized thickening of the derma, commonly called a cyst, which was caused by his former loose socket, is now disappearing.
8. T. W. P., aged 28. Right thigh, 10½ in. Posterior scar not adherent. Seen July 9th, 1925, when the following note was made: “Cannot be pulled off without very great effort, and when he relaxes his muscles; when he opposed I could not pull it off at all.” Seen again October 6th, 1925.
9. C. T., aged 42. Amputation December, 1917, right thigh, 13 in. Very conical stump, muscles much retracted; scar not adherent. Seen July 30th and October 19th, 1925.
10. E. U. Right thigh, 9 in. End and posterior scar partly adherent. Wearing a socket with groove of three turns of a spiral and one ring at bottom. Seen July 18th and October 6th, 1925.
11. J. N. Left thigh, 8 in. End scar not adherent. Seen October 6th and November 2nd. This patient is only now (November 2nd) just beginning to use the leg continuously. Although he has been used to a pelvic band and hip-joint, on account of his short stump, he walks better in this limb without that addition.

FIG. 1.—Diagrammatic drawing of a socket showing the groove and the valve.

The socket is made of duralumin, and experience has shown that a spiral groove of a little more than one turn is enough. The groove is 1½ in. wide and 1/8 in. deep. It is shown in the diagrammatic drawing (Fig. 1). The socket is made to fit without any sock as closely to the stump as is consistent with comfort, but the end of the stump is left free. This fit is so close that a partial vacuum is created when an attempt is made to withdraw the stump, and air rushes in with an audible noise as the socket leaves the stump. To facilitate withdrawal a spring valve is fitted on the socket, through which air may be admitted when it is desired to remove the leg. Attempts to pull