water down one it is quite easy to determine whether the two cannulas lie in the same or in separate cavities. Such precautions are by no means fanciful, and I have met with cases of canalicular optic neuritis where the sinus infection did not correspond with the retinal defects until further careful exploration revealed such irregularities as I have described, and which had very nearly led to the source of the very serious ocular defect being missed and the patient unrelieved.

LITERATURE.

- P. Watson-Williams: The Diagnostic Value of the Suction Syringe in Maxillary Antral Sinusitis, *Journ. of Laryngol.*, 1912.
P. Watson-Williams and E. Watson-Williams: Posterior Ethmoidal Cell Exploration, *Journ. of Laryngol.*, October, 1921.
P. Watson-Williams: Les Sinusites sans Pus, *Arch. Internat. de Laryngol.*, etc., April, 1922.
E. Watson-Williams: The Diagnostic Value of Transillumination of the Maxillary Antra, *Journ. of Laryngol.*, 1912.

A NEW METHOD OF FITTING ARTIFICIAL LEG SOCKETS.

BY

E. MUIRHEAD LITTLE, F.R.C.S.,

CONSULTING SURGEON, ROYAL NATIONAL ORTHOPAEDIC HOSPITAL.

ONE of the greatest difficulties in fitting an artificial limb after amputations in general, and particularly after amputations of the thigh, has lain in the method of attachment of the false limb to the living stump. The ideal must be so to connect these two that the prosthesis shall partake of all movements of the stump in all directions as though it were part of it. Any movement, whether longitudinal or angular, which occurs between the socket of the artificial limb and the stump—what is called by mechanicians “lost motion”—is a source of weakness, gives rise to troubles in the skin of the stump, and is to be eliminated as far as possible.

In the case of amputation of the thigh the attachment has hitherto been made by means of straps or suspenders passing over the shoulders, and by means of belts around the pelvis made of steel, leather, or woven fabrics. When the patient with such an attachment assumes the erect position and rests some or all of his weight upon the limb, the pressure is transmitted almost entirely through the ischium, which bears upon the inner upper edge of the socket. Very little weight is taken on the thigh stump itself in most cases. When, as in taking a step in ordinary walking, the weight is transferred to the sound limb and the artificial limb is swung forward, the socket slides down upon the stump to a greater or less extent, until its movement is stopped by the suspenders or pelvic band. When weight is once more put upon the artificial limb, as the sound one comes forward, the stump descends into the socket until its movement is stopped by contact between the ischium and the top of the socket. When the patient sits down a similar descent of the prosthesis occurs, so that the artificial knee is more prominent than the living one. It has been the constant aim of the limb-maker to reduce this movement to a minimum, but it is nearly always to be detected. Out of thousands of cases of amputation of the thigh which came under my notice at Roehampton and elsewhere, only one was able to dispense with all straps and belts for the attachment of his artificial leg. This was an officer who had a very long stump, and who had found out that he could so jam it into the socket that the prosthesis would not drop off of its own weight. He thrust his stump into it as one might thrust a foot into a riding boot, and walked off without any suspender or belt whatever.

Some two years ago Mr. Ernest Underwood, a pensioner who had suffered amputation through the middle of the

femur, made himself a wooden socket, not shaped *secundum artem*, but actually turned in a lathe with three or four inch-wide annular grooves on the inside fitting the bare stump as closely as possible. Wearing this he was able to give up all suspensory attachments. His stump measures 9 inches in length from the tip of the great trochanter. It has an end posterior scar which is partly adherent.

From past experience I should have been strongly sceptical of the possibility that there could be any practical value in such a method of fitting, but last July Messrs. Charles A. Blatchford and Sons, Ltd., who had for some time been patiently working upon Mr. Underwood's rather crude idea, brought him, and four other amputees wearing sockets fitted on his principle, to see me. There was then no room for doubt that, in their cases at least, the method was a complete success. Since then I have been shown six other cases fitted with equal success, and have had opportunities of watching the progress of all the eleven cases, of which short notes are subjoined.

CASES.

1. D. J. A., aged 32. Amputation, September, 1916, of right thigh, 9½ in. below trochanter. Posterior scar. Seen August 27th and October 6th, 1925.
2. J. F. C., aged 42. Amputation of right arm and right thigh, in 1917, 14½ in. below the trochanter (he is a very tall man). Posterior scar on which he has formerly borne weight. Seen July 27th, September 1st, and October 6th, 1925.
3. S. D., aged 30. Amputation (in Germany) August, 1918, right thigh 10½ in. Posterior scar. Seen August 27th and October 16th, 1925.
4. H. G., aged 26. Amputation October, 1917, 12¼ in. End scar, adherent. Seen August 27th and October 6th, 1925.
5. E. G. G., aged 33. Amputation August, 1916, left thigh, 9½ in. Extensive posterior and internal scars. Seen August 27th and October 16th, 1925.
6. A. L. J., aged 28. Amputation February, 1918, left thigh, 13½ in. Posterior scar. Seen July 27th and October 16th, 1925. This patient is a record breaker for walking in an artificial limb. He walked the mile in 11 minutes 8 seconds, wearing a limb with shoulder control. In this limb he has done it in 9½ minutes, and 4 miles in 45 minutes.
7. J. J. Amputation September, 1916, right thigh, 12 in. Posterior scar. Seen August 27th and October 6th, 1925. A localized thickening of the derma, commonly called a cyst, which was caused by his former loose socket, is now disappearing.
8. T. W. P., aged 28. Right thigh, 10½ in. Posterior scar not adherent. Seen July 9th, 1925, when the following note was made: “Cannot be pulled off without very great effort, and when he relaxes his muscles; when he opposed I could not pull it off at all.” Seen again October 6th, 1925.
9. C. T., aged 42. Amputation December, 1917, right thigh, 13 in. Very conical stump, muscles much retracted; scar not adherent. Seen July 30th and October 19th, 1925.
10. E. U. Right thigh, 9 in. End and posterior scar partly adherent. Wearing a socket with groove of three turns of a spiral and one ring at bottom. Seen July 18th and October 6th, 1925.
11. J. N. Left thigh, 8 in. End scar not adherent. Seen October 6th and November 2nd. This patient is only now (November 2nd) just beginning to use the leg continuously. Although he has been used to a pelvic band and hip-joint, on account of his short stump, he walks better in this limb without that addition.

FIG. 1.—Diagrammatic drawing of a socket showing the groove and the valve.

The socket is made of duralumin, and experience has shown that a spiral groove of a little more than one turn is enough. The groove is 1½ in. wide and 1/8 in. deep. It is shown in the diagrammatic drawing (Fig. 1). The socket is made to fit without any sock as closely to the stump as is consistent with comfort, but the end of the stump is left free. This fit is so close that a partial vacuum is created when an attempt is made to withdraw the stump, and air rushes in with an audible noise as the socket leaves the stump. To facilitate withdrawal a spring valve is fitted on the socket, through which air may be admitted when it is desired to remove the leg. Attempts to pull

off the limb by main force without opening the valve failed in all cases. The patient might be pulled across the floor by the artificial foot, but the limb seemed to be as firmly attached to the trunk as was its living fellow. It appears to me that it is not so much the vacuum as the close contact made by the metal socket and its groove which holds

FIG. 2.—Demonstrates the absence of angular or tilting movement between stump and socket.

the leg on. This opinion is supported by a recent experience of a case in which there was positive air pressure below the end of the stump, so that on opening the valve air rushed out with force enough to blow out a match.

It might have been feared that this close contact and the absence of ventilation would have a bad effect on the skin, or that the constant pressure might cause atrophy of the upper part and oedema of the end of the

stump. In practice I have found that such fears are groundless—at any rate in the cases of the eleven men in question. After wearing these sockets for periods varying from two months to two years the skin is in a healthier condition

than before, and the stumps have increased in circumference. The absence of friction between skin and socket eliminates the risk of sores and callosities, and the increased use as a lever of the stump, which is rendered possible by its close alliance with the socket, appears to have led to increased development of muscle and better nutrition of the stump. It is a commonplace of limb-fitting that the better a limb fits the less is its weight felt. Therefore it is not surprising that the wearers of these sockets make light of their disability.

As regards the question of ventilation, it seems likely that the partial vacuum may be dispensed with, in which case the lower end of the socket might be ventilated in the usual way. As regards the close-fitting metal socket, which in the nature of things allows no ventilation of the skin, it is to be remarked that this is made of thin metal and is suspended inside the metal thigh-piece, which is itself freely perforated. There is thus an air space all round it with free circulation.

Figs. 2 and 3 show how firm is the hold of the socket on the stump.

As far as I know, this method has not been tried in artificial arms, but there seems no reason to doubt that it might be adaptable to certain cases of amputation above the elbow.

FIG. 3.—Shows the strength of the adhesion between stump and socket. The whole of the lower man's weight is borne by the artificial limb.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

INTESTINAL OBSTRUCTION DUE TO GALL STONES.

In my comparatively limited experience I have seen several cases of this kind. One I operated upon in 1914. The patient was an old man, aged 78. He made a good recovery. The case is fully recorded in the *Liverpool Medico-Chirurgical Journal* of 1915. The calculus weighed 418 grains and measured 2 inches in length and $1\frac{1}{2}$ in breadth. The circumference measurements were $5\frac{1}{2}$ and 4 inches. The specimen is in the Royal College of Surgeons Museum in London.

Southport.

FRANCIS J. BAILDON, M.B.

STRICTURE OF THE VAGINA: DELIVERY BY CAESAREAN SECTION.

In May, 1925, Dr. Minett asked me to see a case for her at the Tsang Yuen Hospital.

The patient, aged 36, was pregnant for the second time. She came into hospital because her first labour, thirteen years earlier, had been difficult; she was delivered in a Chinese village; no instruments were used, apparently. The child was full term and was dead.

The external pelvic measurements were: interspinous 9 in., intercrystal 10 in., external conjugate $7\frac{1}{2}$ in. At the time she was about eight and a half months pregnant. There was a very rigid stricture, admitting only the tip of one finger, situated at the junction of the middle and lower thirds of the vagina; it appeared to have resulted from trauma during the previous confinement, but no details were available. The patient was told to return in a month for further examination.

When she next presented herself it was decided to perform Caesarean section shortly before the estimated date of full term, for the following reasons: (1) Any attempt to incise or dilate the stricture would inevitably lead to extensive laceration during the delivery of the head. (2) The patient had no living children, and had not been pregnant for thirteen years; and, further, there was a great probability of the pelvis being contracted, although no internal measurements could be taken owing to the stricture. The child was alive and healthy.

The external measurements of Chinese women appear to be smaller than those of European women; the average of five patients who delivered themselves is as follows: Interspinous $9\frac{1}{2}$ in., intercrystal $10\frac{1}{2}$ in., external conjugate $7\frac{1}{2}$ in.

I wish to thank Dr. Minett for having asked me to see, and treat, the case, and for her kindness in superintending all the arrangements for the operation.

R. E. TOTTENHAM, M.D., F.R.C.P.I.,
Professor of Midwifery, University of Hong-Kong.

"NEURASTHENIC" NEURALGIA.

THE following case is worthy of being placed on record as it demonstrates the importance of a routine x-ray examination in all cases of intractable neuralgia.

Mr. D., a well developed man, aged 49, consulted me with regard to a "new wisdom tooth." He also complained of neuralgia, which had attacked him frequently during the last seven years. He said that about six years previously his doctor commenced medicinal treatment for the neuralgia. As there was no improvement at the end of the month the doctor advised the removal of all his teeth, although they appeared fairly good. This drastic measure gave no relief, and finally the patient was described as neurasthenic, and apparently abandoned.

Examination showed a healthy edentulous mouth except for a small mass of enamel near the middle line, which I diagnosed as an unerupted supernumerary central incisor. Under a local anaesthetic the gum was reflected and a portion of the external alveolar plate removed, exposing, deep in the alveolus, a very large perfectly developed canine, which I removed. The appearance of the periodontal membrane suggested hyperaemia, otherwise the tooth was healthy.

In reply to questions, the patient stated that, as far as he was aware, there had been no abnormality in his dentition. Three months later he returned for new dentures, and then volunteered the information that since the removal of his "wisdom tooth" his neuralgia had completely disappeared.

It would appear that an x-ray examination would have saved the doctor considerable trouble and the patient much suffering.

Lower Bebington, Cheshire.

J. P. FLYNN, L.D.S.