ULTRA-VIOLET LIGHT AND THE ANTISCORBUTIC VITAMIN.

BY

PHILIP EGGLETON, M.Sc.LOND.,

AND

LESLIE J. HARRIS, PH.D.CANTAB., M.Sc.VICT.

THE object of this communication is to present the experimental results obtained in a preliminary investigation, carried out in the autumn of 1924, to determine the relation, if any, of vitamin C to ultra-violet light.* There are several grounds for supposing that there might be some such relationship. In particular, the fact that vitamin C is so largely associated with plant life suggests its possible intimate connexion with phyto-chemical synthesis. Vitamin C had not previously been studied from this aspect when this work was planned, although of course the relation of the antirachitic vitamin (the "D" fat-soluble factor) to light rays has claimed the attention of a number of investigators during the past year or two. Our conclusions, it may be confessed, have been mainly negative—namely, that the C vitamin does not show that interrelation with light which is apparent in the case of vitamin D; we are not at the moment continuing the work, but it is thought perhaps worth while to summarize our conclusions, particularly as other workers are now interested in this field.

HISTORICAL.

The known facts relating to the fat-soluble vitamins are briefly as follows: (1) Coward¹ has shown that "light is necessary for the formation of vitamin A in plant tissues."† (2) It is now well known also that the ill effects caused by insufficiency of fat-soluble vitamins can to some extent be remedied by exposing the animal on the deficient diet to a source of ultra-violet light.^{2,4} (3) Finally, it has been shown by Steenbock and his collaborators,⁵ and independently by Hess,⁶ that it is possible to activate a ration deficient in fat-soluble vitamins by exposing it to light, thereby conferring upon it both growth-promoting and calcifying properties; cholesterol has recently been activated in this way.⁷

Light appears to have no effect upon vitamin B phenomena.⁸

FIG. 1.—Effect of light on synthesis of vitamin C in germinating oats. Average growth curves (all concordant) of a number of guinea-pigs fed on—A, oats, plus cabbage (for vitamin C); B, oats only, ungerminated; C, oats only, germinated in complete darkness (seven days); D, oats only, germinated in light from mercury vapour lamp; E, oats only, germinated in daylight. All animals received in addition daily rations of cod-liver oil (1 drop), bran, rock-salt, and water. The amount of food eaten was weighed daily (same applies to all figures).

EXPERIMENTAL.

Our investigations were concerned mainly with the effect of mercury vapour or other light on:

1. Synthesis of vitamin C in the plant.
2. Animals kept on a scorbutic diet (attempt at prevention).
3. Animals already with scurvy (attempt at cure).
4. Rations with partly destroyed vitamin C (attempt to regenerate).
5. Rations rich in vitamin C (attempt to augment).

*The view (proved erroneous in this paper) that sunlight (or ultra-violet light) is necessary for the synthesis of vitamin C has been taken for granted in some quarters without any scientific evidence.

†According to Luce and Maclean,² however, yeast cells can synthesize vitamin A in the absence of sunlight.

Throughout we used a Cooper-Hewitt quartz mercury vapour lamp. The power of the lamp to emit ultra-violet rays was tested by means of Webster, Hill, and Eidinow's acetone methylene blue solution.¹⁰ No special precautions were taken to prevent oxidation of exposed foodstuffs. It is unlikely that our main conclusions would be affected if oxidation were to be rigorously excluded.

FIG. 2.—Relative efficiency of oats germinating in light or dark in preventing scurvy. The above are average growth curves for sets of three or more animals. All animals consumed the same weight of germinated oats (calculated in weight of original ungerminated cereal) whether germinated in the dark, in light from the mercury vapour lamp, or in daylight. Oats germinated in the dark had sprouted most and were most efficient in preventing scurvy. In the experiment here represented germinated oats were approximately 50 per cent. of the total food consumed. Other details as in Fig. 1.

1. Synthesis of Vitamin C in the Plant.

Ungerminated oats contain no vitamin C. During germination vitamin C is known to be synthesized. We carried out tests to determine whether light was necessary for this synthesis; we found it was not necessary. The diagram (Fig. 1) needs no further explanation.

Relative Amounts of Vitamin C Produced in Dark and in Light Respectively.—Incidentally it was observed during the course of this part of our investigations that oats, germinating in the dark and producing therefore a greater

FIG. 3.—Effect of exposing animals on scorbutic diet to mercury vapour lamp. Broken line=scorbutic controls in dark (diet completely deficient in vitamin C). Continuous line=animals on same diet but exposed to mercury vapour lamp (few minutes to several hours daily; distance from lamp averaged about 4 feet). Normal controls were also taken (not shown in chart).

length of shoot than when exposed to the light for the same length of time and under the same conditions, produce a correspondingly greater amount of vitamin C. In other words, a larger ration is required of oats germinated in light than of oats germinated in the dark, in order to prevent scurvy. This is illustrated in Fig. 2.

2. Effect of Light on Animals on a Scorbutic Diet.

We found that guinea-pigs exposed to daylight, to mercury vapour light, or to very dull light (diffused from filament lamps) all developed scurvy equally and after the same period when fed on the same completely scorbutic diet (Fig. 3). Experiments upon the effect of irradiating

animals fed on a diet only partly deficient in the antiscorbatic vitamin are still in progress.

3. Effect of Light on Recovery from Scurvy.

We found that animals recovered from scurvy at apparently the same rate whether exposed to the mercury vapour lamp or not, when fed on the same curative diet (Fig. 4).

FIG. 4.—Effect of exposing to light of mercury vapour lamp animals recovering from scurvy. The amount of vitamin C supplied during recovery was strictly limited (2½ grams of cabbage per animal in all cases, 1 gram being approximately the minimum to prevent scurvy), so that any curative action of the light might be better detected. Continuous line=animals irradiated during recovery only. Broken line=non-irradiated controls. Preparatory (scurvy producing) diet was of oats, bran, cod-liver oil, rock-salt, and water.

4. Effect of Light on Rations in which Vitamin C had been Partly Destroyed.

In our earlier experiments we thought we could observe a partial regeneration of vitamin C in heated cabbage after exposure to the mercury vapour lamp. The effect, however, was so slight—a possible delay in the appearance of scurvy for a day or two—that we attach no significance to it.

The following chart (Fig. 5) summarizes a typical experiment, in which exposure of previously heated cabbage to mercury vapour light was without apparent effect on its antiscorbatic potency.

FIG. 5.—Animals received, in addition to scorbutic diet (bran, oats, rock-salt, cod-liver oil, water): Animals A, 5 grams raw cabbage; animals B, 5 grams raw cabbage heated to 100°C. for 1 hour; animals C, 5 grams raw cabbage, heated to 100°C. for 1 hour, and subsequently exposed to mercury vapour lamp (at a distance of a few feet for varying periods, up to 4 hours per diem).

5. Effect of Mercury Vapour Light on a Ration Rich in Vitamin C.

We exposed to the light of a quartz mercury vapour lamp rations rich in vitamin C (for example, a proprietary infants' food), and compared their antiscorbatic value with that of the same unexposed foods. We were unsuccessful in finding any augmented antiscorbatic value. (See Fig. 6.)

Absence of Secondary Radiation Effect.

The fact that vitamin C has its main source in the vegetable kingdom raises the question whether there is

any direct connexion between it and radiant energy in the form of light rays. For fat-soluble vitamin such a connexion is known to exist; and some would go further—basing their arguments on the protection against A or D avitaminosis conferred upon animals (a) by direct exposure to ultra-violet rays, or (b) by contact with other

FIG. 6.—Effect of ultra-violet light on antiscorbatic value of food-stuff. Foodstuff rich in vitamin C was mixed with the standard scorbutic diet so as to supply only 60 per cent. of the minimum required to prevent scurvy in guinea-pigs. No delay in onset or modification of the symptoms was found in parallel feeding tests where the foodstuff had been exposed to ultra-violet light immediately prior to feeding. Continuous line=guinea-pigs on irradiated diet. Broken line=guinea-pigs on control (non-irradiated) diet. Normal controls were also taken (complete diet).

animals which have been exposed (it is claimed),⁹ or (c) by consumption of dead food or growing plant which has been irradiated, or (d) by dwellings previously irradiated (disputed)¹⁰—and would maintain that in the prevention of avitaminosis some form of radiant energy plays its part no less than a definite chemical substance, the accessory food factor. There is little or no experimental evidence to support this contention. Our previous results make it highly improbable that the C factor is interrelated with radiation phenomena. The following experiment, yielding, as was expected, negative results, is in further agreement with this view.

We allowed ultra-violet light from the quartz mercury vapour lamp to act on various foodstuffs, some rich and some deficient in the C factor, and tested these foodstuffs immediately afterwards towards a photographic plate. No activity towards the plate was found.

It may be noted that Dr. Janet Clark¹¹ has recently arrived at a conclusion similar to that reached by us in our second section. Honeywell and Steenbock¹² have separately investigated a problem similar to that dealt with in our first section and reached a similar conclusion. They used germinating barley, while we used germinating oats.

DISCUSSION ON INFANTILE SCURVY.

The statement that the synthesis of vitamin C in the plant is dependent upon the action of sunlight has appeared in more than one textbook, but no experimental evidence has been produced in support of it. True, it has been urged in this connexion that infantile scurvy is (according to clinical observation) more prevalent after the darkness of winter than in summer. The explanation, however, may well be—quite apart from all questions of greater susceptibility to disease during wintry conditions—that bottle-fed infants¹³ receive in winter milk less rich in vitamin C than in summer, owing to the fact that the cow's winter dietary is relatively deficient in fresh greenstuff or other sources of vitamin C. The antiscorbatic value of milk has been shown to depend on the presence of the C factor in the

* The disease is rarely encountered in breast-fed infants, for the human mother's winter diet is not so seriously restricted as the cow's; although such sources of vitamin C as salads and fruits are scarce in winter, oranges and potatoes are in most cases readily obtainable and cheap. Further, breast milk is received by the infant with its vitamin value unimpaired, while bottle milk may have had its value diminished by pasteurization, boiling, or drying.

diet. Cow's milk is accordingly richest in this vitamin when the animals are at grass.* The lower antiscorbutic value of winter milk may therefore be attributed to the cow being put on an inferior dietary, rather than to any direct action of sunlight on the diet (or on the animal itself).

CONCLUSIONS.

A noteworthy antithesis is seen to exist between the fat-soluble vitamins and the antiscorbutic vitamin. The antiscorbutic vitamin can be synthesized in complete darkness, and experiment has so far failed to connect it in any particular whatever with ultra-violet light; the fat-soluble vitamins, on the other hand, have been shown to be affected by ultra-violet light in their synthesis and activation as well as in the prevention and cure of avitaminosis. The apparent independence of vitamin C to light is in a sense surprising, in view of its special association with the plant, in the metabolism of which light plays so important a part.

The practical significance of our results is to emphasize the importance of the provision of the C factor in dietaries. While it is possible to replace the fat-soluble factors to some extent by light rays (acting either on the diet or the animal), the prevention of scurvy must—in the present state of knowledge—rest upon the provision of adequate supplies of vitamin C in the food. Of special interest is this in relation to infantile scurvy—Barlow's disease. It is worthy of note that of the numerous proprietary foods sold as substitutes for maternal feeding, we know of only a single one that can claim to contain in itself an adequate supply of the antiscorbutic factor, permitting one to dispense with food adjuncts.

SUMMARY.

The association of vitamin C with plant life suggests its possible connexion with phyto-chemical synthesis. Yet,

(1) Light was found not necessary for the synthesis of this vitamin (in germinating grain), contrary to former supposition and in distinction to vitamin A, which is said to require light for its formation in the plant;

(2) The direct exposure of an animal on a scorbutic diet to ultra-violet light did not appreciably retard the onset of scurvy;

(3) Ultra-violet light appeared to have no curative action on a scorbutic animal;

(4) Cabbage in which the C vitamin had been largely destroyed by heat was exposed to the mercury vapour lamp and tested for regeneration of the vitamin, with doubtful results;

(5) Exposure to ultra-violet light of an infants' food already rich in vitamin C gave no evidence of an augmented antiscorbutic value.

These results indicate that while the effects of fat-soluble vitamin deficiency in, for example, an infant's diet can be remedied to some extent by exposing the infant (or the food) to sunlight, if one is to prevent scurvy one must rely on an adequate amount of vitamin C in the diet.

Cow's milk may be deficient in this vitamin in winter owing to the animal's inadequate stall diet. The observed increase of infantile scurvy in late winter probably results from this cause rather than from absence of sunlight, to which it has been attributed latterly without sufficient evidence.

We wish to express our cordial thanks to Professor Sir F. G. Hopkins, F.R.S., and Professor J. C. Drummond for much kind interest and encouragement.

REFERENCES.

- ¹ Coward, K. H.: *Biochem. Journ.*, vol. 17, 1923, p. 135. ² Luce, E. M., and Maclean, I. S.: *Ibid.*, vol. 19, 1925, p. 47. ³ Hume, E. M., and Smith, H. H.: *Ibid.*, vol. 19, 1923, p. 364. ⁴ Goldblatt and Soames: *Ibid.*, vol. 19, 1923, p. 622. ⁵ Steenbock, H., and Black, A.: *Journ. Biol. Chem.*, vol. 61, 1924, p. 405. ⁶ Hess, A. F.: *Science*, September 19th, 1924. ⁷ Drummond, J. C., Rosenheim, O., and Coward, K. H.: *Abstract in Chemistry and Industry*, March 13th, 1925, p. 123. ⁸ Krizenecky, Jaroslav: *Arch. ges. physiol. (Pflüger's)*, vol. 204, 1924, p. 467. ⁹ Steenbock, H.: *Science*, September 5th, 1924, p. 745. ¹⁰ Webster, A., Hill, L., and Eidinow, A.: *Lancet*, April 12th 1924. ¹¹ Clark, Janet: *Science*, January 9th, 1925, p. 45. ¹² Honeywell, E. M., and Steenbock, H.: *Amer. Journ. of Physiol.*, vol. 70, 1924, p. 322.

* Literature quoted in the Medical Research Council Report on Accessory Food Factors, 1924, p. 72.

CYSTIC DILATATION OF THE COMMON BILE DUCT.

BY

NORMAN H. HILL, M.D.LOND., M.R.C.P.,
ASSISTANT PHYSICIAN, BELGRAVE HOSPITAL FOR CHILDREN;

AND

R. A. RAMSAY, M.CHIR.CANTAB., F.R.C.S.,
ASSISTANT SURGEON, BELGRAVE HOSPITAL FOR CHILDREN.

THE following case is of interest owing to the rarity of the condition, the difficulty of making a correct diagnosis, and the pathology of this and somewhat similar conditions.

H. F., female, aged 2 years and 3 months, was admitted to the Belgrave Hospital for Children on January 2nd, 1925; the child had been vomiting for a fortnight, and a rapidly progressing enlargement of the abdomen had been noticed for the past month. The vomiting started with occasional attacks, and she vomited about four times during the first week. This became worse, and during the week prior to admission she vomited every day, and sometimes three times a day. The vomit was independent of food, and was chiefly nocturnal. She did not complain of any abdominal pain, and was not constipated.

She was said to have had an attack of jaundice when 9 months of age, which lasted for some weeks, but had had no other illness. She was breast-fed for eight months and then given glaxo and cow's milk. She was the youngest of a family of five. Three sisters and one brother were alive and well; one sister, the fourth child, aged 4 years, had jaundice at 2½ years, which lasted four months, but is now healthy and the abdomen normal to palpation, the jaundice having apparently been of a catarrhal nature.

On admission the temperature was 97°, the pulse 116, and respirations 24. The child was fairly well nourished; there was no jaundice, but the complexion was of a rather "earthy" hue. The tongue was moist and covered with a thick white fur. The cranial nerves, heart, and lungs were normal. The abdomen was large and protuberant, and the percussion note was tympanitic all over, including the flanks. The liver could be felt two fingerbreadths below the costal margin; its edge was sharp and very hard; it was not tender. The spleen also could be felt about two fingerbreadths below the costal margin. On the right side of the abdomen a large mass extended from beneath the liver to just above the crest of the ilium; it did not appear to move on respiration; on bimanual examination the tumour was found to extend backwards into the right loin. It was thought to be a tensely cystic swelling in connexion with the right kidney, but owing to the distension of the abdomen it was difficult to palpate; it was not tender. The urine was faintly alkaline, and contained no albumin, sugar, or bile salts or pigments.

During the first week the child vomited nine times. She was not constipated, and the faeces were of a normal colour. She lost weight. A blood count showed red blood cells 5,096,000, white blood cells 7,200, haemoglobin 90 per cent., colour index 0.88. The various cells were in their normal proportions. The Wassermann reaction was negative. The vomiting, which occurred mostly at night, improved with alkalis, and during the second week three small vomits only were recorded. They consisted of partially digested food and a small quantity of bile-stained fluid. During this week weight was regained and became the same as on admission.

During this period in hospital the temperature, pulse, and respiration remained normal, but the abdomen was still very distended and the swelling in the loin was thought to have increased in size. In spite of the facts that the vomiting had stopped, that she took her food fairly well, although never very hungry, and that the bowels were open, she began again to lose weight, and it was decided to perform an exploratory laparotomy in order to ascertain the nature of the tumour.

Operation.

On January 19th one of us (R. A. R.) opened the abdomen under general anaesthesia, by an incision through the right rectus muscle, over the tumour. The tumour was found to be globular and cystic, and about four inches in diameter. It lay below and separate from the liver; the right kidney was clearly felt behind the cyst, and the duodenum lay upon and was closely adherent to its anterior surface. The liver felt hard and its surface rough, and the gall bladder, slightly distended, projected between the liver and the cyst. The cyst was explored with a needle and syringe, and clear mucoid pale yellow fluid was withdrawn. The cyst was sutured to the margins of the upper part of the abdominal wound, and a tube inserted into its cavity through a small incision, to the edge of which it was fixed by a stitch; leakage was prevented by a purse-string suture inverting the edges of the incision in the cyst around the tube. The abdominal wound was closed except for the point at the upper part where the tube emerged.

Examination of the fluid aspirated from the cyst showed that it contained no albumin, no urea, and no pancreatic ferments. It contained bile salts and bile pigments.

After-History.

The day following the operation the child seemed fairly well; there was no vomiting; the bowels were not open, but a small pale motion was passed the day following (January 21st), and on January 22nd two stools of normal colour and consistency. After this the stools were extremely pale, and contained no bile salts or bile pigment. The wound drained profusely, some days more than others, and in amounts varying from 10 to 20 ounces of thin greenish-yellow bile.