conditions of life, I advise the patient to wash his neck and chest in the morning with cold water, not to wrap himself up too much in warm neckties, shawls, etc., but to wear flannel underclothing; to take his usual food, with the exception of strong spices, mixed pickles, etc.; to drink claret in moderate quantity, but to avoid heavy beers, champagne, brandy, whiskey, etc., and never to drink spirits in excess, because this of course augments the congestion. I do not order the patient, if he be accustomed to strong smoking, to give this up all at once, because I find that the sudden leaving off smoking generally brings about dyspepsia, loss of appetite, constipation, and other disagreeable symptoms; but I diminish the quantity to two or three cigars daily, and strongly forbid his staying in a room in which many other people are smoking together, because the inhalation of the cold tobacco-smoke acts decidedly more injuriously than the smoking of a cigar itself.

A very good aid towards a lasting cure is the visit to a suitable watering-place, if used at the proper time. I have stated that the use of the natural mineral waters alone will never cure the affection, but I find that, added to the local treatment, they have a decidedly stimulating effect upon the mucous membranes of the pharynx and larynx, and do excellent services in preserving the patient from a recurrence of the affection, to which this class of diseases is unfortunately so very liable. I can recommend especially the warm soda and chloride of sodium springs of Ems, and the salt-water springs of Reichenhall, Ischl, etc., for the purpose. If the patient leave it to me, to choose the proper time for the treatment, especially in very old cases, in which it does not matter much whether the disease is left a little longer to itself, I like to begin the treatment at the beginning of the spring, and to send the patient immediately after its completion to Ems. There I direct him not only to drink the waters, but also to gargle with them and to inhale them; in other words, to continue the local treatment in a somewhat milder form. If his means and time admit, he may then stay for a few weeks in the Black Forest, in Switzerland, at a moderate height (I find the Engadine not suitable for these chronic laryngeal affections), before returning to this country. Such a line of treatment has, in my hands, hitherto, without any exception, resulted in a perfect success. It is quite true that all this does not give an absolute immunity against the recurrence of the affection if occasioned by some accidental cause, but a repetition of the treatment—which is in such cases invariably much shorter—and a renewed visit to Ems are sure to remove the evil again.

In conclusion, I can say that I have had the pleasure of curing several cases of from ten to sixteen years' standing by one course of local treatment, followed by one or two visits to Ems in the same and in the following year; and as the patients come to see me once or twice a year, I have the opportunity of convincing myself that the effects of the cure are lasting. I can thus conscientiously recommend the line of treatment which has been indicated.

NURSING INSTITUTIONS.

By F. T. PAUL, F.R.C.S., Liverpool.

SINCE the question of nursing has come thoroughly under discussion, it will, I think, be the better settled if the systems which are found to answer best in the more important of our hospitals, provincial as well as metropolitan, are made generally known. The immediate point at issue appears to be the system that shall be adopted at Guy's Hospital. Now, as an old student and house-surgeon at Guy's, I am intimately acquainted with, and thoroughly appreciate, the many good features in its nursing. Still the most partial friend must admit there was room for improvement. The nursing depended almost entirely upon the excellent and well-trained sisters, supported, it is true, by a number of very good nurses, but at the same time by an infinitely greater number of bad ones. The nursing staff in my time was most carelessly recruited. The matron seemed willing to accept a nurse without any regard for her character. As an example—an exceptional one, I admit—I may mention that a sister once complained to me that, of her six nurses, five came from the patients in Patience Ward (venereal), and the sixth had three illegitimate children. The training, too, was quite without method. It is true, I believe, that they had a few lectures from Dr. Steele; but it was an absolute matter of chance what variety of cases they saw, and they were quite dependent upon the goodwill of their sister or fellow-nurse for what teaching they might happen to get. Their position, too, was a very menial one; they had to do all the dirty work of a ward-maid, and to scrub the floors. The old matron had been almost brought up at Guy's, and saw no necessity for a change. A new one comes, no doubt one who has been well trained; she sees these imperfections, and, like the proverbial new broom, immediately

upset the whole concern. Had Miss Burt, fortunately for herself and the hospital, consulted with the staff, a result might have been arrived at which would have been equally beneficial to both. Yet, since the matter has been brought before the public, the practical experience of other institutions may help towards a satisfactory settlement.

In Liverpool, connected with the Royal Infirmary, we have not only a medical school, but also a nursing school; and the latter is not so well known as it deserves to be. The Nurses' Home, as it is called, is a large building in the grounds of the Royal Infirmary, containing excellent accommodation for a great number of nurses and the lady-superintendent and her assistants. It is presided over by a special committee—I believe a subcommittee from the Managing Committee of the Royal Infirmary; and its objects are: 1. To teach and qualify nurses in every branch of nursing; 2. To nurse the Infirmary; 3. To nurse the outside public. The relations of the Infirmary to the nursing institution are, that probationers shall be trained in the wards; that the staff, house-surgeons, etc., shall be requested to assist in teaching them; and that, when necessary, the funds of the Infirmary shall be voted for the support of the institution. In return, the Infirmary has the first choice of nurses, and the services of the lady-superintendent and her assistants. The relations with the outside public are, that they pay from one to two guineas a week for a nurse, by which means the institution becomes almost self-supporting, while the public have the advantage of being able to procure at a moment's notice an intelligent, well-educated, and well-trained nurse, suitable for any emergency.

The probationers are required to enter for three years. The first year is spent in hospital training; the last two years usually, for the most part, in private nursing. The main object in requiring three years' attendance is that, having received one year's free training in the Infirmary, during the next two they may become useful and profitable to it. After this, they are entitled to their qualifying certificate, and can either leave to seek their own fortunes, or be placed on the staff of the Infirmary, or become private nurses at a higher rate of wages. The lady-probationers can, by paying fifty guineas to the institution, obtain a certificate at the end of their year's hospital training, and are not required to serve during the two following years like the free probationers.

The medical world have long ago admitted that, from a medical point of view, the services of a well-trained nurse are second only to their own; and, apart from the selfish requirements of any special institution, the best system must be that which diffuses its good services the most widely. As things at present stand, I think there is no better system than that which we have adopted, and which works so well in Liverpool. It offers the fullest opportunities for benefiting the public.

There was a time when at many hospitals, Liverpool among the number, the lay committee would scarcely tolerate students in the wards. Ultimately, they began to see that this was a short-sighted policy; and that they themselves would very likely one day trust their health to the keeping of those whose medical education they were opposing; and it is only now that people are beginning to see that, if they are to be well nursed in their illness, they must make some provision for properly training private nurses. This can only be well done in our large hospitals. The truth of that being admitted, we may justly ask: What greater right has a public institution to refuse to become a nursing school than it has to become a medical school? I do not apply this directly to Guy's; for the question may not have been put to them. But I think that it ought to be put to every important hospital in the country, and, for my own part, I can scarcely see any just ground for refusal.

My intimate knowledge of the working of the Liverpool nursing system is drawn from having held the position of Resident Medical Officer at the Infirmary, and, as far as my experience goes, I consider the Liverpool public, both rich and poor, to be nursed as well as or better than any other section of the community.

What I have said has, I am aware, very little reference to the present deadlock at Guy's Hospital. But such innovations as neglecting the wishes, or ignoring the authority, of the staff; changing the permanent sisters or head-nurses from ward to ward; ordering patients up at five o'clock in the morning, and such things, admit of no discussion, but have only to be known to be condemned by any one who knows anything of hospital management. One can only conceive that Miss Burt, having been very successful in carrying out her ideas at Leicester, and having been greatly patted on the back by such friends as Dr. Marriott, has become a red-hot enthusiast in her own cause. But, that her enthusiasm has so far misled the clear judgment of Dr. Steele, as some of the letters in the JOURNAL infer, I am loath to believe, until he places his view of the question before the public.

Every one must admit that, in the present state of things, an inquiry into the nursing system at Guy's Hospital is called for. If such an inquiry take place and originate a great change, I believe that, if my old Alma

Mater open her doors for the education of nurses, she will confer a great boon upon the public, and upon a number of women who want and like such work; and that the renown that she is sure to attain as a nursing school will only add an increasing lustre to her fame, while she augments the number of those who lovingly cherish her memory.

ABNORMALLY HIGH TEMPERATURES.

By JOHN W. TEALE, F.R.C.S., Scarborough.

THE entire omission of any reference to my case in Dr. Donkin's recent paper (BRITISH MEDICAL JOURNAL, December 20th, 1879) on abnormally high temperatures (an omission which, Dr. Donkin informs me, was due to the amount of scepticism he found amongst scientific men in London as to its reliability), induces me to ask that I may be allowed to draw attention to some of the leading features of the case, and to draw some conclusions thereon.

In February 1875, I brought before the Clinical Society the particulars of a remarkable case which had occurred in my practice. A young lady having received serious spinal injuries from a fall from horseback, had for sixty days a temperature ranging from 108° Fahr. to 122° Fahr. and upwards, and eventually recovered. The great length of time that the high temperatures lasted, enabled me to take my observations with the utmost deliberation, and the greatest care to guard against error or fraud. A great variety of instruments were used by me, seven in all, exclusive of the one used by my brother, Mr. T. Pridgin Teale of Leeds, who saw the case three times with me in consultation. Most of these afterwards received the Kew certificate of accuracy, and all were supplied by Harvey and Reynolds of Leeds. Observations were taken at all sorts of times, both of day and night, by myself. More than once in the night I took the temperature when the patient was asleep, having carefully placed the thermometer in the axilla without rousing her. Observations were repeatedly taken in both axillæ simultaneously, once in the rectum, once in the mouth, and once in both axillæ and between the thighs simultaneously. On these special occasions, the temperatures were also taken by myself. Trained nurses, three different ones during the illness, were constantly in attendance upon her, specially cautioned to watch her. The patient was never made aware of the extraordinary character of her case, and for the greater part of the time was too ill to take any interest in the matter. Hot-water bottles were never in contact with the body, except sometimes to the feet by my special order. I was in the habit of carefully removing the instruments into the adjoining room, and the height was read off in the presence of two or more witnesses.

In July of the same year, 1875, I wrote in the JOURNAL a letter replying to various questions and sceptical criticisms on this extraordinary, and at that time unique, case; and I was able, owing to my patient having had a relapse under the care of another medical man, to give that gentleman's experience of her high temperature. This gentleman, Mr. Hind of Stockton, told me afterwards that on one occasion the heat in the axilla was so great, that the mercury burst the tube at the top of the instrument, the index being found in the broken-off air-space at the top.

This, so far as I knew, was the first case published bearing on these abnormally high temperatures. It was the first which seriously challenged the then accepted view of physiologists, that such temperatures were incompatible with human life. As long as this case stood alone, it was natural that scientific men, whilst cordially acknowledging the care that had been taken in watching it, should suspend their judgment. I was content to wait, feeling confident that, if my case had been carefully watched and faithfully reported, as I knew it had been, a time would arrive when other cases would be discovered to confirm it. As time went on, first one case and then another was published in the medical journals, all proving the occasional occurrence of temperatures, usually considered to be fatal, in persons who not only recovered, but who, at the time of the excessive temperatures, did not appear to be in imminent danger of dying. This series culminated in that of Dr. Roddick of Montreal, quoted by Dr. Donkin in his paper in the BRITISH MEDICAL JOURNAL for December 20th last, with a recorded temperature of 117° Fahr., in which the mercury was lost in the bulb at the top of the thermometer. The temperature reached 117° Fahr., but how much higher it might have gone cannot be known, as the instrument could not record it.

I claim Dr. Donkin's interesting series of cases as a strong corroboration of the truthfulness of mine. The following points seem now to be clearly established: 1. Temperatures above the degree formerly supposed to be necessarily fatal do sometimes occur without a fatal issue; nay, even without extreme peril to life. 2. Such exceptional

and excessive temperatures as a rule end in recovery. 3. The conditions of body in these cases of excessive temperature appear to be distinct from the conditions existing in fevers, in which the rule as to the extreme peril of temperatures of 107° and upwards remains unassailed.

What the explanation of these rare and exceptional cases may be it is for physiologists to work out, but it can never be settled if they shut their eyes to facts which contradict their theories, and relegate such cases to oblivion.

ANEURISM OF POSTERIOR TIBIAL ARTERY, LAID OPEN BY MISTAKE: ATTEMPT TO TIE THE ARTERY: AMPUTATION.

By G. H. MACSWINNEY, M.D., Brook Green, W.

THE interesting and instructive clinical lecture by Mr. T. Holmes on a case of aneurism opened by mistake, which appeared in the JOURNAL of January 10th, recalls to my mind a case which occurred in my practice in Ceylon, three years ago, which was in many respects very similar.

The patient, a Tamil coolie, was admitted to the Dickoya Hospital early in December. He had got a blow with a stick a few months before on the back of his leg, and soon afterwards noticed a swelling just above the heel. This increased and became very painful, and on admission the tumour was of considerable size, the leg being twice its natural width. The dorsal surface of the foot was oedematous, but there was neither bruit nor pulsation in the tumour. There was, however, a distinct sensation of fluctuation, and the limb was very hot. I ordered a poultice, believing the swelling to be an abscess. Two days afterwards, I found the skin over the tumour very thin, elastic, and glistening. The appearance was exactly the same as that of an abscess about to burst. The man complained so much of the pain that I determined to open the tumour. I made an incision about an inch and a half long, when, to my surprise, there was a considerable gush of bright arterial blood. I immediately plugged the wound with lint, steeped in perchloride of iron, which arrested the hæmorrhage for the time. Some time afterwards, there was a good deal of oozing, so I resolved to tie the artery. Chloroform was administered, and a tourniquet and Esmarch's bandage applied, and I then made an incision four inches long over the tumour. At first, there was a good deal of hæmorrhage, but this ceased almost entirely when the sac was cleared of its clots. I now proceeded to look for the ends of the artery, but during an unsuccessful search of ten minutes, I came on the tibia, which was so softened by the pressure of the aneurism, that I could push my finger through it. I found that the bone was diseased for several inches of its length, and therefore saw that it would be useless to attempt to save the limb. Amputating instruments were fortunately at hand, so I at once amputated the leg below the knee. The man had a fairly good pulse after the operation, which lasted about an hour, and I hoped he would make a good recovery. During the night, however, he became very excited, and pulled off the bandages from his stump, and attempted to jump out of bed. The result was that bleeding came on, which was with difficulty checked, and he continued so violent that the bleeding came on again severely, and the man gradually sank and died, two days after the operation.

This case is interesting because aneurism of the posterior tibial is very rare, and I believe there are not many such cases recorded. A surgeon is much more likely to open a fluctuating tumour in such a situation, in which aneurisms are rare, than in one in which they are known to be frequent. It is, therefore, necessary that attention should be called to the possibility of the occurrence of an aneurism on arteries such as the posterior tibial.

Mr. Holmes has entered so fully into this subject, that I shall only, in conclusion, call attention to two very interesting cases, described by Mr. Solly, in which aneurisms were opened by mistake.

WEST KENT MEDICO-CHIRURGICAL SOCIETY.—The third meeting of the session was held at the Royal Kent Dispensary, Greenwich, on December 5th, 1879: Mr. J. P. Purvis, President, in the chair. Mr. F. B. Jessett of Erith read a paper on Fractures: their Pathology and Treatment. After going fully into the pathology of fractures, he discussed at length the various modes of treatment, especially urging the early application of fixed splints in all simple fractures of the long bones, plaster-of-Paris and starched bandages being especially recommended; and in the treatment of compound fractures, he recommended the same line of treatment, with trap-doors cut over the wounds for dressing.