ferent causes. Observation of fits suggests, I think, that the venous congestion does produce the clonic spasm, not, however, by direct excitation, but by interrupting and ultimately arresting the spasm which before existed. I think that the fixation of the chest in an epileptic fit simply does in an intense manner that which we do in a slighter manner when we arrest a hysterical convulsion by closing the mouth and nose. There is nothing inconsistent in a certain degree of anoxæmia arresting spasm, and a more profound degree, as in actual asphyxia, exciting it. Analogous difference is seen in the action of many stimuli. The view that the tonic spasm is only clonic spasm compressed, that the clonic spasm is only the tonic spasm spread out, is supported by the fact that in attacks which begin slowly, the slight initial spasm is almost always clonic, and as the discharge spreads and becomes more intense, the spasm becomes tonic. In partial epileptiform seizures, the spasm is usually clonic throughout.

The conclusion, then, is that all the phenomena of the fits of idiopathic epilepsy may be explained by the discharge of grey matter; that the hypothesis of vascular spasm is as unneeded as it is unproved; that there are no facts to warrant us in seeking the seat of the disease elsewhere than in the grey matter in which the discharge commences; that this is in most cases within the cerebral hemispheres, although possibly, in some instances, lower down in the medulla oblongata; that epilepsy is thus a disease of grey matter, and has not any uniform seat. It is a disease of tissue, not of structure.

[To be continued.]

INTRODUCTION

TO A

DISCUSSION ON INTRA-UTERINE MEDICATION.*

By W. S. PLAYFAIR, M.D., F.R.C.P.,

Professor of Obstetric Medicine at King's College; Physician for the Diseases of Women and Children at King's College Hospital; President of the Obstetrical Society of London; etc.

TEN years ago, at the meeting of the Association at Leeds in 1869, I read a paper in this Section "On the Treatment of Chronic Uterine Catarrh". Its object was to show that, in a large class of cases of uterine disease, characterised by those general symptoms which are familiar to the gynaecologist as accompanying morbid conditions of the uterine organs, especially when there was much catarrhal discharge from the os uteri and evidence of inflammation of the interior of the cervix, it was a matter of supreme importance that our treatment should not be limited, as was too often the case, to the vaginal portion of the cervix only, but that topical applications of a suitable character should be made to the deeper seated parts of the uterus beyond the external os, in which the disease was often primarily situated, and of which the inflamed, abraded, and granular cervix was, in a large number of cases, merely the external manifestation. I then described to the Section the method of intra-uterine medication which I practised, and which I more fully detailed in a series of lectures on "Intra-uterine Medication in the treatment of Chronic Uterine Catarrh", published in the *Lancet* in 1873. I venture to mention these facts, since they show that my attention has long been directed to this important and still comparatively unstudied branch of gynaecological practice, and afford, perhaps, some explanation of the honour that has been done me by entrusting to me the opening of a discussion on "Intra-uterine Medication".

There are few subjects connected with the diseases peculiar to women of more interest and importance; and this for two reasons. First, there is no kind of treatment which, properly conducted, is of more value. This is a statement of the truth of which but little experience is required to satisfy ourselves of, since all who have successfully carried it out must know of many patients who have by this means been effectually cured of sufferings which may have lasted unrelieved for years in spite of long-continued general treatment, and which rendered life a burden and a misery. Secondly (and this is even of more consequence), like all methods of powerful efficiency, it is rather a double-edged weapon. Capable of infinite good, it may also, when injudiciously, improperly,

and unskillfully used, do infinite harm. Amongst the various cycles of fashion in uterine disease, there was a time, which many here doubtless remember, when practitioners talked of little else than ulceration of the neck of the womb, and when the speculum and the stick of caustic had a wide-spread use, happily now only a tradition. If, as I fear must be admitted, large numbers of practitioners, in the laudable desire of at least finding out something wrong, and using some remedy, applied nitrate of silver to the exterior of the cervix in hundreds of cases in which it was not needed, although they did not do any good, at least they probably did no great harm. Strong advocate though I be of intra-uterine medication, I earnestly hope it may never become a fashionable treatment, or that any will adopt it unless they have made an earnest study of its indications and mode of application. I shudder to think of the consequences to that long-suffering organ the uterus, if it ever become a routine practice to medicate its cavity indiscriminately. I would, therefore, seriously impress on my hearers the importance of not resorting to the practice we are about to discuss, unless they have fully satisfied themselves that the case is suitable for it, and until they have acquired a certain amount of manipulative skill in the treatment of uterine disease; and I can, at the same time, assure them, for such is my strong conviction from no small experience in intra-uterine medication, that, when these essentials have been seen to, there is no class of case which will give more satisfaction both to them and to their patients.

The limited time allotted to me preclude any attempt at discussing in detail either the symptoms or the pathology of the morbid states in which intra-uterine medication is indicated. To do so in any satisfactory way would involve a long dissertation which, were I even permitted to make the attempt, I dare not inflict on my hearers. To make the matter a little more clear, however, to those who have not made a special study of the subject, I must beg leave to mention, in the simplest and most elementary way, the conditions of the uterus which call for or admit of intra-uterine medication. If I appear to make dogmatic assertions on disputed points of either pathology or practice, I will beg you to believe that it is only because time will not allow me either to enter into arguments in their favour, or to bring forward proofs of their correctness.

In order to understand the reasons why, as I believe, topical applications in various morbid states of the uterus require, to be efficient, to be applied to the deeper seated parts beyond the external os uteri, it is necessary to bear in mind the anatomical structure of the lining membrane of the uterus and cervix. You will remember that the mucous membrane of the uterus is thickly studded with minute orifices, not more than one-thirtieth of a line in width, the openings of the utricular glands, which are small *cysts-de-sac* of structureless membrane lined with epithelium, having their blind extremities in the submucous tissue. These complex structures are always more or less in a state of active change, being thrown off in the form of *débris* at each menstrual epoch, and re-formed by the proliferation of the cells of the muscular and connective tissues. The mucous membrane lining the cervix is even more complex in its arrangements. It is covered with a series of prominent folds or ridges, which collectively are known by the name of the *arbor vite*; and these, more especially on the lower half of the cervix and on its external surface, are studded with innumerable white papillæ, each containing a vascular loop. All over the interior of the cervix, both on the ridges of its mucous membrane and between their folds, are a very large number of mucous follicles, which secrete a thick, tenacious, and alkaline mucus, which is generally found filling the cervical cavity. Here, then, it will surely be admitted, are structural elements of sufficient complexity to lead us to expect that they are likely to be the seat of pathological alterations of no slight importance. Unfortunately we are met with this difficulty, that we are compelled to base our study of them more on the observation of symptoms and the results of treatment, than on carefully recorded pathological facts derived from accurate *post mortem* examination of the structures themselves in a state of disease.

The diseases we are considering, although they give rise to much suffering and ill-health, are never fatal, so that opportunities for *post mortem* investigation do not arise; and the study of gynaecology is too much in its infancy to have enabled those who pursue it to avail themselves of the rare opportunities that may occur. Still, symptomatology and treatment are not without a real value; and, failing more scientific methods, we are surely justified in learning what we can from them. Modern writers on gynaecology have given very elaborate divisions of the various forms of inflammatory affection of the endometrium, which, I am afraid, the state of our pathological knowledge hardly justifies; such, for example, as cervical as distinguished from corporeal endometritis, chronic hyperplastic endometritis, hæmorrhagic endometritis, fundal endometritis, cystic metritis, and the like. Satisfactory as it would be to be sure that all these conditions actually exist as distinct

* Read in the Section of Obstetric Medicine at the Annual Meeting of the British Medical Association in Cork, August 1879.

and recognisable forms of disease, distinguishable one from the other; I believe that the less ambitious plan of simply describing the cases as we meet them in practice, without too strictly binding ourselves to a theory of their pathology, is, in the present state of our knowledge, the wisest course to pursue. By no one has this been better done than by the distinguished gynaecologist who at present holds the office of Master of the Rotunda Hospital, whose very practical work I would strongly recommend to the study of those who are interested in the question. Although I venture to differ from Dr. Lombe Atthill in some of the methods of intra-uterine medication which he prefers, I am sure that I cannot improve on his description of the conditions in which it is indicated, which is obviously the result of much careful study and accurate clinical observation. These differ considerably, as he points out, I am convinced with perfect justice, as they are met with after pregnancy or in abortion, or in the sterile and unmarried.

The former are the most common, and are certainly the most amenable to treatment. In this, as in most forms of uterine disease, symptoms will have developed gradually after a labour or abortion, and may very generally be traced directly to imperfect involution of the uterus, brought on, very probably, by want of due rest or other imprudence. In most of the cases coming under observation, the patient seeks advice after long-continued ill-health, and the failure of tonic remedies and general treatment, so that the disease has become thoroughly established and chronic. Nor are there of necessity such well-marked local symptoms as obviously to direct attention to the uterine organs as the source of mischief. It is rarely that the general health will not have suffered materially; there are debility, loss of appetite, inability to take exercise, and prostration on the slightest fatigue, with the existence of pronounced sympathetic disturbance, such as various forms of neuralgia, flatulent distension, and other dyspeptic symptoms, some of which are often extremely distressing, and may throw the practitioner off his guard by diverting his attention from the true *fons et origo mali*. This is, of course, mere truism to those familiar with the protean disturbances which disease of the uterine organ is apt to produce. Still it will bear insisting on, since the absence of pronounced local symptoms is so often considered to be sufficient to put local disease out of court. Generally, however, careful inquiry will show that local phenomena are not absent, and often they are very distinct. They are of the kind common to very various forms of uterine disease,—bearing down, backache, pain on exertion, and often complete inability to take any kind of exercise; profuse leucorrhœal discharge; disorders of menstruation, more often menorrhagia than any other, frequently to such an extent as to produce much debility, and not rarely associated with dysmenorrhœa; irritability of bladder and frequent micturition; constipation; these and others of the same class, variously modified and combined, will very generally be found.

On vaginal examination, in the more common cases occurring in multiparæ, the local signs are very well marked and distinct. The uterus is generally large, boggy, and tender; and, more often than not, we are able to detect some error of position, either a retroflexion or antelexion, one or other of these displacements being a very frequent complication. The cervix itself is increased in size, the margin of the os is irregular, and its canal often so patulous that the examining finger may be inserted into it, and the hypertrophied ridges of the arbor vitæ distinctly felt. On exposing the cervix by the speculum, it will often be observed to be of a florid red colour, and covered with granular abrasion, resulting from the removal of the epithelium, and the exposure of the hypertrophied villi covering its surface. This sometimes exists to a very marked amount, the part bleeding on being touched; and this, it may be observed, is the so-called ulceration or abrasion which has been the subject of much discussion and controversy. From the os uteri, the characteristic mucous discharge may be observed to exude in great abundance, varying in appearance according to the nature of the case, most frequently being transparent and gelatinous, like white of egg, at other times yellowish and puriform, and, it may be, streaked with blood. On passing the sound, the cervical canal is almost always found to be abnormally patulous, and the passage of the instrument is frequently attended with slight hæmorrhage. The patent condition of the cervical canal is very characteristic, and is of great importance in facilitating intra-uterine medication, inasmuch as it readily allows the passage of the probe, etc.

In the other class of cases already alluded to, the local phenomena are by no means so distinct, and may very easily be overlooked. The chief local sign is the viscid tenacious mucus which pours abundantly from the small contracted os. The cervix itself need not exhibit any sign of disease, being neither hypertrophied nor abraded, although it is often conical and elongated. The body of the uterus is not necessarily tender nor enlarged, although there is very generally a marked flexion, most often an antelexion.

It will be observed that, in the foregoing observations, I have said little as to the pathological changes to be observed in the mucous membrane, or as to the precise parts which are affected; and this I have done intentionally, since I do not believe that the state of our knowledge justifies any positive assertion on these points. Even the granular condition of the endometrium which is frequently described in severe cases of metrorrhagia, although it very probably exists, has not, so far as I know, been demonstrated by reliable microscopic examination. I think it likely enough, as most writers on the subject assert, that, in the majority of cases, the disease commences in the cervix, but it is certain that it soon extends by continuity of tissue, and involves the lining membrane of the body of the uterus as well. Moreover, the question is more of theoretical than of practical interest, since no forms of intra-uterine medication at our disposal can be readily limited at will to one part of the endometrium, but must generally be applied equally to all portions of the mucous membrane beyond the external os uteri.

I proceed, therefore, to consider the more practical part of the subject before us, viz.: Can we by any topical applications to the endometrium hope to remedy the condition of things I have described? How can it best be done? And what are the dangers and drawbacks, if any, to the treatment?

The most obvious plan which suggests itself is the injection of medicated fluids into the uterine cavity, a practice which has been abundantly experimented on from the time of Hippocrates downwards. Those who are interested in the matter will find a very complete historical summary of all that has been done with regard to the subject in Thomas's *Treatise on the Disease of Women*, chiefly condensed from an article on Chronic Metritis by Dr. Cohnheim of Berlin. Without entering into details, it may suffice to say that there is overwhelming evidence to prove that intra-uterine injections are so dangerous, that they are clearly not justifiable as an ordinary method of treatment. Not only are there many cases on record in which they have been followed by severe inflammation and even death, but, short of such serious results, their use has been commonly attended by very grave symptoms, apparently of uterine colic, intense cramp-like pain, prostration, rapid and feeble pulse, and faintness. These symptoms are probably due to retention of the injected fluids in the uterine cavity, either in consequence of occlusion of the canal of the cervix by the pipe of the syringe, or from a concomitant flexion. It has been suggested, and the practice is still adopted by some authorities, that, previously to the use of intra-uterine injections, the cervix should be dilated by sponge or laminaria tents, so as to give a free exit to the fluids. To my mind, dilatation of the cervix is at all times a serious proceeding, never to be lightly undertaken. Any method of treatment involving so grave a preliminary, especially since to be of service it must be repeated frequently at short intervals, is obviously altogether inapplicable for ordinary use. Without farther comment, I may, therefore, say that such injections should, under no circumstances, be used as a means of intra-uterine medication. Various procedures have been employed for the purpose of applying caustic or alterative medicaments to the interior of the uterus, either in the form of ointments, or of solid substances which melt in the uterine cavity, and spread themselves over its lining membrane. Many instruments have been invented for this purpose, by some of which a piece of solid nitrate of silver or some other substance is pushed into the uterine cavity and left there (a method of treatment strongly recommended by Courty of Montpellier); by others, a medicated plasma or ointment is similarly introduced. Sticks or crayons of various substances, such as sulphate of zinc, alum, and the like, are by some inserted into the cervical canal, and allowed to melt. I have made many experiments with these plans, and have long discarded them, believing them to be far more troublesome to use, more painful to the patient, and less efficacious than the method of applying the chosen medicament to the uterine cavity in a fluid form, and swabbing its mucous membrane freely with it, which I shall now briefly describe.

For this purpose, I use small probes of flexible metal, roughened with emery paper, two inches and a half in length, and attached to a wooden handle. (See Figure.) Before operating, three or four of these are covered

with a thin film of cotton-wool. A small piece of cotton is flattened between the palms of the hands; the probe is dipped in water to make the cotton adhere, and, by a little practice, the cotton is wrapped round it so

firmly that it cannot be stripped off by any ordinary force, and so smoothly and evenly as scarcely to add to the bulk of the probe. The patient is placed on her left side, and the cervix exposed by a full-sized Ferguson's glass-spectulum, which answers better, I think, than either the duck-billed or glass-spectulum, as it embraces and steadies the cervix. Before commencing, it is advisable to pass the sound at least once, to ascertain the direction of the uterine cavity, and the probes must be bent to accommodate themselves to it. With one or two of the mounted probes, the cervical and uterine cavities are thoroughly swabbed out, so as to clear them as much as possible of the discharges which might otherwise prevent the medicaments used from reaching the mucous membrane efficiently. If the discharge be very tenacious and adherent, it may be advisable to moisten the cotton with water or glycerine to facilitate its removal. Another probe is then dipped into the solution to be used, and the cavity is thoroughly swabbed with it. In the majority of cases of endometritis, there is no difficulty whatever in passing the probes, since the cervical canal is abnormally patulous, and they enter with the greatest ease. Indeed, increasing contraction of the cervical canal, and consequent difficulty in passing the probes, is frequently a sign of improvement. Dr. Atthill, in the cases in which he applies fuming nitric acid to the uterine cavity, presently to be considered, uses a small vulcanite speculum, which he passes into the canal of the cervix, and through which the probe is introduced. I believe, however, that the use of this unnecessarily complicates the procedure, and that it has no great practical advantage.

Such is the plan of intra-uterine medication I myself adopt; and I now proceed to say a few words as to the agents used and the time for using them. Before doing so, I would wish to make a preliminary observation of great importance, which is that, even in cases otherwise eminently adapted for intra-uterine medication, it is often a matter of moment to prepare the uterus for it by certain preliminary treatment. This is of peculiar importance in cases in which there is appreciable tenderness of the organ on examination, indicating some degree of active congestion. A few days, or even a week or two, devoted to relieving this will be well spent. No agent is so useful for this purpose as the nightly application of a tampon of cotton-wool well soaked in glycerine, an invaluable addition to our resources in the treatment of uterine disease, and one which has the advantage of being capable of being employed by the patient herself. Frequent hot water douches, acupuncture, or the application of one or two leeches to the cervix, are also often useful preliminary measures.

A great variety of substances have been employed. Amongst these may be mentioned nitrate of silver in solution, tincture of iodine, chromic acid, carbolic acid, solutions of the sulphate of zinc or of copper, the acid nitrate of mercury, and fuming nitric acid. After many experiments, I have arrived at the conclusion that the safest and most reliable agents are tincture of iodine, and a mixture of equal parts of crystalline carbolic acid and glycerine, the latter being that which I generally use. I conceive that what we require in most cases is not a solution which will cauterise and, if sufficiently strong, destroy the tissues to which it is applied, but one which will so modify the circulation as to cause a healthy reparative action in the diseased structures. Hence some of the substances I have enumerated above are too powerful for ordinary use, and, in fact, do more than is required. No better evidence of this is needed than the case recorded by Mr. Wigglesworth in the *Obstetrical Journal* for January 1879, in which the application of fuming nitric acid was followed by cicatricial contraction and complete occlusion of the os, and subsequent very serious results. Now, carbolic acid has the advantage of being a powerful alterative, causing shrinking and mummifying of the tissues when applied in a concentrated form, but never producing an eschar. It is therefore free from all risk of producing cicatricial contraction. When I first commenced the practice of intra-uterine medication, I was in the habit of employing it once, or sometimes twice, a week, without reference to the menstrual period. I soon found that, when used too near the period, it was very apt to bring it on prematurely, and sometimes profusely. I therefore now limit its use to the first ten days after menstruation; and I find, as a rule, that two applications, at the interval of three or four days, are quite as effectual as a larger number, and much less irksome to the patient. It is probable that they have a better effect at this time, since the superficial layer of the endometrium, shed during menstruation, has not yet been reformed, so that the alterative action of the remedy more directly affects the deeper layer of the mucous membrane. In the more common cases, occurring after parturition, in which the cervical canal is patulous, such a course of treatment, systematically carried on after three or four successive periods, will generally prove effectual in relieving the patient from her sufferings. In the rarer cases in a nulliparous patient, with a conical cervix and contracted os, much greater difficulty is experienced, and the result is frequently disappointing. The amelioration is often

only temporary, and, after a time, the symptoms are apt to recur as severely as ever. It has been pointed out, especially by Thomas and Lombe Atthill, that the reason of this is, to quote Dr. Atthill, "that the conical cervix and contracted canal, coupled with ante flexion, prevent the menstrual discharge and viscid secretions of the diseased lining membrane from obtaining easy exit. The first step, then, towards effecting a cure is to ensure the free escape of the contents of the uterus. There is but one way of effecting this—the cervix must be divided." This, of course, renders the case much more complex; and, as I am always unwilling to be over-active in treatment, it was long before I could bring myself to believe that this procedure might be necessary. I have, however, seen such beneficial results follow in cases in which the more ordinary method had been long and carefully tried, that I am constrained to admit that division of the cervix is, in some cases, absolutely indicated. The full consideration of this subject is, however, of too much importance to admit of being treated in a cursory manner, and I merely allude to it in passing.

I have hitherto said nothing of general treatment; not because I underrate it, but because my special object, at present, is to consider the local management of such cases. Suitable constitutional measures, varying according to the exigencies of each individual case, are of the greatest moment, and will always materially facilitate the cure.

So, again, any complications, more especially the flexions which are frequently met with, should receive careful attention. It is totally impossible for me to endorse the theory of my friend Dr. Graily Hewitt, that endometritis is simply a secondary result of uterine flexion (*Diseases of Women*, 3rd edit., p. 41); but I do not differ essentially from him in my estimate of the beneficial results to be obtained from the judicious use of mechanical appliances when the uterus is distorted, which can be used *pari passu* with intra-uterine medication. What I maintain is, that those who rely too exclusively on one or other deprive their patients of an important aid to recovery, and that the truly scientific method is to get the maximum of benefit from both combined.

I have alluded very briefly to the employment of fuming nitric acid. The use of this agent originated in this country; it has been strongly advocated by the Dublin school of gynaecologists; it has been as strongly protested against by others, who have probably given the matter but little thought, and who have no practical knowledge of its effects; and, therefore, I feel that I ought to state my own opinion with regard to it. The suggestion that fuming nitric acid should be freely applied to the interior of the uterus, at first almost horrified by its boldness, and the most disastrous consequences were predicted. The fact that such men as Kidd, Atthill, and many others testified to its safety and utility, of itself was sufficient, to my mind at least, to ensure for the proposal the most respectful consideration, and I have not been backward in giving it a fair trial. My conclusions as to its use I will briefly state to the Association. In endometritis with profuse catarrhal discharge, I believe that nitric acid is in no way superior to milder and safer remedies; and, although I have used it in a sufficient number of cases to justify my forming an opinion on the point, I have not satisfied myself that it possesses any advantages which counterbalance the bad effect which may possibly follow the application of so powerful an escharotic. It seems to me to be open to the objection I have already indicated, that it is apt to do too much; that it tends to destroy the tissues to which it is applied, while, in fact, our aim ought to be to modify vital action, not to produce an eschar. In cases of severe and otherwise intractable menorrhagia, in which nitric acid has been chiefly used, it stands on a different footing. In them, it is assumed, and I believe with considerable probability, that the entire lining membrane of the uterus is in a thickened, rough, and granular condition; and here the destruction of its lining membrane may have a distinctly beneficial effect. The evidence produced on this point by our Dublin colleagues seems to me too strong to resist; and, indeed, my own experience, so far as it goes, is distinctly favourable. I have seen many cases of otherwise intractable menorrhagia completely and rapidly cured by a single application of nitric acid to the uterine cavity. It must not be forgotten, however, that there is a distinct risk of producing occlusion of the cervical canal. It is true that in no case coming under my own observation have I seen the slightest ill-effects following the use of this agent. Still, the one I have cited from Mr. Wigglesworth is, of itself, sufficient to prove that the risk is not entirely imaginary. I am not sure that preliminary dilatation of the cervix is essential. I have often used it without this being done, and with the best results. I have never, however, applied nitric acid unless I had satisfied myself that the cervical canal was abnormally patulous, so that some degree of contraction would be of little consequence; and, unless this were the case, I should prefer to dispense with this plan of treatment. I am by no means sure, moreover, that the application of nitric acid possesses any advantages in cases of the kind over the use of the dull wire curette, as recommended by Thomas, Mundé, and others; a simple means of

removing any granulations of the mucous membrane which is painless, requires no preliminary dilatation of the cervix, and is oftentimes very effectual.

Before concluding, it is necessary to consider whether intra-uterine medication is attended by any risks which might outweigh the good effects to be derived from it. If roughly and injudiciously used, I can fancy that intra-uterine, like any other form of uterine treatment so carried out, might produce serious inflammatory mischief. When judiciously applied in suitable cases, I am satisfied that no risk of the kind is run. I have been daily practising intra-uterine medication for many years, and I have only in one case seen any trouble of the kind. Every gynaecologist is, however, aware that even the simple passage of the uterine sound may, now and again, produce a like effect; and certainly the possibility of this is no argument against the use of that instrument. Practically, the only bad effect that is at all likely to occur is the occasional production of a hemorrhagic discharge. This most commonly happens either when the treatment is first commenced, presumably because the mucous lining of the uterus is then most likely to resent any mechanical interference, or when the application is made too near the expected advent of menstruation. The discharge, however, never lasts beyond a day or two, and never leads to any serious trouble.

Mr. Wigglesworth, in the paper already alluded to, has stated his fear that intra-uterine medication might produce sterility, by destroying the uterine mucous membrane. If there were any foundation for this (and Mr. Wigglesworth brings forward no cases in support of his theory), it would, of course, be a very strong argument against the practice. As a matter of fact, however, I am satisfied that the apprehension is quite groundless. So far from this being the case, I have long been in the habit of considering the cure of chronic endometritis by intra-uterine medication as one of the most certain means of removing acquired sterility. I have seen so many cases of the kind that, as I stated in a short paper on "Intra-uterine Medication and Sterility", published in the *Obstetrical Journal* of February last: "It has been a matter of every-day experience with me to meet with many cases of chronic endometritis, with subinvolution, after a labour, perhaps years before, in which intra-uterine medication has been followed so rapidly by impregnation as to leave no doubt of the result being due to the removal of the cause which led to the sterility. So commonly have I found pregnancy follow the cure of these conditions, that I have over and over again remarked to patients, who have expressed themselves as rather aggrieved at finding themselves in the family way, that I had come to look upon pregnancy as the nearly certain result and proof of a satisfactory cure." It may suffice to dispose of this theory if I state that, since writing that paper, only a few months ago, I have had no fewer than five cases in my own private practice in which pregnancy has rapidly followed the use of intra-uterine medication in patients who had been long sterile, in one case eleven years, in another seven, having elapsed since the last pregnancy.

In conclusion, I feel that an apology is due to the Association for the very superficial, and therefore necessarily imperfect, way in which the short space of time at my disposal has rendered it necessary for me to treat this important subject. I trust, however, that enough has been said to elicit from others their opinion and experience on a topic of great and increasing practical interest.

ON INTRA-UTERINE MEDICATION.*

By EDWARD JOHN TILT, M.D.,

Past President of the Obstetrical Society of London.

As we have been left to interpret the scope of the subject of discussion, I do not suppose we are asked to discuss the well understood treatment of cervical diseases, nor to consider whatever kind of treatment the inside of the womb may require for cancer or other diseases; and I shall limit my remarks to the treatment of the several varieties of internal metritis.

These varieties are characterised anatomically by great congestion of the mucous membrane lining the body of the womb, by more or less distension of its blood-vessels, by softening or ulceration of tissue; lesions giving rise to purulent, mucopurulent, or sanious discharges. The mucous membrane may be also found roughened by the enlargement of its utricular glands; they may be so enlarged as to look like fungoid bodies, and they were called "fungosités utérines" by Recamier, who recognised in them a not uncommon cause of frequent loss of blood. Admitting that, without some morbid ovarian influence, there is no membranous dysmenorrhœa, that disease cannot well be detached

from internal metritis, to which it often leads; for I have found a history of membranes having been habitually passed, in ten out of fifty cases of internal metritis, long before they came under my care.

Intra-uterine medication may be doubtless required for the cure of the above-mentioned diseases, but very seldom indeed; and that is the object of a first remark. There may be severe internal metritis while the cervix is comparatively healthy, but as a rule there is a considerable amount of cervicitis whenever internal metritis is sufficiently severe to make one think of intra-uterine medication; and I have no hesitation in asserting that this is unjustifiable, until the cervix has been restored to health. Here is a kind of case that I have often had to treat. The cervix is two or three times its average size, harder than usual, and covered with a dusky red mucous membrane. The lips are more or less ulcerated, and the epithelial desquamation extends to the whole length of the cervical canal. The body of the womb is too large, painful on pressure, and, about every two or three months, there is a gradual increase of pelvic bearing-down pains, and then violent forcing pains suddenly expel from the body of the womb about half a teacupful of pus, which is sometimes spoken of as the bursting of an internal abscess. I find that, if the inflammation of the cervix be rightly treated, this forcible expulsion of pus from the womb occurs less frequently, and ceases altogether when the cervix is thoroughly cured. I also find that what remains of pain and enlargement of the womb generally subsides if the patient take moderate care of herself at all times, and great care at menstrual periods, during the following year or two. I thus cured, without intra-uterine medication, fifty well-marked cases of internal metritis, that I carefully studied for my work on *Uterine Inflammation*; and, if I be not mistaken, the same principles had already guided Dr. H. Bennet, and he also found that internal metritis generally yielded to the thorough cure of inflammation of the cervix.

Such, then, is the safe and rational mode of treating the vast majority of cases of internal metritis; and intra-uterine medication should be reserved for very exceptional cases of the following forms of disease.

1. Incoercible blood-loss, resisting milder remedies and menacing life;
2. When internal metritis menaces life or reason, rather by the intensity of its reaction on the system, than by the amount of purulent discharge; for insanity may be sometimes fairly attributed to an impairment of brain-tissue, due to long protracted internal metritis;
3. When, independently of ovariitis, internal metritis leads to an aggravated combination of dysmenorrhœa and menorrhagia menacing life or reason;
4. Membranous dysmenorrhœa of exceptional severity;
5. Habitual abortion, independent of syphilis and ovariitis, and seemingly caused by some morbid condition of the lining membrane of the body of the womb.

It is now admitted, as an essential preliminary of all intra-uterine medication, that it is indispensable to establish a free passage through the cervix for all fluids injected into the womb or by it secreted; and I have found the dilatation of the whole cervical canal by laminaria-bougies, a day or two before menstruation, to diminish greatly the pain and blood-loss attendant on membranous dysmenorrhœa and the uterine irritation that follow the expulsion of the membrane.

With regard to treatment, when internal metritis is leading to death by blood-loss, I recommend the injection of the undiluted tincture of iodine; for, by doing so, I have repeatedly checked flooding, and also uterine exfoliation in membranous dysmenorrhœa. Tincture of iodine commends itself by antiseptic as well as by hæmostatic properties, and by the fact of its having been found trustworthy, for the purpose under consideration, by Drs. Savage, Routh, Fordyce Barker, and Emmet. I have only heard of its use having been followed by one fatal issue, in Dr. Gaillard Thomas's hands; I am not aware of its having led to any pelvic disease of a serious nature; and a manageable amount of acute metritis is the worst result I have observed in my own practice. Similar immunity from serious results cannot be recorded of the injection of a solution of nitrate of silver in cases of internal metritis. The plan has often led to death, and still more frequently to pelvic peritonitis, from which patients have recovered with difficulty. I am struck by the fact, that the liability to serious disease after using intra-uterine injections is not proportional to the energy of the fluid injected, for formidable accidents have followed the injection of glycerine and of plain water. For that reason, when internal metritis has necessitated intra-uterine medication, I have sometimes had recourse to the solid nitrate of silver, and I have dropped into the womb about five or six grains of it, by means of Simpson's "porte-caustique"; thus returning to what Recamier did fifty years ago. I have seen this followed by acute peritonitis; and, as there are cases on record where death has ensued, I am ready to welcome a better plan of treatment. I trust our meeting at Cork will enable us to settle this best plan of treating such cases, for we are now able

* Read in the Section of Obstetric Medicine at the Annual Meeting of the British Medical Association in Cork, August 1879.