an hour later, I found the uterus distended with clots. These being removed, ergot was administered. As contraction did not take place, hot water was injected into the uterus at a temperature of 115° , as verified by the thermometer. In a few seconds, the uterus became soft, distended, and completely paralysed. In this condition, the induced current was resorted to with instantaneous and permanent success.

No. XI was a case of twins; both children were delivered by natural powers. Very alarming hæmorrhage occurring on the expulsion of the placenta, I was sent for. When I saw the woman three-quarters of an hour later, the hæmorrhage had not ceased. The uterus reached to the umbilicus, and was full of clots; these were removed by the hand, and one grain and a half of ergotin was injected under the skin of the abdomen. As the uterus had not contracted ten minutes later, and oozing was still going on, hot water at 120° (by thermometer) was now injected. While the water was beginning to be injected, and before four ounces could have passed through the tube, firm and permanent contraction ensued. How much of this was due to the grain and a half of ergotin is difficult to say.

My object in placing the above cases on record is not to stamp as useless, on the experience of eleven trials, a treatment for *post partum* hæmorrhage which has been found of such signal service in the hands of others, but rather to induce those who have an opportunity or desire of trying the treatment to test with a thermometer the temperature of the water injected in every instance. By this means we shall alone be able, after a more extended experience, to decide whether this treatment should or should not be resorted to for *post partum* hæmorrhage.

The recent experimental researches of Dr. Max Runge on the influence of heat and cold when applied to the uterus (of which there is a short epitome published in the *American Obstetrical Journal* for last April) tend to prove that, if success is to follow the hot-water treatment of *post partum* hæmorrhage, the temperature of the water must not be raised to as high a degree as was done in the above cases.

The phenomena which seemed uniformly to follow the injections in the cases under my own care (No. 11 excepted) were relaxation and dilatation of the entire uterus. If contraction did occur, it was but temporary, and was rapidly succeeded by relaxation. I should mention that, as regards the pulse and general condition of the system, no appreciable alteration was observable from the injections. The researches of Max Runge on the application of hot water to the uterus are very interesting. He experimented extensively on rabbits; the first step being to open the abdominal cavity, draw aside the intestines, and expose the uterus to view. Into the pelvic cavity was then poured hot water at a temperature of 122° . In a few seconds, he observed that the uterus became bloodless, performed peristaltic movements, and then contracted; but this contracted or tetanic condition was soon changed for one of total relaxation and paralysis, and several minutes elapsed before irritability returned to the organ. If the temperature of the water used were as high as 149° , the uterus suddenly contracted once, and the most absolute paralysis followed. When, however, the temperature of the water did not exceed 104° , the uterus was observed to contract, without being followed by paralysis.

Before concluding, I wish to draw attention to the success that followed the use of the induced current in the above cases. Indeed, obstetricians owe a debt of gratitude to Dr. Radford, who as early as 1844 brought this treatment of *post partum* hæmorrhage prominently before the profession. It is, besides being the speediest in action, the most reliable and safest of all methods of treatment.

ON THE TREATMENT OF *POST PARTUM* HÆMORRHAGE BY INJECTIONS OF HOT WATER.*

By JOHN WALTERS, M.B., Reigate.

THE treatment of *post partum* hæmorrhage is a subject which must always possess great interest to those engaged in the practice of midwifery, and the introduction of any new method demands our most earnest consideration; that by the injection of hot water, first introduced by Dr. Fordyce Barker and supported by the high authority of Dr. Atthill, is at once so novel and bold, that, were it not for the high position of its introducers, few practitioners, I fancy, would have ventured to adopt it.

My object in bringing the subject before you this evening is to narrate some of the cases in which I have used it; that, by comparing our experience of its use, we may endeavour to form an opinion as to the

class of cases for which it is most suitable, and how far we may trust it to control the severer forms of uterine hæmorrhage occurring after delivery.

I will now briefly relate some of the cases in which I have employed this method of treatment; and afterwards state the conclusions at which I am disposed to arrive from the experience I have had of its use.

CASE I.—Mrs. D., aged 41; twelfth confinement. She was a pale flabby woman, and had several times nearly died from *post partum* hæmorrhage, which had required once or twice the injection of perchloride of iron. I was called to her on October 29th, 1878, and found the child just born. The placenta was expressed, and ergot given. Free hæmorrhage continuing, and the uterus alternately contracting and relaxing, the hand was introduced and the clots removed, cold being applied externally and pressure kept up. These measures failing to induce contraction and control the hæmorrhage, warm water was freely injected into the cavity of the uterus, after again removing the clots, with the effect of inducing firm contraction of that organ and checking the hæmorrhage. She made a good recovery.

CASE II.—I was called on March 24th, 1879, to Mrs. D., aged 24, in her second confinement. The child had been born two hours, the placenta following quickly. I found her cold and pinched, with very feeble pulse; the uterus was flabby, and free hæmorrhage was still going on. The administration of ergot, removal of clots, cold, and pressure externally, failing to induce uterine contraction and check the hæmorrhage, I injected warm water, as in the former case, with complete success. The uterus contracted well; the hæmorrhage ceased; she soon began to rally, and made a good recovery.

CASE III.—Mrs. W., aged 26, was attended by Mr. Hallows in her second confinement, July 18th, 1879. Chloroform was given during the second stage. Afterwards, she had severe hæmorrhage, which could not be checked by the ordinary means. When I saw her, one and a half hours after the child was born, it was still going on. The uterus was relaxed; she was cold, faint, restless, and blind. Pulse 120, very feeble. A free injection of warm water caused the uterus to contract and checked the hæmorrhage. Stimulants were given by the mouth and rectum; and after two hours she had fairly rallied, and made a good recovery.

CASE IV.—Mrs. H., a primipara aged 24, six and a half months pregnant, was attacked with severe and repeated convulsions at the commencement of labour on September 1st, 1879. Chloroform was given, and delivery effected by version after dilatation with Barnes's bags. The placenta was adherent and attached low down; free hæmorrhage occurring, it was removed by the hand, cold and pressure being continuously applied. The hæmorrhage still going on, warm water was freely injected into the uterus, but it required several injections, and in addition the application of cold externally, to completely stop it and ensure contraction of the uterus. A little albumen was subsequently found in the urine. She made eventually a good recovery.

Dr. Holman tells me that he has used warm-water injections in two cases of severe *post partum* hæmorrhage, with the effect in both cases of checking the severe loss; it was finally arrested completely by the application of ice externally and into the vagina. One was a case of placenta prævia.

Mr. Berridge tells me that he has found hot-water injections successful in causing contraction of the uterus and arrest of severe *post partum* hæmorrhage in two cases, one of which was complicated with concealed hæmorrhage, and the other with placenta prævia.

The following points seem to me of importance in carrying out this treatment. 1. The uterus must be as far as possible emptied of its contents. 2. The water must not be too hot; 100° to 110° are the limits of temperature which ought to be used; 103° or 104° being perhaps the most suitable. 3. The tube must be passed quite up to the fundus, taking care that the os uteri is kept open to allow the water to escape readily. 4. A considerable quantity of water must, as a rule, be employed. By following out these rules, we shall, I think, seldom fail to induce uterine contraction and control the hæmorrhage; when not completely successful, the use of cold water or ice externally or introduced into the vagina alternately with the warm water would seem to be beneficial.

The action of warm water seems less satisfactory when the placenta is attached low down. It seems to stop the hæmorrhage by causing contraction of the uterus. Since using warm-water injections, I have never had occasion to employ perchloride of iron; though, in at least three out of the four cases I have narrated, I should have had no hesitation in doing so formerly before this method was introduced. Cases will, no doubt, occasionally occur in which we must still resort to the use of iron.

The advantages of the warm-water treatment are, that it is cleanly,

* Read before the East Surrey District of the South-Eastern Branch.

always at hand, antiseptic (for every clot, piece of membrane, or placenta is completely washed away), and; so far as I am aware, perfectly safe. I have never seen or heard of any ill effect whatever following its use. It seems to rouse the patient from the cold collapsed condition often induced by severe uterine hæmorrhage.

I will conclude by stating the following propositions. 1. Most forms of *post partum* hæmorrhage can be controlled by warm-water injections properly carried out. 2. The use of cold water or ice alternately with the warm water is in some cases beneficial. 3. There are some cases of severe and alarming hæmorrhage in which, so far as our present experience teaches, we must still resort to the use of perchloride of iron.

ON THE MODE OF CONVEYING STYPTICS TO THE BLEEDING ORIFICES IN *POST PARTUM* HÆMORRHAGE.

By W. BERNARD, F.K.Q.C.P.I., Londonderry.

THE subjoined woodcut represents the gauntlet which was shown at the Obstetrical Societies of London and Dublin, and by myself at the meeting of the Association held in Manchester in 1877, and which I have since modified and improved.

Fig. 1.

The idea propounded is that a styptic free from undiluted blood can, without intervention, be directly sprinkled through the medium of the hand during the removal of membrane or coagula; that by no other method can an application be brought into such immediate contact; and that the two acts—the clearing out of the uterus, and the scattering of the spray—are, *pari passu*, performed synchronously.

The subject has long occupied my attention; but I had not an opportunity of bringing my idea into practical use till some weeks ago; for, as a rule, *post partum* hæmorrhage is of rare occurrence; and, moreover, when we meet with it, we are able to control it by the ordinary means discussed from time to time in the pages of this JOURNAL. Nevertheless, one will come across an occasional case more violent than usual, and apparently uncontrollable.

About a year ago, I was sent for to assist a practitioner in a tedious primipara case. After delivery with forceps, hæmorrhage preceded and followed the expulsion of the placenta. We restrained it with injections of hot water into the uterus and other means, but were forced to fall back on the perchloride. Such was the tendency to bleed, that oozing and coagula continued for three days. On being consulted some months afterwards, I found the lady very pale, and again pregnant. She was then put on the prophylactic treatment, which, up to the time of her confinement, was sedulously and uninterruptedly continued. The pallor and general condition were benefited; yet I had reason to be anxious, not only from the experience acquired in her own case, but from that of a sister also, who (under my care) had hæmorrhage after her three confinements, lasting three or four hours each time. From the use of cold applications, I fear pelvic cellulitis was superinduced. Inflammatory action ended in abscess. She was never again pregnant.

But, to return again to my prophylactic case; labour commenced with strong and regular pains at 6 A.M., and was over at about 12 o'clock.

Though the membranes were ruptured early, the uterus was persistently followed down, and ergot was administered, yet, on the expulsion of the shoulders, blood began to pour. The uterus contracted well, and the placenta was quickly thrown off. In this short space, the system sustained so severe a shock that I was beginning to fear the worst. Stimulants were given, and ergot and ether were injected hypodermically. On this occasion, I was ready to encounter the enemy; for I had appliances in readiness, and consequently, in a few moments, my hand and glove were in the uterus. On the cock being turned, I felt the fluid trickling over the gauntlet, and after a short interval, by a strong contraction, my hand was pushed from the uterus. Not a single bad symptom followed. There was no return of the hæmorrhage, and no bloodless face remains to be seen, as in her previous confinement.

Fig. 2.

Judging from the past history of herself and sister, I feel bound to assert that the hæmorrhage was sooner and more efficiently arrested by my mode of procedure than by the syphon. To me, the introduction of the hand soon after labour is not a dangerous act. I never witnessed any bad results from it, either in my own practice or in that of others. On the other hand, I regret not having often insured a more firm contraction; for it, and it alone, will excite this when nothing else will do so. Had I done so, I should not have gone away from some of my cases with a feeling that anæmia or autogenetic fever might supervene.