polypus, cancer of the body, etc.; but from digital examination, he can get no information to guide him. In such a case, if hæmorrhage be not great, it is not necessary to do more than keep the patient at rest, freely open the bowels, and give ergot. The degree of hæmorrhage which we ought to consider profuse depends, not upon whether the patient thinks it excessive or not, nor even upon the absolute amount of blood that has issued from the vagina (supposing that this point could be ascertained), but upon the effect of its loss upon the patient. If she be evidently becoming anæmic, her mucous membranes getting pallid, then the hæmorrhage, even though it may be described to us as only slightly exceeding the patient's customary menstrual loss, yet demands treatment; and conversely, if, although the patient complain of profuse discharges of blood, yet her complexion and mucous membranes remain florid, and her bodily functions be well performed, no interference involving even a slight risk should be resorted to. It is usually desirable to make a vaginal examination, unless there be strong reason against it (as in the case of a young unmarried woman), because uterine hæmorrhage, not enough *per se* to produce grave disturbance of health, may yet be an early symptom of disease, such as cancer of the neck, or polypus, in which the sooner local treatment is carried out the better for the patient. But, should vaginal examination reveal nothing, we should not go further till we have objective evidence that the hæmorrhage needs to be stopped.

Artificial dilatation of the cervix is sometimes, though seldom, followed by ill consequences; and, therefore, we should not expose our patient to the risk of those evils, unless we are satisfied that the disease is injuring her health to such an extent that it is worth her while to incur the possibility of ill results. We are not at present in possession of any data from which to state in what cases it is that dilatation is not well borne, or in what proportion such cases are met with; these are points that yet have to be made out. But, speaking generally, I believe that, in the large majority of cases, this process is accompanied with a good deal of pain and some pyrexia; sometimes by much retching; occasionally perimetritic inflammation results; and death has been known to follow. The pain and pyrexia usually subside when the dilatation is finished, and the risk of more serious consequences is small; still, it is large enough to be kept in mind, and to induce us to refrain from this measure, unless we ourselves are quite certain that the disease present is impairing health.

The treatment practised and advocated by Récamier and his followers for hæmorrhagic endometritis was to scrape out the interior of the uterus with the curette. Various kinds of curettes have been devised, the latest, and probably safest, of which is the "dull wire" curette of Dr. Thomas of New York. Published cases show that, by this means, conditions presenting similar symptoms to the disease in question, may be cured. But this does not seem to me the best possible practice. The first thing, in my opinion, should be to ascertain as accurately as possible what is the morbid condition we are dealing with. The hæmorrhage may, for instance, be caused by a small fibrous polypus, in which case the curette would be quite useless, and might be injurious. And, according to Aran (*Maladies de l'Utérus*, p. 478), who speaks from experiments on the dead body, it is not possible with the curette to be sure of touching every part of the inner surface of the uterus. It therefore seems to me but a haphazard practice to scrape the uterine wall without having first found out what is the morbid condition present; for such treatment may not be suitable to the case, and if it be, we may fail in carrying it out successfully.

If, in a case of the kind, the general measures to which I have referred, which cannot make the patient any worse, fail to check the hæmorrhage, the next step should be to dilate the cervical canal, either with sponge or laminaria tents, until it is big enough to admit the finger. Then the cavity should be digitally explored, and treatment suitable to the condition present carried out. If chronic hyperplastic endometritis be the cause of the hæmorrhage, it is to be recognised by the enlargement of the uterine cavity, the thickening and ready detachment of its lining membrane, and the absence of any other morbid condition.

The treatment recommended by Routh and by Olshausen is the removal of the diseased lining membrane of the uterus with the curette. This measure has in their hands proved very successful. In my case, the application of strong nitric acid alone effected a cure. It would obviously be going too far to say, from one case, that the application of nitric acid will cure every case of the kind. But, in some of the cases in which the curette was used, a relapse took place; therefore this mode of treatment is not invariably successful. Bischoff, on account of the possibility of a relapse, thinks it desirable, after the use of the curette, to apply some mild caustic to the uterine cavity. It appears to me that, by passing a swab charged with nitric acid through the dilated cervical canal, and pressing it against the interior of the uterus, we are at least as likely to modify the nutrition of the lining membrane as if

we scraped it off with a curette; that we are more certain of reaching every part of the internal surface; and that this treatment involves a less degree of rough interference with the uterine cavity than does the use of the curette. These considerations, joined to the favourable result of the case I have narrated, would induce me, should another case of the kind come under my care, to treat it in the same way.

CHOREA SANCTI VITI, OR ST. VITUS'S DANCE:

ITS TREATMENT WITH THE BATH MINERAL WATERS.

By A. B. BRABAZON, M.D.

Physician to the Bath Mineral Water Hospital.

I WILL preface my remarks on the above mentioned subject with an apology to the numerous readers of the JOURNAL, for inflicting upon them the results of observations, which have nothing to recommend them but a truly honest desire to place before them what I consider sufficient grounds whereon to rest a simple, agreeable, and generally available mode of treatment, for an affection or disease which is always troublesome, and occasionally intractable. While engaged in the consideration of the subject, I was irresistibly led on to seek information regarding the early history of "Chorea Sancti Viti". An historical sketch of the disease, interesting though it might be, would occupy too much space in the JOURNAL, to the exclusion of more valuable material; but I would strongly recommend to those interested a perusal of the historical sketch of this and other analogous diseases, given in Hecker's *Epidemics of the Middle Ages*, translated from the German by Dr. Babington and published by the Sydenham Society in 1844.

I have traced the history of this very extraordinary affection so graphically described in those pages, and extending from a date almost prior to the Christian era, down through the middle ages, to the present time, through all its varied phases of development, as the dancing mania, or Chorea Sancti Viti, through the bacchanalian orgies dedicated to St. John the Baptist, the tarantism of the Italians, the ecstatic movements of the dancing dervishes of Asia Minor, witnessed by myself, the "convulsionnaires" in France, the hysterical results of camp meetings in America, and of religious revivals in our own country; and I have come to the conclusion that unhealthy, morbid, mental excitement now, as in past ages, is powerful to produce evil effects; and the fact that such effects are now only met with in isolated cases—though these cases are sufficiently numerous—is due, *not* to want of material on which to work, but to the removal of the superstitions of the dark ages, to the advance of civilisation and education, and to the teaching of sound and practical religion, instead of enthusiastic spasmodic fanaticism.

The earlier writers on the Bath mineral waters are, as far as I have been able to ascertain, silent on the subject of St. Vitus's dance, or chorea, with one exception. I have in vain searched for a record of any such case in the works of Drs. Wilkinson, Barlow, Lucas, Charlton, Phillips, Wm. Falconer, and Gibbs. The single exception is Dr. Peirce, who mentions five cases of Chorea Sancti Viti to show "How many strange infirmities our crazy carcasses are here liable to"; of these patients four were males and one a female—the latter did not use the baths. The ages of the males were thirteen, fifteen, and twenty-four, the female was under fifteen; all the males are reported to have been cured. To come down to our own period, in Dr. R. W. Falconer's treatise on the *Bath Mineral Waters*, he states that, in chorea or "St. Vitus's dance", bathing and douching down the whole back is productive of good results. In Dr. Tunstall's essay on the *Bath Mineral Waters*, of which an edition revised by Dr. Carter has recently appeared, chorea is referred to under its common name of St. Vitus's dance; and it is therein stated, "the Bath waters are so useful in themselves, that there is no need to enlarge upon auxiliary remedies, such as strychnine and sulphate of zinc." In my paper on the "Uses and Abuses of the Bath Mineral Waters", published in the BRITISH MEDICAL JOURNAL, 1878, I gave the statistics, as far as was possible with the means at my command, of the curative effects of these waters in chorea. I recorded sixteen cases of this disease discharged from hospital during the year ending May 1877. My remarks are to the following effect. "The total percentage of 'cured' is 75 per cent. The results of treatment of this troublesome disease by mineral baths were to me quite unexpected; and, had I not the opportunity of personal observation in many cases, I should have been sceptical."

I regret that the individual history of patients admitted into the Bath Mineral Water Hospital, as regards the previous subjective symptoms, is generally very imperfect and obscure, being dependent upon the brief and usually *ex parte* statements of friends and professional advisers

naturally anxious to gain admission to hospital for their clients. The clinical history in hospital is, I fear, not as closely attended to as is the case in general hospitals. However, although the appended list of cases, which have been selected at random from the records of the hospital, may not contain much which will contribute to scientific knowledge, I think much useful and valuable practical information will be gained on the subject of a simple and successful mode of treatment of a disease, which in past ages became developed occasionally into a formidable epidemic, and even in the present age must be regarded as an affection, always troublesome, often intractable, and sometimes incurable.

My remarks on the appended cases, eighteen in number, will be brief. All but one were females; their ages varied from eight to twenty; their residence in hospital varied in duration to from three to four months. The existence of the affection before admission, as far as was ascertainable, varied from three weeks to fifteen months. Exciting cause is given as fright in four cases, acute rheumatism in three, rheumatism and hysteria in one case, hysteria alone in one case, sympathy in one. Fourteen out of eighteen were discharged cured, four were discharged "much better". Eleven cases were treated by baths and internal use of mineral waters alone. In the remainder, the supplementary treatment was iron, arsenic, iron and quinine, zinc, bromide of potassium, and, occasionally chloral hydrate. One case only, twenty-seven years of age, was readmitted.

were most unsensational. There happened to be, in the same ward with this girl, a child suffering from chorea, but not of an exaggerated type. After being in hospital about a month, this girl was suddenly seized with chorea in its most severe form. In attempting to walk, she knocked herself black and blue against furniture or whatever came in her way; she could not hold a cup in her hands, or feed herself, and, at night, was sometimes jerked as it were out of bed. Her facial contortions and grimaces were horrible to behold. No votary of St. John, no disciple of St. Vitus, no victim of tarantism could have proved a more perfect specimen of "Chorea Sancti Viti" of the middle ages. It became necessary to isolate this girl completely from her fellows, some of whom, in accordance with the history of the disease, would in all probability have been similarly affected, and the Mineral Water Hospital would have been assimilated to an asylum for "jumpers" or "shakers". This patient was placed in a room by herself, with one attendant, the room being so padded and arranged that she could not injure herself. Mr. Coppinger, the Resident Physician, has favoured me with notes of the case. I should mention that in this, as in all the cases which I have seen, the thoracic movements were bilateral. The medical treatment employed was bromide of potassium and chloral hydrate in full doses at night. When the symptoms were most violent, six leeches were applied to the inside of the thighs; and, after a time, when the acute symptoms had subsided, she had tincture of belladonna perchloride of iron, and a reclining bath daily at 100° Fahr. This was

TABLE.—Eighteen Cases of Chorea.

No.	Sex.	Age.	Admitted.	Discharged.	Cause.	Treatment.	Results.	Duration of Disease before Admission.
1	Female.	8 years.	September 28, 1875.	December 23, 1875.	Fright.	Baths and waters.	Cured.	2 months.
2	Do.	16 "	March 11, 1876.	June 22, 1876.	None.	Baths and waters.	Cured.	1 month.
3	Do.	15 "	October 22, 1877.	January 10, 1878.	Do.	Liq. arsenicalis, baths and waters.	Cured.	"
4	Do.	16 "	March 10, 1876.	June 17, 1876.	Do.	Baths and waters.	Cured.	"
5	Do.	14 "	December 3, 1874.	February 11, 1875.	Do.	Baths and waters.	Cured.	3 weeks.
6	Do.	8 "	August 23, 1875.	October 21, 1875.	Fright.	Baths and waters.	Cured.	"
7	Do.	15 "	January 18, 1872.	May 9, 1872.	Acute rheumatism.	Baths and waters.	Cured.	3 weeks.
8	Do.	16 "	" "	May 31, 1878.	None.	Baths and waters.	Cured.	15 months.
9	Do.	20 "	March 18, 1878.	May 2, 1878.	Hysteria.	Iron and quinine, baths and waters.	Much better.	"
10	Do.	" "	" "	September 2, 1878.	None.	Baths and waters.	Cured.	3 weeks.
11	Do.	9 "	July 26, 1879.	October 22, 1879.	Fright.	Baths and waters.	Cured.	"
12	Do.	27 "	" "	October 9, 1879.	Hysteria, rheum.	Val. zinci gr. iii, ter die, baths and waters.	Much better.	"
13	Do.	18 "	" "	September 18, 1879.	Acute rheumatism.	Baths and waters.	Much better.	"
14	Do.	15 "	" "	September 10, 1879.	Valv. disease, acute rheumatism.	Baths and waters.	Cured.	"
15	Do.	14 "	May 19, 1879.	September 16, 1879.	"	Baths and waters.	Much better.	14 months.
16	Do.	16 "	July 18, 1879.	September 23, 1879.	Fright.	Liq. arsenicalis, zinc. ferri, baths and waters, etc.	Cured.	14 months.
17	Do.	13 "	August 10th, 1879.	October 29, 1879.	Sympathy.	Baths and waters.	Almost well.	Never had it.
18	Male.	12 "	March 22nd, 1880.	May 10th, 1880.	"	Reclining bath: liquor arsenicalis.	Cured.	"

Of the eighteen cases recorded, six arose from causes directly or indirectly influencing the nervous system, as far as regards volition and co-ordination; to this category I might add four of the six cases in which no causes are given; thus, there would be ten cases to be placed to the credit of functional nervous lesion, real or imaginary; in four cases, the symptoms had been preceded by rheumatism, acute or chronic. The part which rheumatism plays in producing chorea is apparently a *questio vexata*. Taking a broad view of the matter, I do not think that rheumatism, *per se*, has much influence directly in producing this disease, but I think these affections may be and are frequently concomitant; and I believe that both diseases are the outward and visible signs of a poisoned and consequently deteriorated condition of blood, causing, in the case of chorea, defective nerve-nutrition, and its consequent train of symptoms. I think, too, that we may ascribe the frequent connection of amenorrhœa and chorea to the same cause, viz., a vitiated and impoverished condition of blood, producing its effects on those functions which, considering age and sex, are most likely to be disturbed, viz., the functions of the cerebro-spinal system. If we divide choreic cases into two classes: (1) those in which the symptoms are connected with actual cerebro-spinal disease, accidental or congenital; (2) those in which there is no such causation present, my opinion would be that cases of the second class are instances of perverted or inco-ordinate muscular actions, owing to loss of volitional power, due to functional disturbance of the nervous system; further that, generally speaking, owing to impaired or defective blood-making, nerve-nutrition is "below par", and consequently the nerve-functions are easily morbidly affected by exciting causes, such as fright, sympathy, or imitation.

The seventeenth in my category presented in a most remarkable manner the apparently uncontrollable power of morbid sympathy or imitation to produce chorea in its most violent form. This patient, a girl aged 18, was admitted to hospital for chronic rheumatism and amenorrhœa. She was a stout healthy looking girl. Her home was in the quiet village of Wrington in Somersetshire; her mode of life and general surroundings

palpably a case of acute chorea, induced by morbid sympathy, in a female at the age of greatest nervous susceptibility, and under circumstances of functional derangement adapted to aggravate the symptoms. I think there could be no better example than this of a case of acute chorea, and no case better suited to form a link in the chain of connection between the epidemics of the middle ages and the sporadic cases of the present. It also bears out to some extent the theory of Dr. Sturges as regards this and other analogous diseases, regarded by him as simply an affection of function.

I have an extreme aversion to making a statement of successful treatment of disease, unless I can adduce some good reasons for removing that treatment from the accusation of being "empirical". Past history helps me materially in my effort to prove that there are good reasons to expect successful results from such treatment.

There is incontrovertible evidence that this disease is the same now as in ages past, only differing in degree, that difference being due to surrounding circumstances, to which I have alluded. I think also the history of chorea goes far to prove that the immediate causation is a "functional" disturbance of the balance of nerve-power, and that this nerve-storm may be caused by conditions involving impaired nerve-nutrition, such as are contributed by depraved blood-making, as in rheumatism, amenorrhœa, and possibly other allied diseases.

There exist abundant proofs of the beneficial effects produced by the internal and external use of the Bath mineral waters in improving depraved conditions of blood; thus their use would tend to remove the causes leading to the production of choreic symptoms. Again, it is well known that these waters exercise a decidedly sedative influence on the spinal nerves; thus their application would tend directly to subdue or diminish aggravated involuntary muscular movements. I believe myself that time, care, careful moral training, proper nutrition and isolation, may probably effect a cure; but, from personal experience in private cases, I know that cure will not be effected in anything like the

time within which it can be, with the addition of the use of the Bath mineral waters.

I hope that my remarks may prove a just claim that, in this troublesome affection, treatment by these waters should be regarded as based upon a rational and not merely an empirical foundation.

ANNULAR STRICTURE OF THE INTESTINE.

By STEPHEN MACKENZIE, M.D., F.R.C.P.,

Assistant Physician to, and Lecturer on Pathology at, the London Hospital.

HAVING recently had another case of annular stricture of the intestine under my care, and successfully diagnosed, I am glad of the opportunity in reporting it to answer some observations which my former paper on the subject* called forth. In the article in question, I wrote, "The fact that full sized properly formed fæces are occasionally passed of course shows that there can be no organic stricture". It is this statement which has been called in question. Dr. Heath of Torquay, in the BRITISH MEDICAL JOURNAL for September 27th, 1879, page 496, draws attention to the fact that Dr. Walters of Reigate, in the same number of the JOURNAL as that in which my article appeared, communicated a case of annular stricture of the junction of the sigmoid flexure and rectum, for which colotomy was performed. In his description of the symptoms, Dr. Walters says, "Sometimes the evacuations were not larger than the little finger; at others, large bulky motions would be passed", and, commenting on the case, he writes, "In stricture in this region no positive reliance can be placed on the size of the stools: fecal matter may dribble through a stricture situated high up in the rectum as this was, accumulated in the lower part of the bowel, and be discharged in a large mass." Dr. Sawyer of Birmingham, after quoting my statement, says his experience is directly contrary to it, and writes that "The presence of a tight stricture of the intestine, so low down as the junction of the sigmoid flexure and rectum, is compatible with the occasional passage of fecal masses of normal size and shape". He narrates the case of an elderly gentleman whom he saw for intestinal obstruction, and writes, "A solid motion which he had recently passed was shown to me: it was of normal shape and size. I was able to reach an annular stricture at the lower extremity of the sigmoid flexure."

These striking cases show that my statement on this point was a little too unguarded. I was quoting from my own experience, and am very glad to have this supplemented and corrected by that of others. I am much obliged to those who have drawn attention to the point and raised the question as to the value to be attached to the appearances (and statements as to appearances) of the fæces in stricture of the intestine. It will be found that standard authors, as a rule, draw attention to the altered character of the stools in stricture, but do not speak with distinctness as to the absolute value attaching to the sign. Dr. Hilton Fagge, in his valuable article in *Guy's Hospital Reports*, however, writes, "I am, for my own part, indisposed to place much reliance on the statement of a patient, that the motions have for some weeks been abnormally narrow. To expect a trustworthy answer to an inquiry upon this head is to demand more than can ordinarily be looked for, at least from the common run of hospital patients. It has not yet, I think, been determined how near the anus disease must be for such an effect to occur"; he adds, "some careful observations in reference to this question appear to be needed". Dr. Hilton Fagge, in deprecating the importance attaching to alterations in the character of the stools, only intends his remarks to apply to the statements of patients. He considers the symptom of some value when the facts are verified by the surgeon. Whilst admitting Dr. Sawyer's proposition, that "the presence of a tight stricture of the intestine, so low down as the junction of the sigmoid flexure and the rectum, is compatible with the occasional passage of fecal masses of natural size and shape", I still think that the passage of small, lumpy, flattened, or otherwise deformed fæces, when properly formed and sized fæces are never passed, is a sign of great value in the diagnosis of stricture of the intestine. Of course, it is not the only sign, nor invariably, when present, pathognomonic of this condition of the bowel. It would be valuable to elicit in the JOURNAL the experience of different observers on this point, as derived from cases with *post mortem* examinations. The case I have now to report shows, I think, that the symptom is of decided significance.

CASE.—R. C., aged 61, with no hereditary tendency to disease, was naturally loose in the bowels, going twice or thrice a-day. For the

three or four last summers she had attacks of diarrhoea and vomiting lasting about a fortnight; there was no blood in the stools. For eighteen months preceding her death, her bowels became constipated, and she had to take three or four pills a-week to procure evacuations. About September 1879, she began to have pain, like knives cutting her across, about the hips and bottom of the abdomen, which later became localised in the abdomen, and associated with flatulence. Her bowels became more and more constipated and never acted without pills. The motions were hard and lumpy, but not ribbon-like; never properly formed. She was seen by me on December 1st, 1879, when she was suffering from constant pain, flatulence, distended abdomen, constipation, and anæmia. Some sulphate of soda was ordered for her to be taken each morning. The first dose procured a copious pulpy motion, but subsequently it failed in its effects. The obstruction increased, the pains became aggravated, and vomiting occurred. She became a patient in the London Hospital under my care, on December 31st, 1879. Her abdomen was distended, and tympanitic throughout the course of the colon and elsewhere; there was much borborygmus; some tenderness on pressure: no lump could be felt in the abdomen. Rectal examination revealed no disease. Her tongue was coated; appetite lost. Pulse 72, regular, moderately full, soft, and compressible. Respiration 14, temperature 99°. Her bowels had acted the previous day after taking two pills; she then passed with much pain four hard lumps about the size of marbles: the pain continued for some time after the evacuation. A linseed poultice was applied to the abdomen, and a simple enema given; the latter brought away a few scybala. The following day, January 1st, another injection brought away more hard fæces. She vomited. On January 2nd, she was ordered a bismuth mixture and a drachm of Glauber's salts every other morning, in warm water. On January 5th, the bowels not acting, an enema was given which opened the bowels fairly. The fæces were hard, and of the size and shape of walnuts. During the next few days, action of the bowels was only obtained by means of enemata: the motions always hard and lumpy. On January 10th, she had a good deal of pain in the abdomen; a drachm of castor-oil was ordered each morning. On the following day, January 11th, she had a motion, partly liquid, and the remainder scybala about the size of walnuts. She vomited this day. January 12th, she had a motion in the night, and again this morning. There was no vomiting, the abdomen was less tender. January 13th, she had a great deal of diarrhoea in the night due to the castor-oil. There was no vomiting. On January 14th, 15th, 16th, there was more or less free action of the bowels: no vomiting. She felt weaker but thought she could eat a chop on the 16th. January 19th, there was much wind and pain: the bowels were open. She could only take liquid food. January 20th, she complained a great deal of pain, diffused all over the abdomen. She had frequent vomiting. The abdomen was distended but not tense: the stools copious and liquid. She looked hectic, but the temperature was only 98.2°. January 23rd, she had liquid stools and frequent vomiting. Temperature 98.6°; pulse 114; respiration 26. She was vomiting frequently. She gradually became weaker, the pulse increasing in frequency, but the temperature remaining normal till the day of her death, when it rose to 101.5°. The daily excretion of urine ranged between twenty and thirty-five ounces.

At the necropsy, the abdomen was distended; on opening it, the colon was greatly dilated throughout its visible course, and, on displacing the remaining intestines, the dilatation of the colon was found to cease abruptly at the sigmoid flexure. There was slight recent peritonitis. On removing the intestines, the stricture was found to be situated twelve inches above the anus. The bowel was constricted throughout its circumference over an area not greater than one third of an inch, and its calibre reduced to the size of a No. 8 catheter. The stenosed part showed a slightly raised red border of neoplastic nature. The whole of the colon above the stricture was greatly distended and filled with yellow turbid fluid; its mucous membrane was much congested: there was a small superficial ulcer just above the stricture, and some pin-head-sized ulcers were found in the cæcum. The mucous membrane of the ileum was congested, and presented peculiar changes in its epithelium. There was no enlargement of lymphatic glands, no secondary deposit in any organ. The remaining *post mortem* appearances were unimportant to the case.

I regret very much that I did not recommend colotomy, for the case was an exceedingly favourable one for the operation. It is doubtful if surgeons would have consented to perform it with the free evacuations the patient was having. It is quite clear, however, that the relief was insufficient and too late. Changes had taken place in the bowel from over-distension and straining, and inflammation had extended to the peritoneum. I shall have greater confidence in advising colotomy on another occasion. In such cases, as advocated by Dr. Sydney Coupland and Mr. Morris (BRITISH MEDICAL JOURNAL, January 26th,

* "Annular Stricture of the Intestine: its Diagnosis and Treatment" (BRITISH MEDICAL JOURNAL, May 31st, 1879, p. 805).