attached to peristalsis of the intestines. Diminished intestinal peristalsis is a recognised cause of constipation, for which remedies are daily prescribed. It is precisely in such cases of dyspepsia that constipation might be theoretically expected, from the probable existence of a common condition throughout the intestinal tube. In the case of imperfect secretion, on the other hand, digestion is not merely sluggish, but the gastric juice being unable to effect the necessary changes in the aliment, the ill-prepared chyme is unsuited for contact with the intestines. The consequent irritation not only prevents constipation, but sometimes causes diarrhoea.

In some cases, the symptoms which accompany impaired gastric movements are of so general a nature, that the question arises whether the gastric affection may not be incidental to a state of relaxation and want of power in the whole system. The circulation is slow and weak, indicating a relaxation of the vaso-motor nerves, and feebleness of the heart. But the organ is easily excited, and its action is often intermittent. For the latter, there are two causes, namely, mechanical pressure of the distended stomach on the diaphragm, and irritation reflected from the stomach through the pneumogastric nerves to the heart. Sometimes this irritation affects the lungs, when dyspnoea and cough are produced. The tongue, large and flabby, is deeply indented by the teeth, while the muscular structures of the throat, like those of the stomach, are so relaxed, that the part is often a source of constant trouble. There is a general sluggishness of the whole man. The tendency to sleep after meals is, in some cases, irresistible. The mind participates in the torpor of the body, and yet, like the circulation, is subject to be morbidly excited. It is in this form of dyspepsia that the pains of indecision, depression, and apprehensiveness, are most fully experienced. Such is an outline of the symptoms caused by, or perhaps sometimes only coincident with, impaired gastric peristalsis. Mixed cases, in which the effects of defective secretion are combined with those of impaired movements, are to be met with. But, in general, the affections are distinct.

The distinction between their causes is also well marked. The causes of impaired peristalsis may be summed up as those by which nervous energy in general is impaired. Such causes are, hard study, mental strain, depressing passions, prolonged bodily fatigue—in a word, whatever uses up nerve-force in such a way as to leave an insufficient amount of it for an organ, the action of which, although intermittent, requires a large share.

But, in order to maintain its vigour, the stomach requires absolute rest at regular intervals, and, for this purpose, must be empty. When at rest, the organ hangs motionless and nearly perpendicular in the abdomen. The practice of eating too frequently or at irregular intervals is, therefore, a common cause of dyspepsia from impaired motion.

Tea-drinking is a very common cause of impaired gastric peristalsis. This is mainly due to a specific effect on the nerves, and partly to the practice of taking the infusion as warm as possible, by which the tonicity of the muscular coat of the stomach is lowered.

The proximate cause of defective secretion is congestion and consequent gastritis. A primary affection of the vaso-motor centres is probably a frequent cause of congestion; but local irritation, such as from strong alcoholic drinks, hard indigestible food, and food taken in excess of the gastric juice secreted is the ordinary cause.

In the treatment of all forms of dyspepsia, attention to diet claims a prominent place. Articles known to be slow of digestion must be avoided, and a lessened amount of food must be taken only at proper times. But, as a rule, absolute strictness in diet is more necessary in dyspepsia from defective secretion than in that from impaired motion; for, as already said, in the latter affection, digestion is sluggish rather than imperfect. One dietetic rule is, however, of the greatest importance in the present case. The principal meal should be taken early in the day, before the nervous system has been exhausted either by mental or by bodily exertion. In some instances, the power of digestion seems to diminish in proportion as the day advances. A distinguished literary lady consulted me who had, by incessant brain-work, fallen into a state of great suffering from gastric oppression and flatulence after meals. At my suggestion, she dined early instead of late in the day. This change was beneficial, but was not effectual in affording relief. I then advised that she should eat meat at breakfast only, and that no writing should be done before the meal. This plan succeeded perfectly.

From its well-known power in causing muscular contraction, strychnia suggests itself as the remedy for impaired gastric peristalsis. It affords the most powerful means we possess of restoring the gastric functions. I may, perhaps, take some credit for having helped to make known its value. So long ago as 1860, I wrote: "Speaking from extensive experience, I know no single medicine of more value It acts by increasing the tone of the muscular coats of the stomach and intestines. When these coats are relaxed, gases are

generated, mainly owing to retardation of the aliment in the cavities, No remedy has in my hands proved so permanently effective as strychnia against this inconvenience." (*Imperfect Digestion*, 1st ed., p. 186.) In 1864, the late Dr. Brinton, following Chomel, condemned the use of strychnia in stomach-diseases as unnecessary and dangerous. (*Diseases of the Stomach*, p. 334.) But, notwithstanding the condemnation of these authorities, strychnia has held its place in these affections, because, although too often given without discrimination, it proves beneficial in many instances. The secret of its successful administration lies in the recognition of the cases. It is suited for cases characterised by the symptoms of impaired motion; namely, uneasiness, but not actual pain, after food, and flatulence. It is not suited for cases of impaired secretion, characterised by pain after food and little or no flatulency.

Some precautions are of course necessary, and more so because the patients are seldom under daily observation. A dose of one-twentieth of a grain should rarely be exceeded. It should never be given in pills, on account of the difficulty of exact subdivision in that form. The susceptibility of the alkaloid to precipitation by alkalis and some other substances must be kept in view. If so precipitated, the whole of the drug would, of course, be contained in the last dose in the bottle. For the rest, the pharmacist must be responsible. But, after having prescribed strychnia some thousand times, I never knew any harm to arise from its use.

It might be supposed that electricity would prove useful for lesions of peristalsis; but, after many trials of faradisation and a few of the direct current, I am compelled to say that I do not regard it as a useful agent in this affection.

It is sometimes desirable to check flatulence by some agent which hinders fermentation. Formerly, I prescribed carbolic acid for this purpose; but its unpleasant taste is a great drawback. Of late, I have used thymol with, I think, better results; and the taste is far less objectionable.

Many cases are met with in which the stomach is unable to expel flatus in consequence of temporary paralysis from over-distension. Various drugs given to promote contraction of the organ—carminatives, as they are called—sometimes fail in their purpose. It is in such cases that charcoal proves useful. Charcoal possesses a remarkable power of absorbing gases; but this power, as I have elsewhere shown, is very much lessened by long keeping and by wetting. This led me to the plan of giving, in hermetically sealed gelatin capsules, charcoal prepared from vegetable ivory, which kind was proved by experiment to possess the best absorbing power. If, in cases of obstinate gastric distension, three or four such charcoal-capsules be swallowed, a few cubic inches of carbonic acid gas will be speedily absorbed. Tension being now removed, the muscular coat of the stomach generally resumes its power, and flatus is freely expelled.* In a few obstinate cases, however, chiefly when the stomach-affection is secondary to diseases of the liver or kidneys, the muscular paralysis is so complete that, as happens in case of the overdistended *rumen* in cud-chewing animals, mechanical interference is the most effective mode of treatment. For this purpose, I have had made a small India-rubber tube (tube shown) two feet in length, having one extremity closed, and perforated like a drainage-tube to the distance of four inches from the end. Such a tube can be safely and easily introduced into the stomach, and will prove effectual in relieving the distended organ.

I am convinced that certain symptoms, described as a sinking sensation in the gastric region, craving for food soon after meals, etc., are generally to be attributed to hyperperistalsis or exaggerated movements of the stomach. But this constitutes a distinct disease, to be separately discussed. The same remark applies to dyspepsia of liquids and the flaccid state of the stomach which exists in that affection.

CASE OF TONSILLITIS, MEASLES, MEMBRANOUS LARYNGITIS, AND CATARRHAL PNEUMONIA.

By HORATIO DONKIN, M.B.,

Senior Assistant-Physician to the Westminster Hospital; Physician to the East London Hospital for Children, etc.

I THINK the following case worthy of noting at this time, when the natural history and proper classification of membranous inflammations in the throat are under consideration.

A boy, two years and a half old, well nourished and strong, was admitted into the East London Children's Hospital on March 20th, 1879, suffering from a frequent "croupy" cough, and slight inspiratory dyspnoea, but the voice was very little affected. The glands in the left

* Perhaps the best proof of the utility of this charcoal so administered is to be found in the fact, that the capsules are manufactured on a very large scale, and are exported largely.

submaxillary region were enlarged and tender. The fauces were slightly red, but no membrane was seen. The physical signs of the chest were normal. Temperature 102.2 deg.; pulse 136; respirations 36.

This attack had apparently begun with a slight cough, two or three weeks previously to admission. This had been becoming worse till three days before that date, when he was said to be suddenly seized with a "new kind of cough", like a dog's bark. At the same time, the child began to be feverish and drowsy, with some difficulty of breathing, much thirst, and loss of appetite.

On the 24th, the symptoms were not much altered. The dyspnoea was less; the glandular swelling was more marked, especially on the left side. The tonsils were swollen. The voice was not affected, and no membrane was seen. The urine was turbid with urates; it contained no albumen.

On the 29th, all the symptoms were much improved. Temperature 99.

On April 6th, all glandular swelling had disappeared, and the tonsils were much diminished in size.

On April 7th, the temperature began to rise, and there was some coryza, with cough. A well-marked measles rash appeared on the 10th (the disease being in the ward). The cough became again more croupy in character.

On April 12th, at 9.30 A.M., the breathing became markedly affected for the first time; the cough was husky, and the voice was quite lost. The dyspnoea increased till 11 A.M., when there was marked recession of the epigastrium, etc.; and inspiration was very long-drawn and laboured. Very little air entered the chest. The tonsils were seen to be much enlarged, and covered, in common with the back of the pharynx, with thick viscid mucus. No membranous patches were distinguished. At 12.30 P.M., the child began to show signs of lividity. The trachea was opened by Mr. Hayward above the thyroid isthmus. Hardly any blood was lost. Before inserting the tube, feathers were passed through the wound upwards and downwards, with the result of removing small pieces of membrane. Great relief followed the operation. About an hour afterwards, the child sat up in his cot, which was surrounded by a steam tent, and seemed strong and lively. In the evening, he seemed exhausted, and signs of consolidation were observed at the bases of both lungs.

On the 13th, the lung-symptoms had extended. On changing the tube and cleaning out the trachea, some pieces of membrane were extracted. There was very little moist secretion. During the day, diarrhoea occurred, and the child rapidly sank. Death occurred on the morning of the 14th, about thirty-six hours after the operation. A *post mortem* examination was not allowed by the friends.

As far as the clinical symptoms went in this case, which are the chief data on which we must depend for practical purposes, the child apparently suffered from a bronchial catarrh at the outset, in the course of which the larynx became involved, and the tonsillitis occurred. This state of things apparently lasted all March, improving towards the end; and, by April 6th, the boy appeared nearly well. Measles then came on; and, with its usual tendency, attacked the mucous membrane. The child's weak points were its tonsils and larynx, which now became seriously affected, as evidenced by the appearance of the former, and the loss of voice and nature of the cough. Moreover, catarrhal pneumonia, so common in measles, in all probability occurred, which in itself was a sufficient cause of death.

But the operation of tracheotomy revealed the presence, not before made out, of membranous exudation in the trachea. Has this fact, taken in connection with the whole course of events, any etiological significance? Does the occurrence of membrane point to diphtheria as the disease from which the child suffered, and imply a zymotic origin? Or is it to be looked upon merely as an accident in the course of a prolonged catarrhal attack, aggravated by measles?

BEDFORD RURAL.—Estimating the population of this district at 25,203, Dr. Prior makes the death-rate for 1878 to be 17.6 per 1000. The infantile rate and that from zymotic diseases were both satisfactory. No less than 160 out of the total of 444 occurred in persons aged 65 years and upwards. Measles occasioned 8 deaths, scarlatina 4, diphtheria 5, and enteric fever 4 deaths. Dr. Prior's comparison of the mortality from phthisis on the cold impervious clays of North Bedfordshire with that on the warmer soils in other parts of the county militates somewhat against the generally received opinion that consumption is less prevalent on pervious than on impervious soils. Sanitary work appears to be well maintained in the district, but much improvement is still needed in the water-supplies.

CLINICAL MEMORANDA.

ON MEASURING THE PUPIL.

MEN who draw inferences from the size of the pupil like to know that the measurements are taken in the way least liable to error, while those who make the observations desire that the method used should be the simplest compatible with exactitude. Perhaps the most accurate way would be to hold a very fine graduated wire across the centre of the pupil, and so take its diameter; but this is not a practicable method; for, if the scale were easy to read off, it would require to be of such a size as might interfere with the retinal supply of light, and so, by tending to enlarge the pupil, be a source of error. The plan by which the pupil might be approached as closely as possible, without obstructing the light from entering it, is to hold a thin graduated scale (say of ivory) which, I think, Dr. Gowers uses, or the perforated card (Cooper), or the grey paper with black spots on (Nettleship), just in front of the lower margin of the cornea, and see to how many markings on the scale or to what aperture or spot the pupil corresponds. The former of these modes can probably alone be termed a measurement, the other is rather a comparison with spaces of known size; for, if the pupil do not happen to correspond with the size of one of the spots, there is no means of determining its size other than by a kind of guess.

That the measurement by scale is the less liable to error is evident, for the following reasons. The scale can be brought nearer the pupil than the spots, as the graduations are marked quite at the edge. In this case, we measure by mentally letting fall vertical lines from the ends of the horizontal diameter of the pupil, as far as the edge of the scale, and read it off. In the case of the card, since the spots cannot be quite at the edge, the brain has to retain the impression of the size of the pupil for a period longer in proportion to the distance of the spot from the eye, and error is in proportion to the time the impression is retained. In measuring circles, or plane circular surfaces, it is simpler and more practicable to compare the size by comparing their diameters than their circumferences. The pupil may approximately be regarded as a circle. Mr. Cooper's method would compel us to go the roundabout way of measuring the length of a circle, when we might measure the length of a straight line. In catheters, as we cannot directly take the diameter, we take the circumference; but few would undertake to test a catheter-gauge. Mr. Hutchinson's method being an arbitrary one, he himself, as a scientific man, would not expect or wish it to come into general use.

This measurement of the pupil by the length of its diameter is the only one that can be used, assuming that the pupil is not round; and surely no one would wish us to employ two modes of expressing its size—one when the pupil is round, and a different one when not round. Moreover, it is not desirable to increase the number of instruments a man must carry about, unless absolutely necessary; and, as a scale, graduated on one side in *millimètres*, and on the other in fractions of an inch, is used for other measurements, men who do not care to use the metric system will still be enabled to record their observations in every-day language.

C. W. EVANS, Shropshire Eye Hospital.

SURGICAL MEMORANDA.

A CASE OF CYCLODIALYSIS.

THE right eye of Kate B. of Walton, near Warrington, thirteen years old, presents the following features, unaltered from what they were in 1874, when I first saw her; in fact, the mother holds that the condition of the eye has been the same from birth, and ascribes the defect to a fright by a rat in her pregnancy. The eye appears smaller than its fellow, from the palpebral aperture being less widely opened, and further separation of the lids soon proves the eye to be rather intolerant of light. The iris is of a greenish hue, whilst that of the emmetropic and altogether sound left eye is blue. The eye counts fingers with uncertainty. The pupil is somewhat oblong vertically, and its inner third, at some distance behind the plane of the pupillary margin, in front of which the inner edge of the pupil appears unattached, is occupied by a greyish-white mass. The oblong shape of the pupil persists after repeated instillations of atropine, whilst its now kidney-shaped form, with the hilus towards the nose, proves the inner margin to be slightly adherent. On the eye being turned extremely inwards, the white cataractous mass is seen to extend further in that direction than what could be noticed before. The ophthalmoscope, showing the path of the pupil also in its apparently unobstructed two outer thirds to be impervious to light, they are found, under great illumination, to be occupied by a congeries of brownish striæ, about thirty in number, and of a pyramidal shape. These extend from underneath the outer pupillary margin, where