THE LUMLEIAN LECTURES ON THE PATHOLOGICAL RELATIONS OF THE VOICE AND SPEECH.

Delivered at the Royal College of Physicians of London.

By J. S. BRISTOWE, M.D., F.R.C.P.,
Senior Physician to St. Thomas's Hospital.

LECTURE II.—(Concluded.)

Loss or impairment of the laryngeal voice is one of the common incidents of many of the diseases which involve the larynx or its ministerial nerves. It is a common temporary consequence of catarrhal inflammation of the vocal cords and surrounding parts; it is a frequent and often abiding result of syphilitic or tubercular destruction of the vocal apparatus, or of the involvement of the same parts in neoplasms. But the most interesting causes of pathological modifications of the voice are those which are referable to affections of the nerves and nerve-centres concerned in the actions of the larynx. Diphtheria and hysteria are common causes of paralysis of the motor and sensory nerves distributed to these parts, the muscles which govern the tension of the cords being most frequently involved. The movements of the vocal cords are not often compromised in cerebral hemiplegia, yet occasionally in such cases the voice becomes feeble and monotonous. Much more frequently the central causes of laryngeal paralysis are such as involve the medulla oblongata, and especially, therefore, glosso-labio-laryngeal palsy, disseminated sclerosis, progressive muscular atrophy, and the presence of tumours in or about the medulla. Again, any lesion involving one of the pneumogastric trunks, or its superior or inferior laryngeal branch, or any of the subdivisions of these branches, necessarily leads to paralysis of the muscle or group of muscles to which it is distributed. This subject is one of great interest, and might well repay analytical investigation; but I shall confine my remarks to a single point, which to the physician is probably the most interesting of all, namely, paralysis in the domain of the recurrent laryngeal nerve. This nerve, as is well known from its long course and peculiar relations to certain important structures, is very liable to be involved in the progress of material diseases of the root of the neck and upper part of the chest, among which we may include inflammations of the pericardium and pleura, and especially of the connective tissue in the locality referred to, and goitre. But by far the commonest and most important are aneurisms of the aorta or innominate artery, and malignant tumours. Recurrent laryngeal paralysis is generally unilateral and attended with loss of motor power in all the muscles which are supplied by the nerve; namely, all the intrinsic muscles except the crico-thyroid, which is the special tensor of the vocal cord, and is concerned in the production of the higher notes of the voice. The powers of adduction and abduction are lost, and so much of the permanent tension of the cord as is due to the action of the thyro-arytenoid. If such a larynx be examined with the laryngoscope, the paralysed cord will be seen to remain passive about midway between the position it should assume in phonation and that attending ordinary respiration, and motionless excepting so far as it is affected by the breath passing over it. The other cord, however, will be seen to execute its accustomed movements, and probably in phonation will be seen to revolve beyond the median line, so as to come into actual contact with the paralysed cord, the glottis acquiring an oblique position, and the healthy corniculum usually passing in front of the other.

In the bilateral affection, both vocal cords are of course paralysed; both assume the characteristic position; and the aperture of the glottis, which is open to about half the extent it should have in quiet respiration, acquires, as von Ziemssen expresses it, the cadaveric position. The symptoms of recurrent paralysis are well-marked, and, in association with the results of laryngoscopic examination, leave no room for doubt as to the cause. In bilateral paralysis, there is usually complete aphonia; accordingly the voice is reduced to a whisper, and even that is effected with unusual waste of breath, and consequent difficulty. Similarly the acts of coughing and expectoration are impaired in effectiveness. In unilateral paralysis, the voice is also affected; but it is not usually reduced to a mere whisper. It becomes weak, impure or husky, and of heightened pitch—conditions due to the combination of a nearly closed glottis (which, while it allows the escape of air in vocal puffs, allows, also, of more or less escape of unvoiced breath); of vocal

cords in different degrees of tension, and therefore unable to vibrate in unison; and of a vocal cord (the healthy one) which, from the increased effort which the patient makes to speak, is rendered more tense than it would be under ordinary circumstances, and so tends to vibrate with unwonted rapidity. The voice sometimes presents a falsetto character. There has been a general belief (at any rate, I think, among English physicians) that difficulty of breathing, also, is one of the symptoms of affection of the recurrent laryngeal nerves. At any rate, it has been held, and is, I believe, still commonly believed, that not merely loss of voice, but the paroxysmal cough and dyspnoea which are so often observed in aneurism of the arch of the aorta, are the conjoint consequences of implication of the recurrent laryngeal nerves. It is now eight or nine years ago that several cases came under my care, which satisfied me that this view is erroneous, and that the paroxysmal attacks occurring in these cases have a different origin from the loss of voice with which they are frequently associated. Especially I had under my care, within a very short period of one another, two middle-aged men who were suffering from carcinoma, and had, also, unilateral paralysis of the larynx. The first patient had cancer of the oesophagus, a little below the situation of the thyroid body, and extending some distance downwards; he also had several enlarged and hard glands in the lower part of the neck. Besides the usual symptoms of oesophageal stricture, he suffered from hoarseness of voice, which had only come on recently, and was unattended with soreness of throat or cough. On examination, the vocal cords were seen to be structurally healthy; but the right one moved freely, while the left was motionless, nearly in the position of closure. The patient was under treatment for nearly three months, and then died suddenly from perforation of the left common carotid. At the necropsy, the left recurrent nerve was found to be entirely destroyed in a good inch of its course by involvement in the cancerous growth. He had had no difficulty of breathing from first to last. In the next case, the patient had carcinoma of the retroperitoneal glands, stomach, and liver, to which his earlier symptoms were mainly due. Subsequently the lungs became affected, and the immediate cause of death was pneumonia. On admission into the hospital, his voice was a hoarse whisper, and it had been so for six weeks; but he had never had, nor had he then, either cough or dyspnoea. The right vocal cord acted well, but the left was motionless. The paralysis of the left cord, and the weakness and hoarseness of voice, were persistent up to the time of death, which occurred three months after admission. Yet, though during the last three or four weeks symptoms of pulmonary disease, with cough and expectoration, were developed, he never suffered from dyspnoea, still less from paroxysmal attacks. During the last two months, he had a marked difficulty in swallowing fluids, which had a tendency to pass into the windpipe. At the *post mortem* examination, the lungs, with the trachea and oesophagus, were found healthy; but the left recurrent laryngeal nerve was involved in a carcinomatous tumour of the size of a filbert. In both of these cases, it was clearly proved that there was complete destruction of the recurrent laryngeal; and it is certain that, during life, they had presented those symptoms which, *à priori*, one would be inclined to attribute to that lesion—namely, impairment of laryngeal intonation, and (at any rate in one of them) difficulty in opposing the entrance of fluids into the trachea; but there was no dyspnoea, either persistent or paroxysmal.

The next case I shall quote is that of one of my patients, a man thirty-seven years of age, who died in the hospital about six years ago from aortic aneurism. His illness was of about six months' duration, the last three weeks of which time he passed in the hospital. At first, he supposed he had caught cold; and he had from that time up to the day of admission steadily increasing dyspnoea and cough, attended latterly with expectoration. He was a spare man, with dusky face and anxious aspect. The local indications of an aneurism of the aorta, about the point at which the innominate is given off, were quite obvious. His chief sufferings for some little time before he came into the hospital, and during the whole time he was under treatment, arose from constant difficulty of breathing, with occasional sudden terrible exacerbations, during which he had a hard metallic cough, lasting until his face became extremely livid, and relieved by the expectoration of muco-purulent fluid. His breathing was remarkably stridulous. But there was never any tendency for food to go the wrong way; short-breathed though he was, his voice maintained from first to last its natural quality; and there was no indication whatever of paralysis of the vocal cords. His death was due to pulmonary congestion, with lobular pneumonia. At the *post mortem* examination, both recurrent laryngeals were found unaffected; but the trachea opposite the site of the tumour (which was as large as a small orange, and arose from the transverse arch of the aorta close to the origin of the innominate) was much compressed and narrowed; and already ulceration, preparatory to the opening of the aneurism into the trachea, was far advanced.

Another case of special interest has recently been under my care. The patient was a ship's steward twenty-seven years of age, who came into one of my beds in March 1878, and died on the 1st of February last. He had had syphilis. About four months previously, he strained himself in lifting a cask; and shortly afterwards was attacked suddenly in the night with loss of voice. He went for this to the Valparaiso Hospital for six weeks. At first, it was thought that he was suffering from laryngeal catarrh, and he was treated accordingly; but before he left he was told that his malady was really aneurism. On admission into St. Thomas's Hospital, there was distinct evidence of the existence of an aneurism, in the presence of dulness and pulsation over a limited area beneath the inner part of the right clavicle, burning pain in the same part, distension of the right jugular veins, pain down the right arm and up the same side of the neck, and weakness, impurity, and elevation of pitch of voice, associated with paralysis of the right vocal cord. He had no difficulty in swallowing, no obvious dyspnoea, and there was only a slight cough which aggravated the intrathoracic pain. About a week after admission, he had a sharp and severe attack of pericarditis, in which for a time his life was despaired of. From this, he recovered in the course of two or three weeks. About a month afterwards, he had intense abdominal pain followed by diarrhoea. These symptoms lasted for three or four days; and recurred a few weeks later. For several months after the pericardial attack, he thought he was getting better—he felt, at any rate, easier. The pain in the arm, neck, and chest diminished, and his appetite and spirits improved. But his disease was extending. A systolic murmur was detected beneath the inner extremity of the right clavicle, the dulness and pulsation gradually increased in area, and extended beneath the manubrium to the left side. The right radial pulse became smaller and smaller, until it could scarcely be felt, the veins in the neck became larger, and slight oedema of the face appeared. Moreover, though he did not actually complain of shortness of breath, his breathing became manifestly stridulous when hurried or deep drawn. Sometimes also he had to sit up in bed, and had attacks in which he feared he should be choked. Occasionally he complained of a little difficulty in swallowing, and a tendency to choke when drinking quickly. At the end of December, the aneurism had extended considerably to the left, and had caused prominence of the manubrium with infiltration and thickening of the lower part of the neck, and he began to complain of pain down the left arm. And, although he had had cough, with occasional shortness of breath and sense of choking for some months previously, it was now especially that he began to suffer from marked dyspnoea with stridulous breathing, and paroxysmal attacks, coming on at first at intervals of several days, but subsequently several times a day, and lasting from a few minutes to an hour or more. In these attacks, he was compelled to sit up in bed, sometimes sat on its edge, struggled for breath with frantic efforts, grew livid in hands and face, with starting eyes, and a look of intense horror; his pulse became imperceptible; perspiration poured from his face; and occasionally insensibility followed. The attacks were attended or followed by cough, and more or less abundant expectoration. His death occurred quietly some hours after one of them. At the necropsy, an innominate aneurism as large as a cocoa-nut, and filled with clot, was found to have caused erosion of the sternum, to have obliterated by pressure the right innominate vein, and in some degree to have obstructed the left, to have destroyed the right recurrent laryngeal nerve, and to have compressed and eroded both the trachea and the oesophagus, which tubes communicated freely with one another and with the interior of the aneurism, through ulcerated openings which were obstructed (so far, at any rate, as the aneurism was concerned) by the protrusion of laminated clot. The aorta was much diseased; and three other aneurisms were found: one, of the size of half a walnut, in the transverse arch, which caused considerable stretching of the left recurrent nerve; another somewhat smaller in the ascending arch; and another in the lower part of the thoracic aorta, which was obliterated by laminated clot. The right common carotid artery was obliterated. The subclavian was pervious. The heart was attached by old adhesions.

In the former of these cases of aneurism, it will be observed that the recurrent laryngeals were not at all implicated, and that there was no affection of the glottis and no impairment of voice, but that the trachea was compressed by the tumour, and the patient suffered from paroxysmal attacks of cough and dyspnoea of the most aggravated kind. In the latter case, on the other hand, almost the first symptom which the patient complained of was aphonia from affection of the right recurrent; and it was only after some months that he began to show manifest signs of difficulty of breathing, which were evidently connected with the gradually increasing pressure of the tumour on the trachea; nor was it until a later period still, at a time when the direct implication of the trachea had reached serious proportions, that paroxysmal attacks of dyspnoea and cough arose.

The several cases just narrated show, first, that destruction of the functional activity of the recurrent laryngeal nerve is attended with paralysis of the corresponding vocal cord (which can be recognised by the aid of the laryngoscope), with impairment of the musical quality of the voice, and apparently with some difficulty of swallowing, owing to the tendency of food to get into the trachea, but is not necessarily attended with stridor or dyspnoea; and, second, that compression of the trachea involves stridor, and dyspnoea which is often paroxysmal, and liable to end in sudden death, but does not interfere with phonation, excepting in so far as it may render the voice weak by diminishing the supply of wind to the vocal organ. Exacerbations of dyspnoea occurring in narrowing of the trachea may be due in some measure to spasm of the muscular fibres of the affected part, but are referable in some degree to the accumulation of mucus in or below the point of narrowing, and to difficulty of dislodging it owing to the mechanical impediment existing there to the performance of an effective cough, and to temporary swelling of the mucous membrane.

The above discussion raises a question of some importance, namely, as to whether tracheotomy should or should not be performed for the relief of dyspnoea dependent on the pressure of an intrathoracic tumour. Dr. Gairdner many years ago proposed this operation for the temporary relief of the dyspnoeal attacks often complicating thoracic aneurism; and his suggestion has been largely acted upon, I believe. Now, assuming that in such cases the difficulty of breathing depends on pressure on the recurrent laryngeal, and, therefore, occurs at the laryngeal orifice, tracheotomy is not only justifiable, but it becomes our duty to the patient to urge its performance upon him. But, if it be shown that the impediment is not there, but in the trachea, below the lowest possible point of operation, the question assumes a very different aspect. In such a case, my own opinion is certainly, as a rule, against its performance.

I pass on now to make a few observations about those affections of speech, in which the mechanical acts on which articulation depends are interfered with. As a general rule, the greater the skill one acquires in so combining and adjusting the movements of particular groups of muscles as to make them the ready instruments of our will, the more automatic do these movements become, and the less need is there of the direct guidance of the mind. And the perfection of such skill is attained by the performer only when, losing sight of all intermediate processes, he becomes conscious of nothing relating to his performance but the will to execute and the actual execution. In fact, the greater the attention we bestow on our own mechanical operations, the less will probably be the accuracy of the immediate result. Thus, no one in health finds it difficult to maintain his balance, and to walk evenly under ordinary circumstances; but few manifest equal precision when the having to walk over a narrow plank placed over an abyss diverts their attention from the end in view to the actions of their legs and those needed for maintaining equilibrium. Thus, again, the skilled pianist performs the most difficult passages with consummate skill, while she thinks simply of the music to which she is giving life; but when, from nervousness or other cause, her attention becomes divided between her music and her fingers, she is specially apt to become inaccurate and to stumble. So it is with the orator, whose fluency is liable to be checked and whose language to become confused, whenever his mind wanders from the matter of his speech to the grammatical construction of his sentences and the careful choice of his words; and to ordinary speakers, when their attention is distracted from the words they wish to utter to the organs on whose action utterance depends. It is owing to this that soreness and dryness of the mouth are often attended not only by a little thickness of utterance, which is their normal consequence, but by a tendency to leave out or transpose letters or syllables, and to stammer. It is owing partly to this that stammering becomes aggravated whenever the sufferer is under observation, or tries to speak specially well, or is nervous; and, doubtless, the enforced attention which persons whose organs of speech are paralysed pay to the enunciation of their words brings with it a tendency to blundering enunciation and stammering. The condition here referred to is a more or less important factor in the defects of articulation, which are manifested by all those persons who suffer from paralysis or spasm directly implicating the mechanical functions of speech.

It is curious that hemiplegic paralysis of the organs of articulation, and even one-sided paralysis due to diseases of the motor nerves supplied to them, interfere comparatively little with the function of articulation. In either of these cases, the patient can usually pronounce every literal sound without difficulty, manifesting merely a slight degree of thickness in their articulation in connection with one another. One of the most striking forms of general paralysis of the organs of speech is that observed in cases of glosso-labio-laryngeal palsy, and in other morbid conditions of the medulla oblongata. In these affections, the

patient may lose absolutely all power over the muscles concerned in phonation and articulation, and be unable also to masticate and swallow. In glosso-labio-laryngeal palsy, the loss of speech is gradual, and varies in the order of its loss according as the lips, tongue, or fauces are first implicated. If the lips be paralysed, the labial consonants naturally become difficult or impossible of utterance; if the tongue suffer, those which need the use of that organ succumb; if the laryngeal muscles be affected, phonation will be annulled or modified in character. I must say, however, that, both in this case and in allied forms of disease which I have carefully examined in regard to articulation, I have often found that, even when the patient's speech was an inarticulate babble which only those in constant attendance on him could interpret, when his lips were scarcely capable of voluntary motion, and the tongue could not be protruded beyond the teeth, he could still, when asked to do so, repeat with fair distinctness and force all the separate articulate sounds which in combination he found it impossible to evolve. The utterance of the separate sounds, however, has generally been uncertain, so that I have left the patient at one time with the impression that the labials or some other class of letter-sounds were specially defective, and on my return have found them perhaps more distinctly pronounced than any others. In testing patients who cannot pronounce complex groups of letters, and whose organs are not quite under their control, it is important to make them utter the letter-sounds pure and simple, or in easy combinations specially devised, and not to ask them to repeat the names of letters, which, like those of *z*, *h*, and some others, are not letters at all, but words comprising several letters. There is, however, a general tendency for the confusion of the explosive consonants with the corresponding oral or nasal continuants; and for an undue expenditure of breath, which imparts a sibilant or wheezy character to the utterance.

Other affections attended with paralytic impairment of articulation are tabes dorsalis, disseminated sclerosis, general paralysis of the insane, and Sir W. Gull's cretinoid condition of adults. The peculiarities of utterance which attend these are not unlike those displayed in bulbar paralysis; nor is there, I think, any trustworthy difference in respect of speech among themselves. In all, there is a more or less marked feebleness and indistinctness of utterance, which becomes most apparent and probably blundering when the patient is excited or speaks rapidly. In all there is a greater or less tendency in the lips to tremble during speech, or immediately preceding speech, and for fibrillar tremblings or amoeboid movements (if I may so express myself) of the tongue to take place when the organ is protruded. In all of them there is a manifest difficulty in beginning words and syllables, which leads to imperfect utterance and slurring of the letters, but leads also (when the patient tries to speak distinctly) to the divided or scanning enunciation which is so common in some of these cases, or even to explosive enunciation. It should be added that the attempt to speak distinctly induces not only scanning enunciation, but increased efforts of the patient to put his muscles of articulation in motion, which reveals itself either by needless violence and amplitude in the movements of the lips, or by a curious general elevation of the upper lip attending the opening of the mouth, associated with deepening of the naso-labial sulci, elevation of the *alæ nasi*, and even with more extensive co-operation of the facial muscles. Moreover, such patients have a common tendency to speak in a monotone. Of course, in all these cases, implication of the soft palate renders the voice nasal, implication of the larynx may lead to more or less aphonia, and difficulty of swallowing is liable to supervene. I do not mean to imply, by the foregoing observations, that there are no clinical differences in regard to speech between the different paralytic affections which I have grouped together. I believe that such differences do in many cases exist; but I believe that they are rather differences of degree than of kind, and have no specific value.

The most interesting of the spasmodic affections causing defect of speech are chorea and that condition or those conditions to which stammering is due. In most cases of chorea there is distinct affection of speech, which may depend on spasm of the lips, tongue, larynx, or respiratory muscles, or on several of these factors combined. The words are sometimes indistinctly articulated, sometimes drawled out, sometimes uttered with explosive impetuosity, sometimes unfinished, being interrupted by a sudden spasm of the lips or tongue, some sucking or other sound produced in the mouth, a grunt or croak effected at the glottis, or a sudden inspiration or expiration. Rarely, speech is absolutely abolished. I have recently had under my care an intelligent little girl of about ten, who, while suffering from severe general chorea, lost for a period of three weeks all power to utter articulate sounds and to phonate. She understood everything that was going on around her, yet she remained absolutely dumb. During the same time, she had great difficulty in retaining food within the lips, and in deglutition.

Stammering comes on usually in childhood, but occasionally at later

periods of life, probably caused by some sudden impression on the general system or on the nervous organism. It is characterised by a sudden hitch in the act of speaking, a momentary or prolonged arrest in the flow of sounds. It is generally said that it is at the explosive consonants that the sufferer comes to grief, that the continuous consonants are for the most part pronounced readily, and that vowels are uttered freely. These statements, however, do not express the whole truth. For a little examination of a series of stammerers will make it clear that, while some will halt specially at the labial, others at the lingual, and others at the guttural explosive consonants, most are liable at times not only to stumble at every one of the explosives, but even at the continuous consonants, and some will even find an impediment in the utterance of simple vowels. If the check arise at the lips, the evolution of the labial sounds is arrested by a spasmodic closure of the mouth; if at the point of the tongue, this becomes fixed in the position for the utterance of lingual sounds; if at the base of the tongue, this part becomes spasmodically approximated to the palate. In some instances, and particularly when the vowel-sounds are implicated, the spasm takes place at the glottis; and frequently the stammering, whether occurring in connection with vowels or consonants, is due to a sudden inspiration. But whichever part the spasm may affect primarily, the arrest may consist either in a sudden simple more or less prolonged spasm, or in a sequence of such spasms in rapid succession, during which the implicated letter undergoes more or less frequent repetition. Further, there is a tendency, especially if the spasm be prolonged, for a greater or less number of other groups of muscles to become involved. Thus if, for example, the spasm commence at the base of the tongue, we may find that the mouth opens widely and remains in that position, the muscles of expression work convulsively, the glottis contracts, respiration becomes arrested, and at last spasmodic movements of remote parts are also excited. It would be easy to write a minute description of some of these cases, and to show how individual examples differ from one another. But the result would be to prove that their resemblances are far more important than their differences, and that their pathological explanation is the same. Speech, as I have already explained, is, even in its mechanical details, a very complicated art; it requires for its perfect performance, not only that the respiratory apparatus, the laryngeal instrument, and the organs of articulation shall uniformly co-operate with the greatest nicety, under ever-varying conditions, but, as regards the last organs especially, that they shall pass, from arrangement to arrangement, with unflinching ease, certainty, and rapidity. It is in connection with complicated rapidly varying muscular combinations, which have become easy of performance, and more or less automatic by long practice, that spasms or hitches are liable to occur and mar the performance. It is thus that occasionally the swordsman's arm becomes arrested in some particular attitude; the pianist has to give up his playing because of some sudden inco-ordination; and the writer especially is seized during his work with momentary spasm or cramp which checks or spoils his hand. And it is doubtless that that spasm in the utterance of articulate language takes place; and that this spasm (although it may commence in either of the three mechanisms concerned in speech) almost always begins either in the proper organs of articulation or in connection with the mutual co-ordination of the several factors of speech—respiration, phonation, and articulation, and at the commencement of words or syllables. It has been held that stammering does not occur when the sufferer whispers, or when he sings or intones. These statements, however, are inaccurate. Whispering is often attended with stammering; and, although the singing voice is usually free from it, even in aggravated cases, it certainly sometimes suffers.

In concluding these remarks, I may observe that I do not pretend to throw any light on the treatment of this affection. But, inasmuch as it does not depend on any structural defect of the organs of speech, local remedies, with the possible exception of the continuous galvanic current, are not likely to be serviceable. It may, however, be improved, if not cured, by close attention to the management of the breath in speaking, by carefully practising those combinations of letters which are specially liable to give rise to stammering, and by the adoption of a deliberate and carefully balanced method of enunciation.

TESTIMONIAL TO DR. D. J. CUNNINGHAM.—At the close of the session, the students attending the classes of Anatomy in the University of Edinburgh met in the Anatomy class-room for the purpose of giving the demonstrator Dr. D. J. Cunningham an earnest of the good will they bore him, especially in relation to his recent marriage. Professor Turner presided, and Mr. J. Macdonald Brown made the presentation, which consisted of an extremely handsome epergne in silver. Dr. Cunningham suitably acknowledged the gift; and Professor Turner expressed his satisfaction with the success and object of the meeting.