mountain air towards the contraction of the pulmonary capillaries and a certain degree of interference with the pulmonary circulation.

Although we had no snow this last winter at Cannes, there was much more ice than usual. The effect of the sea as a source of heat was interesting to note; the mud and road-puddles being on several occasions frozen a few hundred yards from the shore, while soft and free from ice at the immediate seaside.

The temperature of the surface of the Mediterranean, taken daily at Cannes during the last six winter months, yielded the following monthly mean :

		1878-79.	1874-78.
November		58.3	61.2
December		56.6	57.6
January		55.4	56.1
February		55.7	55.5
March		58.1	56.3
April		58.4	59.5
Mean for the season		54.2	57.7

Hence it will be seen that the temperature of the sea at its surface was less available than usual towards checking the fall of temperature of the air; still, while the mean atmospheric temperature for the winter was 49.8°, we find that of the sea higher by 4.4°; so that the store of heat from the preceding summer taken up by the Mediterranean certainly assisted towards the maintenance of a comparatively mild winter temperature of the atmosphere.

The following table gives the number of days the wind blew last winter at Cannes, the notes being taken at 9 A.M. and 2 P.M.:

	Calm.		Light to Fresh.		Strong to a Gale.	
	9 A.M.	2 P.M.	9 A.M.	2 P.M.	9 A.M.	2 P.M.
November..	17	9	8	16	5	5
December..	13	14	13	12	5	5
January..	18	6	10	20	3	5
February..	18	8	4	15	6	5
March..	18	1	12	26	1	3
April..	11	1	15	21	4	9

Different localities at Cannes are far from being exposed to the wind in the same degree: a great many of them are sheltered from the dry westerly wind; and at some places hardly any wind at all may be felt, while a gale is blowing at sea. The mistral, or north-west wind, is, I understand, stronger and more frequent at Toulon and Marseilles than at Cannes.

Many people in England form an idea of the character of the air on the Riviera from that of the Atlantic seaside in the South of England; they compare it with that of Torquay or Bournemouth, or consider it perhaps somewhat like that of the Scilly Islands. This is quite a mistake; Cannes and the Riviera cannot compare with any place on the British coast. The climate is much drier, and the weather finer, with much more sunshine. A residence at Cannes, selected some little distance from the seaside, and somewhat above the sea-level, exerts a bracing influence, though not equal to that of Brighton or Hastings in the summer or early autumn. Such a climate has a decided influence towards equalising the circulation, and thereby checking congestions leading to hæmorrhage. As spring weather sets in, however, cases of hæmoptysis are occasionally met with; and I am in the habit of advising those who have suffered from pulmonary hæmorrhage to leave Cannes about the end of April. A residence at some sheltered place in Switzerland, such as Bex or Montreux, before moving further towards the British Channel, is I find attended, as a rule, with much benefit. Nothing, however, is so pleasing as the fine, fresh, bracing air of the British Isles, that welcomes one home like an old friend.

THE THERMAL SPRINGS OF ROYAT, LA BOURBOULE, AND CHÂTEL-GUYON.

By G. H. BRANDT, M.D., OPORTO.

THIS interesting group of mineral springs, situated in the middle of France, Department of Puy-de-Dôme (Auvergne), is remarkable for the great value of the distinct therapeutic action possessed by each of them. These three thermal stations possess waters of determined chemical properties; those of Royat are characterised by the large percentage of lithia they contain in combination with various salts, and iron and manganese, constituting a tonic alkaline water. La Bourboule springs are, no doubt, the most remarkable of all the known mineral waters in Europe, their chief characteristic being that they are the strongest arsenical waters yet analysed. Châtel-Guyon possesses the only laxative

waters in France; and they can compare favourably with the numerous springs in Germany and Hungary.

My object in bringing before the profession the importance of these three springs is, firstly, because little as yet has been said about them in England; and, secondly, to show the valuable therapeutic agents we have at our command to combat almost with certainty three classes of diseases, all of which are most intractable to the usual methods of treatment at present in use. Few diseases have been so thoroughly investigated as gout; in this class of disease, British authors have shown themselves pre-eminent, no doubt in some measure owing to its prevalence in England, and to the fact of many distinguished members of our profession having been themselves victims to the malady. Among the numerous authors, there is one whose name heads the list. Dr. Garrod has thoroughly worked out the subject, and all agree with his conclusions. His practical experiments have shown that there exists a solvent for the gouty deposits of urate of soda, and that that solvent is lithia; this question being settled, it remains to find out the best method of applying the remedy to the disease, so as to obtain the best results. Two conditions are requisite for carrying out this purpose; first, to place the patient in the most favourable conditions for treatment; secondly, to apply the remedy in its most potent form. To carry out the first indication, all, I believe will agree, that a healthy locality, with interesting scenery, rather a high temperature, where exercise can be taken in the air on flat ground, so as to encourage a free action of the skin, are most favourable conditions. To obtain the second indication—that of introducing the remedy under its most active conditions—few, if any, will deny that natural waters, containing the remedy in combination with other adjuncts prepared by nature, and offered to us under a proper temperature for external and internal use, are conditions we cannot obtain with all the wonderful chemical knowledge we possess; and why? Because there exists in these products of nature an active principle, invisible, imponderable, impalpable, which we cannot detect; and which gives power, action, animation, and certain virtues, which man can but poorly imitate and never equal.

Royat.—The numerous experiments made by competent observers, on thousands of patients under treatment at Royat, prove that the action of these thermal springs, applied in various forms and in a methodical way, are not only most beneficial in that obstinate disease called gout, but also in all the forms of the diathesis which modern pathologists have described under the name of "arthritis". I hope, by the end of next season at Royat, to be able to publish some interesting observations on the effects of mechanical massage on arthritic patients whilst under the influence of the mineral waters.

La Bourboule.—The mineral thermal springs of La Bourboule were known several centuries ago, as we see by a description, in 1540, of a rudimentary bath, in which patients from an infirmary in the hamlet of La Bourboule used to bathe. Owing to the small supply of water, the reputation of these springs did not extend much beyond the limits of the locality; and it is only within the last twenty years that their value has been justly appreciated, owing chiefly to the efforts of Monsieur Perrière, a most active and industrious man, who, by continued labours, discovered the great source to which he gave his name. This source affords an immense supply of water, at a temperature of 140 deg. Fahr., each quart containing over one-tenth of a grain of metallic arsenic, or one-third of a grain of arseniate of soda. Two other springs (Sedaigues and La Plage) of water of the same kind, at a moderate temperature of 91.4 deg. Fahr. and 86 deg. Fahr., afford about one hundred gallons per minute. These secondary springs have a great importance at La Bourboule, as they serve for graduating the temperature of the water from the great spring (Perrière). The advantages of reducing the temperature of a hot mineral water with water of the same kind, but of a lower temperature, are obvious; nothing is lost by evaporation, and it retains unaltered its active principles. These waters are employed as a drink, douches, lotions, inhalations, and pulverised as a spray. According to M. Lefort's analyses, these waters, when properly bottled, keep for a considerable time without losing their principal properties, owing to the small quantity of iron and organic matter they contain. Besides chloride of sodium, bicarbonate of soda, and arsenic, these waters contain other substances—such as iron, lime, silica, etc.—so admirably combined, that Dr. Gubler compares them to a true mineral lymph. Strange to say, the extraordinary cures performed by these waters for so many years only attracted the attention of chemists in 1852, when Thénard, surprised at their marvellous effects, investigated the cause, and discovered the presence of arsenic in a medicinal dose. Since then, a new era began for La Bourboule, and it now possesses one of the finest and most complete establishments in Europe.

The diseases in which arsenic, in the shape of arseniate of soda, plays such an important part are anæmia, chlorosis, certain forms of cutaneous diseases, malarial cachexia, chronic syphilis; and last, but not least, in

struma and all its varieties; in tuberculosis, and in certain forms of lung diseases in lymphatic individuals, affected by a slow form of phthisis, where circumscribed indurated spots are detected by percussion and auscultation—in fact, where caseous deposits or internal scrofula exist—these waters are most valuable, and very many successful cases have been recorded. Dr. Dobell, in his recent work on *Loss of Weight, Blood-spitting, and Lung-Disease*, refers to the good effect of arsenic observed by himself, quoting other authorities in favour of it. If we look back to remote periods, we find that the value of arsenic has been recognised by great observers, from Dioscorides down to Trousseau, de Haen, and particularly Isnard. Arsenic has been considered as one of the most potent tonics and alteratives known, and, being found in combination with other mineral products prepared in nature's laboratory, is no doubt the most perfect form of preparation. La Bourboule is situated at about two thousand five hundred feet above the level of the sea, in the centre of France, and in the midst of a most fertile and picturesque country. It thus enjoys a most exquisite climate, the air being pure and bracing, without being irritating; the place is provided with every comfort that an invalid can possibly require; these are advantages which must contribute largely to the efficacy of treatment.

Châtel-Guyon.—The thermal springs of Châtel-Guyon are the only laxative waters in France, and deserve special notice from their peculiar action. Situated at about four miles from Riom (railway station), on the borders of the Sardou, in a most picturesque part of Auvergne, full of interesting curiosities and pleasant promenades, it is a most inviting place. These waters owe their laxative properties to the presence of chloride of magnesium; several springs exist, of which some are used as baths. The diseases specially treated at Châtel-Guyon are obstinate constipation of the bowels, dyspepsia, congestion of the liver, and particularly cerebral disorders in which intestinal derivatives are indicated. For some years past, Dr. Barraduc has obtained most satisfactory results in cases of obesity.

The place affords excellent accommodation at very reasonable prices. It will be observed that I use the term "laxative" intentionally; for the action of these waters is not purgative in the common acceptance of the word, and differs essentially from the purging waters of Hungary and Germany; for these act like any common solution of sulphate of magnesia, purging the patient and leaving his ailment as it was before. The waters of Châtel-Guyon are laxative, and require to be administered in a methodical way to obtain the desired effect. The patient ought to take one or two glasses of the water before meals and *at meals*; if necessary, this quantity may be increased to three or four glasses *per diem*, until a free and regular action of the bowels takes place; this should be kept up for ten or twelve days, when the water may be gradually left off. It will be interesting to know the result of experiments now being carried out on dogs, at the École Pratique in Paris, as to the *modus operandi* of these waters. According to the last analyses, these waters owe their peculiar action to the large percentage of the chlorides of magnesium and sodium.

SURGICAL MEMORANDA.

THE TREATMENT OF SPINAL CURVATURE.

MR. E. CARR JACKSON is mistaken (see BRITISH MEDICAL JOURNAL, July 19th) in dating the treatment of spinal curvature by extension and hanging only one hundred years ago; for it is more than two hundred years since it was put into practice by Glisson, an English Fellow of the London College of Physicians. One can hardly take up any German surgical manual, which treats of orthopædic surgery, without coming across Glisson's swing being mentioned as one of the methods of treatment of spinal curvature. (See *Handbuch der Chirurgischen Praxis* of Dr. von Bruns, Professor of Surgery at Tübingen, vol. ii, pages 1217-1223). In the BRITISH MEDICAL JOURNAL of February 9th, 1878, I published a letter describing a similar apparatus to that of Dr. Sayre, which is found in several orthopædic institutions on the Continent. I cited those of Dr. Klapsch, Professor of Surgery at Breslau, and of Dr. Schildbach at Leipzig, where the gallows has been used for many years past. I also mentioned that Glisson was an English physician, who published a work *De Rachitide*, a second edition of which is dated 1660. The first edition was published in 1650, and was in Latin; but it was translated into English, in 1651, by Culpepper. In addition to other treatment, Glisson recommended (see English edition, 1651): "artificial hanging of the body.....which is performed by the help of an instrument usually made with swathing bands, first crossing the breast and swung under the armpits, then above the head and under the chin, and then receiving the hands by two handles, so that it is a pleasure to see the child hanging pendulous and moved to and fro by the spectators.

This kind of exercise is thought in many *waies* conducive in this effect; for it *helpes* to restore the crooked bone, to erect the bended joynts, and to lengthen the short stature of the body; it is rather a pleasure than a trouble to the child. Some, that the part may be more stretched, hang leaden shoes upon the feet and fasten weights to the body, that the parts may the more easily be extended to an equal length."

It is curious that, at Dr. Schildbach's Institution, I saw bags of sand, weighing several pounds, attached either to one hip or the other of the patients, according as one side of the trunk was required to be more extended.

I may remark that much unnecessary confusion is caused by speaking of cases of spinal caries (Pott's disease), and of spinal lateral curvature (scoliosis), indiscriminately, in reference to treatment by Sayre's plaster bandage and suspension. These diseases must be considered separately in properly appreciating the effects of Sayre's method, which is undoubtedly the greatest improvement yet effected in the treatment of spinal caries; while it is still an open question whether it is as successful in lateral curvature.

BERNARD ROTH, F.R.C.S., Brighton.

CLINICAL MEMORANDA.

A RARE CAUSE OF OEDEMA.

A RETIRED merchant, fifty-six years old, who is suffering from a very slight paralysis of the legs and occasional failing of memory, for which I had attended him from time to time, sent for me on account of symptoms of dropsy. I found his legs nearly double of their natural size, the skin of the abdomen oedematous up to the umbilicus, and the abdomen itself occupied by a large tumour, extending in the median line from the symphysis to the pit of the stomach. The whole of the pelvis was filled by a tense tumour, which fluctuated on pressure from above. He did not complain of scarcity of urine. I sent for a catheter, and evacuated nine pints of clear acid urine. For a few days, the bladder filled again to some extent; but the oedema began to diminish, and was gone at the end of a week. The daily introduction of the catheter and faradisation; iron and nuxvomica internally; injections of permanganate of potash while the urine was offensive and alkaline, as it turned immediately after the first catheterisation, have restored the bladder to its normal condition.

A. HESS, M.D.

INFANTILE DIARRHŒA.

I HAVE read Dr. Carey Coombs's article in the JOURNAL of July 5th, and I have no doubt that the facts which he states as the cause of the epidemic of infantile diarrhœa, which occurred in the town of Castle Carey last summer, are correct. From my own experience, I find that there is always more or less of an epidemic of infantile diarrhœa during the hot summer months amongst infants who are brought up by hand, especially amongst the children of the lower orders, the occurrence of which I believe to be almost entirely attributable to causes which, to a great extent, are preventable. The most common cause without doubt is, that sour milk is given to the infants; another, and no less frequent one is, that the feeding-bottles are never properly scoured out, so as to remove every trace of milk from the bottle after the infant has fed. If the same bottle be used two or three times without being previously well washed out, the milk will very likely be rendered sour, especially in the hot summer, when a very short period suffices to turn milk sour, and will assuredly cause diarrhœa and mortality amongst infants brought up by hand. I believe these to be the causes *par excellence* of infantile diarrhœa, from the fact that these epidemics rarely occur amongst the children of the well-to-do, where more care is taken to observe scrupulous cleanliness in the feeding-bottles, and in the use of milk which has not turned. May not the reason that the diarrhœa was arrested by the substitution of condensed milk for ordinary cows' or asses' milk (in Dr. Coombs's cases) be explained from the fact that the large quantity of sugar which is contained in the former renders it perfectly free from any acid, which probably might have existed in the cows' milk with which the children were previously fed? I have observed that diarrhœa hardly ever occurs in infants who are fed upon condensed milk, excepting, of course, the disorders of the alimentary canal, the result of dentition; and I believe that this freedom from the disease is entirely due to the fact that the condensed milk never turns sour. In infants who must be brought up by hand, I have seen such good results from its use that I much prefer it to ordinary cows' milk, providing I find by experiment that it agrees with the child's stomach. Another great advantage is its uniform quality.

WILLIAM ALEXANDER BUDD, L.R.C.P., etc., Exeter.