years of age, and was admitted into Bright ward under my care on March 1st, 1878. She is a married woman, of phthisical family. She has had six miscarriages and four living and said to be healthy children. Her last confinement was thirteen months ago. She appears to have suffered many things of many physicians, or rather surgeons, and has been an inmate of at least four of the metropolitan hospitals for typhus, rheumatic fever, erysipelas, etc. Seven months before her admission, she noticed pain in the right umbilical and epigastric region whilst mangling. She vomited, and the pain was persistent. She tacked about between various dispensaries for two months or more; and a swelling was noticed in the abdomen by Dr. Stocker. A fortnight before she came here, she was seized with severe abdominal pain; vomiting, which all along had been occasional, had become much more severe; and for three days she had been unable to retain anything on her stomach. The bowels had been very confined, rarely acting under intervals of from four to twelve days. She looked pale and ill; and on examining the abdomen we found a tumour to the right of the umbilicus, a little lower than in the former case, but like it giving some resonance on percussion, and there was still more resonance between it and the margin of the ribs. It was ill defined at its margin, very tender to manipulation, and not displaced by deep inspiration. The abdomen was not distended. The morning temperature was over 100° F.

The difficulties in diagnosis here are similar to those in the man. The tumour has not in this case any of the hardness present in the other, and its position is somewhat lower; and for that reason, the ascending colon and cæcum must be added as possible seats of disease; and conning over the case, I look upon it as one of localised peritonitis, the cause of which is not evident. Possibly it is due to some ulceration of the colon or displaced cæcal appendix, possibly to disease of the gall bladder. We gave her half a grain of opium and a quarter of a grain of extract of belladonna in a pill three times a day, with hydrocyanic acid and bismuth, milk, lime- and soda-water; and, under that treatment, she had no return of sickness, though before it had been constant. Since that time, the chief additional symptoms have been that the tumour has shifted up a little, so that if you examine it now you might think it to be attached to the liver; that she has had throughout very considerable abdominal distension; that the bowels have been decidedly confined; and that she has had decidedly more distension and abdominal pain on a fuller diet than on a restricted one. There is, moreover, a history that, on April 3rd, she passed a stone about as big as the top of one's thumb in her motion. The nurse stupidly threw it away, so we can only conjecture that it may have been a gall-stone. If this be so, a pretty good case would be made out, for gall-stones encysted in the gall-bladder; ulceration of the gall-bladder; adhesion by inflammation to the part around; ulceration into colon or some part of the intestine low down; and some amount of constriction of the bowel as a result of the inflammation. But this is all very conjectural, because she has repeatedly passed scybala, and the supposed gall-stone may have been no more than a hard fecal nodule; and our diagnosis practically rests where it did at first at chronic peritonitis catching up the colon and producing partial obstruction. The treatment has been such as I described, and she has latterly had copious enemata by the rectum. The prognosis in the case is somewhat gloomy; she is getting tired of the hospital, and will go out before long. She will then pay no attention to her diet or bowels, and will in that case soon get complete obstruction; though I believe if she could be kept under observation and guidance she might get well.

ABDOMINAL SECTION IN INTESTINAL OBSTRUCTION.

By T. R. JESSOP, F.R.C.S.,

Surgeon to the Leeds General Infirmary.

THE columns of the BRITISH MEDICAL JOURNAL could hardly be opened to the discussion of a more important question than the propriety of surgical interference in cases of acute hidden obstruction of the bowels.

When the papers of Dr. Clifford Allbutt and Mr. Pridgin Teale appeared in the number for January 11th, 1879, I was encouraged to hope that a difficult and perplexing surgical problem was about to be well thrashed out, that our leading teachers of medicine and surgery would eagerly seize the opportunity and freely enter the lists in so inviting a controversy.

The communications published in the JOURNAL of May 31st once again bring the question to the front—Is an exploratory operation called for when life is threatened by an acute concealed intestinal block, and, if so, under what specific circumstances?

More than twenty years ago, I assisted in making a *post mortem* examination of the body of a fine little fellow, the youngest son of one of our most respected citizens, whose disease, rightly and early diagnosed during life, was found to be a simple invagination at the ileo-cæcal junction. The case had run its course with much rapidity; there was a singular absence of all inflammatory changes, and by the simplest manipulation the gut was restored to its normal position. The conviction was forced upon everyone present, that an opportunity of saving the boy's life had been lost. Similar examples of intussusception are not rare; I can distinctly call to mind the particulars of three others, in which it was shown on *post mortem* examination that a reduction might readily have been effected.

Several cases are now recorded in which, by means of gastrotomy, invaginations have been successfully reduced; and no one I imagine would question the propriety of opening the abdomen in these cases when the diagnosis is assured, and after it has been made clear that other means have failed.

Many years ago, Mr. Seaton related to the members of the Leeds Medical Club the details of a case which, with the one given above, made a lasting impression on my memory.

A vigorous, healthy, and muscular man of thirty was suddenly seized with abdominal pain, vomiting soon followed, the bowels refused to act, and, in spite of appropriate treatment continuously applied, the patient died on the fifth day, with all the symptoms of a strangulated gut. On *post mortem* examination, besides evidences of recent peritonitis in the lower half of the abdomen, a coil of the ileum was found suspended from the front abdominal wall, not far from the umbilicus, by an organised band an inch and a half long. In his remarks on this case, Mr. Seaton writes: "To me it appeared, and still does, that, apart from the danger of opening the abdomen, the division of the band would have cured the patient."

Examples like this, again, will have been met with in the experience of many physicians and surgeons. I have selected the two for special reference out of a considerable number within my recollection, because they afford typical illustrations of conditions recognisable in some measure at least, even at an early stage, and by general acknowledgment admitting of relief from surgical interference.

It is only within late years that I have put the operation of laparotomy to the test, and as yet I have not met with a case of invagination suitable for or requiring surgical aid. I have had the satisfaction, however, of advising in one case in which death was averted by the timely division of a constricting band. It occurred in the practice of Dr. Land of this town, by whom the operation was most ably performed.

The patient was a man aged 45, who, for several weeks prior to his seizure, had suffered daily from abdominal pains and from retching before breakfast. Whilst walking in the street, on the 24th of October, 1875, he was suddenly attacked by most violent pain in the body, so that it was with difficulty he made his way home. Dr. Land was soon in attendance, and by means of fomentations and the subcutaneous injections of morphia he was able to afford some relief to the man's intense suffering. Daily, however, the symptoms of obstructed bowel became more marked:—uncontrollable vomiting, passing from cibal to bilious, and finally becoming distinctly stercoraceous; all escape of fæces and flatus by the rectum absolutely stopped; a gradually increasing distension of the abdomen. On the 31st of October, seven days from the commencement, Dr. Land asked me to share the responsibility with him. Matters had now become grave in the extreme. We both felt that, if unrelieved, the man had not twenty-four hours to live. The progress throughout had so closely resembled that of a strangulated hernia, without the external accompaniments, that we decided to recommend an exploratory operation. This was performed after such delay only as was necessary to obtain assistants and instruments. An incision large enough to admit the hand was made through the linea alba below the umbilicus, with the immediate result of a large protrusion of distended small intestine. After vainly searching the several internal rings, our attention was being given to the ileo-cæcal region, when, on turning aside the intestines, a band thicker than a quill and several inches long was discovered, and beneath it a loop of bowel. Just as the cord was brought from the midst of the distended coils to the surface, stretched over Dr. Land's forefinger, it was seen to give way. The exact attachments of the band were not sought for; there was no bleeding; the visceral protrusion was ever on the increase; there had already been a prolonged exposure, and we were naturally anxious to complete the operation. It probably sprang from a spot corresponding with the base of the sacrum or the last lumbar vertebra and stretched thence to the right brim of the pelvis. The intestines, deeply injected, were replaced, the wound was carefully closed by means of wire sutures, made to include the peritoneum, and the abdomen was covered with cotton wadding and encircled with flannel. The after-

progress of the case was all that could be desired; the symptoms of obstruction at once ceased, and he slowly convalesced. His only inconvenience now arises from a feeling of weakness in the anterior abdominal wall.

It is not, however, in cases of intussusception, which are, generally speaking, easy of diagnosis and readily followed through their various stages, that the surgeon will experience his greatest difficulty in deciding whether or not to interfere, nor is it in cases like Dr. Land's, so closely resembling a strangulated hernia in its symptoms, that he will feel himself most perplexed. There is, indeed, a fair amount of unanimity amongst authors as to the advisability of opening the abdomen in cases both of intussusception and of constriction by bands where the diagnosis is clear. But there are others, more numerous perhaps, of a much more doubtful character, in which the cause of the obstruction can only be determined by an exploratory incision; and it is in these that the question assumes its greatest importance—Is an operation of inspection called for? Are we right, that is to say, in opening the abdomen in an extreme case of bowel obstruction, when the nature of the obstruction is not apparent, on the possibility of finding some removable cause? The answer to this question, it appears to me, must depend upon the replies given to two others: What is the general experience of observers as to the causes of fatal obstruction? and are the chances of recovery in any degree lessened by an exploratory incision, where the cause is found to be irremovable by operation?

Mr. Teale answers the latter question by a decided negative. In four instances, I have searched the abdomen for a cause of obstruction; and, whilst I have not in any one of the four succeeded in discovering a state of things admitting of permanent relief, I have satisfied myself in each that death would, under any circumstances, have been inevitable; and in two, at least, life was prolonged considerably beyond what seemed possible at the time the operation was decided on. Now, I cannot share in Mr. Messenger Bradley's astonishment at this prolongation of life. It admits of a reasonable explanation, and it may fairly, I think, be reckoned upon in a majority of instances. No one would for a moment contend that, antiseptic precautions notwithstanding, an opening through the abdominal wall and a search amongst the intestines could be effected upon a healthy person without incurring considerable risk of inducing a dangerous or fatal peritonitis. But the conditions are wholly changed when we have to deal with viscera in a state of local or general acute inflammation; and it is probable that these are the states most likely to be met with in cases of obstruction from other than mechanical causes. To set against the element of danger introduced by the operation itself, are the facility and, if carefully managed, the safety, with which distension of intestines—a potent factor in the perpetuation of the diseased action—may be relieved, and the advantages to be gained by the removal of purulent collections and the general cleansing away of inflammatory products. One of the four cases above alluded to may serve as an illustration.

On the 8th of August, 1878, Messrs. John Teale and Waudby of Scarborough summoned me to join them in the treatment of the case of a young man visiting at the seaside from Sheffield. Mr. Waudby kindly furnishes me with a detailed account of the case, from which I extract the following. The patient, aged 33, was a fine strong man. On the 1st, 2nd, 3rd, and 4th of August, he complained of abdominal pains, at first in the region of the stomach, and afterwards in the right abdomen. Relief followed the use of an aperient. On the 5th, after having had a tepid bath, he played on the sands with his children, and subsequently, whilst sailing in a pleasure boat, he was seized with pains in the body so violent that he had at once to be conveyed to his lodgings. From this time, he began to vomit, and the bowels ceased to act. Mr. Waudby was now called in, and on the following day, August 6th, finding matters worse, he sought the help of Mr. John Teale. In spite of well directed treatment, the case assumed a graver aspect; and on the 8th, when I saw him, we all three considered him in the most imminent danger. Besides the block to the passage of both fæces and flatus, and the vomiting which took place after every act of swallowing, and which was, moreover, becoming decidedly feculent in character, the abdomen was tightly distended, the countenance was pinched and collapsed, and the pulse a mere flicker. After a most careful consideration of the case in all its bearings, we concluded that, as his symptoms might possibly be caused by an internal strangulation, and as his condition was rapidly becoming hopeless in spite of all that was being done for him, we were bound to offer him the chance of relief by operation. On opening the abdomen, it was at once evident that we had to deal with a case of diffuse purulent peritonitis. The intestines were everywhere thickly coated with semiliquid lymph, and numerous purulent patches were scattered in the furrows formed by the contiguous coils. In the neighbourhood of the cæcum was a large collection of pus, with which we interfered but little for fear of causing a fecal extravasation. So much

of the peritoneum as was exposed we cleansed, and, deciding that further interference was useless, we stitched up the wound. At the time of the operation, the severe symptoms had lasted three days, and he was then rapidly sinking. Mr. Waudby informs me he lived seven days after the operation, until the 15th; that the bowels acted on the 9th, the day after the operation, and that the patient always expressed himself as much relieved.

Upon a careful review of my own cases, I feel bound to affirm that my experience leads me to the conclusion that a patient's prospect of recovery is not prejudiced by an exploratory operation when the symptoms are found to be due to an acute local or general inflammation.

The former question is one which can be answered only by an appeal to the general experience of the profession, or by searching the records of reported cases, as has been done by Mr. Gay, Dr. Hilton Fagge, Dr. Brinton, and others, with the result of showing that in a large percentage of cases the cause of obstruction is purely mechanical, and, therefore, remediable. Invagination, concealed hernia, constriction of bowel by a band or omentum, volvulus, impaction of concretions or of foreign substances, together constitute, no doubt, a not inconsiderable proportion of acute obstructions; and typical instances of each of these present for the most part sufficiently distinctive features. Whilst, however, it is not a difficult matter to place on paper a categorical array of signs and symptoms, which shall apparently in the most satisfactory manner differentiate these various pathological states, and make it moreover plain as A, B, C, that to confound any one of them with, say a limited enteritis or a perityphlitis, would be an unpardonable blunder, every honest clinical observer must acknowledge the occasional correction in the *post mortem* room of a diagnosis confidently made and unwaveringly adhered to during life. One of my most valued friends, a physician who had, even whilst young, obtained a wide reputation amongst his professional brethren as a thoroughly accurate and trustworthy clinical observer and teacher, in the full belief that he was himself suffering from a concealed hernia in his right groin, begged for an exploratory operation; and yet, when the abdomen was opened after death, his request not having been acceded to, I found that the obstruction had been due to a typhlitis set up by the impaction in a patent appendix vermiformis of a fecal concretion, just as was the case in the little patient I attended with Mr. Pickles, as reported in the JOURNAL for May 31st.

Cases like these do not afford any very strong argument, it is true, in favour of operative interference, and yet it is fair to remark that the little boy did not die more rapidly than did my poor friend, upon whom no operation was performed. Both were equally beyond the aid of physician and surgeon; with or without operation, they were doomed, I believe, to a certain and rapid death. In the case of the child, we entertained the question of gastrotomy at our first visit, only to postpone it until it seemed clear that medical help was unavailing—until, in fact, he was sinking from obstruction of doubtful origin. Had we been able to diagnose the case correctly, we should not have deemed a surgical operation called for, unless, indeed, it could have been undertaken at a much earlier stage.

The question is—What proportion do these and other irremediable conditions bear to the more purely mechanical? Are not cases like Mr. Seaton's and Dr. Land's more common than is generally imagined?

On the evening of the 21st of October, 1877, Mr. Newstead of Leeds called for me to accompany him to a patient's house under the following circumstances. An engine-driver, aged 34, took his train as usual from Leeds to Hull on the 20th of October, arriving at the latter place in the evening. Whilst waiting for his return train to be made up, he was suddenly seized with violent abdominal pain, accompanied by vomiting and faintness. Utterly incapacitated, he was conveyed back to Leeds as a passenger; and when seen by Mr. Newstead at midnight he was still vomiting, in pain, and collapsed. Next day, the symptoms remained unabated; and when I saw him on the evening of the 21st, about twenty-four hours after the commencement of his illness, he was restless, tossing from side to side, vomiting a dark fluid with a steracaceous odour, moaning continually, and almost pulseless. There was some abdominal tension, but the man was too ill for us to determine the presence or absence of tenderness. He died at eight o'clock on the following morning, about thirty-seven hours after the seizure. We learnt from his wife that he had for some years been subject to attacks of diarrhoea, for which he had several times been compelled to seek medical aid. This information, the suddenness of the attack, together with the extreme and continuous prostration, led us to conclude that an old-standing intestinal ulcer had penetrated the peritoneum, and given rise to fecal extravasation—an opinion we deemed fully warranted by the facts, and confirmed by the rapid termination in death. On *post mortem* inspection, our diagnosis was not verified. Stretched horizontally across the lower abdomen, the appendix vermiformis was

attached by its tip to the sigmoid colon; a knuckle of ileum had become adherent to the posterior and upper surfaces of the displaced appendix; whilst beneath the whole, and tightly held down and constricted, lay a coil of jejunum, black with congestion. The adhesions were readily separated by the finger without the occurrence of any appreciable lesion in the gut; and the neighbouring peritoneum, though congested, was not marked by any inflammatory products. Who can doubt that, had it been possible on Mr. Newstead's first visit correctly to diagnose the state of things discovered in this case after death, the operation of gastrotomy would have offered a very probable means of cure? Are such cases as this and the others I have mentioned more frequent than is generally supposed? I have myself seen several others; my medical friends have related to me many examples. What is the general experience of the profession? If it be in accordance with my own and many of my colleagues and friends, and if I am right in concluding that, under circumstances unfavourable for operation, the exploratory incision, by enabling the surgeon to relieve tension and to cleanse away harmful inflammatory products, is found, on the whole, to prolong life without adding to the sum of suffering, the exploratory incision ought to be more frequently resorted to than heretofore. At any rate, I protest against the term "unjustifiable" which has been by some applied to the operation, and apparently upon theoretical grounds only.

A CASE OF INTESTINAL OBSTRUCTION SUCCESSFULLY TREATED BY PUNCTURE OF THE SMALL INTESTINE.

By W. H. BROADBENT, M.D., F.R.C.P.,
Physician to St. Mary's Hospital.

THE following case may be related as one of the series now being published illustrating various methods of treatment of intestinal obstruction.

In February 1877, I was called by Mr. Rayner to a maiden lady, aged about 60, who presented the usual symptoms of intestinal obstruction—severe tormina, vomiting, and great distension of the abdomen. The seat of the obstruction was made out to be in the neighbourhood of the cæcum, the transverse and descending colon being empty. Opium and belladonna were given, food withheld, and enemata administered; and, the pain and vomiting being quieted, an examination of the rectum was made at a subsequent consultation, when a solid mass was felt pressing down into the pelvis on the right side. It was pushed up as far as possible, and very shortly flatus and fæces began to pass. It was ascertained, on inquiry after the examination, that about nine or ten years previously, the patient had been tapped, and that the contents of a suppurating ovarian cyst had been withdrawn. The shrivelled cyst had formed a tumour in the right inguinal region, where she had been accustomed to feel it, and where it was found after the attack. The obstruction was attributed to displacement of this tumour downwards, giving rise either to pressure upon the bowel, or more probably to dragging by adhesions.

There was a slight return of obstruction in March, and again in June, brought on by imprudent exertion; in the autumn, she had a more severe and prolonged attack while at the seaside.

On May 14th, 1878, I was again called to this lady by Mr. Rayner. She had then been suffering from complete obstruction of the bowel for several days, which persisted in spite of the treatment which had on the several previous occasions been successful. The tumour formed by the old cyst could now be indistinctly felt in the right inguinal region notwithstanding the great distension of the abdomen, and was only just reached by the rectum; the rectum was held open by adhesions, so that on passing the sphincter ani the finger moved freely in all directions in a large cavity. A vaginal examination was impracticable, on account of the contracted state of this passage.

The treatment was continued: in addition to the opium (gr. j) and belladonna (gr. $\frac{1}{3}$) pill taken every three or four hours, and poultices with opium and belladonna over the abdomen, morphia was injected hypodermically. No food was given by the mouth, but enemata of beef-tea and brandy were administered every three or four hours, while once a day a copious enema of water or gruel was employed, for the double purpose of contributing to the relief of the obstruction and of washing out the rectum and removing particles deposited from the beef-tea, these being liable to become acid and set up irritation which causes the nutrient injections to be expelled.

No good result was obtained; and at length, after nearly a fortnight of anxious watching, and when complete obstruction had lasted three

weeks, the intestine was punctured by a long aspirator-needle. The aspirator was used at first, but was found to be unnecessary; an enormous amount of gas gradually escaped, the tension of the abdominal walls, which had seemed ready to split, was relieved, and two days later (May 20th) fæces and flatus began to pass naturally.

Another attack came on in January 1879, and when I saw the patient with Mr. Bridges on January 4th, complete obstruction had existed for several days. The attempt was made to procure relief by enemata, opium, hypodermic injections of morphia, external applications of opium and belladonna on poultices, and in effect by treatment similar to that already described, but in vain. Ice also was applied over the right iliac region; it gave relief from suffering, but did not set the bowels free. The patient begged from the first that the gas might be drawn off, as in the previous attack; and, as there was no improvement, this was done, without waiting for the same length of time, on January 8th. The distension was removed, and during the following night a copious evacuation from the bowels put an end to the danger.

At this time, indistinct fluid vibration was detected in the right inguinal region instead of the solid-feeling tumour, and, on the subsidence of the abdominal distension, it was well marked. It was accordingly determined that on any recurrence of obstruction an attempt should be made to remove the cause by aspiration of the cyst, which was evidently refilling after an unusually long interval. The recurrence happened in May of this year, and I was called to the patient on the 7th by Mr. Gawith, to whose care she had been transferred. She had been suffering from bronchitis, and still had a severe cough. There were all the indications of obstruction; the signs of fluid in the right inguinal region were more distinct; and, on examination *per rectum*, an elastic tumour could be felt high in the pelvis, from the surface of which projected two or three small secondary cysts. On the 9th, a moderate sized aspirator-needle was plunged into the cyst, and about six pints of dark brown fluid were drawn off without any unfavourable results. At first, it seemed as if the desired effect were about to follow: a little flatus was said to escape when the enemata were used, and a small quantity of fecal matter was felt in the rectum on examination. This promise was, however, illusory; and, as the distress increased, the intestine was once more punctured on May 20th. A considerable amount of gas had escaped; and the extreme distension was relieved, though not to the degree required, when, in the act of coughing, the coil in which the needle was planted was suddenly displaced to the left and slightly downwards, carrying with it the needle, which now lay flat on the abdominal wall. It was immediately withdrawn; but this was followed by a free escape of gas, which first issued with a hissing sound from the aperture in the skin, but soon rushed into the subcutaneous cellular tissue, and produced emphysema, which rapidly spread. This was circumscribed by pressure made by the hands, which also had the effect of directing any gas which had escaped into the peritoneal cavity to the internal opening in the abdominal wall. Another needle was at once plunged into a neighbouring coil of intestine, to take off pressure and prevent gas from passing into the coil from which it had escaped after withdrawal of the needle. Finally, when all issue of gas from the intestine appeared to have ceased, the subcutaneous emphysema was removed by squeezing towards and out of a puncture made by a lancet.

Strange to say, no bad effects of any kind followed; but the pressure had not been sufficiently reduced to permit of the removal of the obstruction. The distension soon became as great as ever; and, with considerable hesitation, the operation was repeated four days later. More gas escaped; the abdomen became soft and yielding, almost flaccid; when exactly the same accident occurred again, the patient coughed and displaced the intestine, the needle had to be hastily withdrawn, it was followed, as before, by gas which escaped from the puncture and rapidly diffused itself under the skin, where its progress could be both seen and felt. Similar precautions were taken, and happily no inflammation followed, but, on the contrary, a copious stool, and the bowels have since continued to act.

Since the above was written the patient has again suffered from obstruction, which was not relieved till the intestine had been punctured. This was done on August 1st with the same fortunate result as before.

I have recommended and practised puncture of the intestine several times in intestinal obstruction, and have never seen injurious effects. The precautions which I consider necessary are the following:

1. To secure, if possible, absolute freedom from peristaltic action of the bowel. This is done by giving an extra dose of opium by the mouth, or a considerable hypodermic injection of morphia, or both, three or four hours beforehand. No food of any kind should have been taken for some time.

2. To select, if possible, a coil of intestine which shall contain only gas, and not liquid. This will be in the jejunum, and is to be found above the umbilicus rather than below it. An indispensable condition