be no doubt that the drug had a direct influence in checking the attacks. In the present instance, this could hardly be due to the elimination of any material from the blood, since there was no uræmic element in the causation of the convulsions. Nor did the effect upon the pulse, as shown by the sphygmograph, appear to be sufficient to make the explanation probable that its mode of action was by causing relaxation of any arterial spasm. As an oxytocic, the pilocarpine appeared in this case to be absolutely without any effect.

No further convulsions occurred after delivery, and the patient recovered well.

INTRA-UTERINE MEDICATION.

By CLEMENT GODSON, M.D.,

Assistant Physician Accoucheur to St. Bartholomew's Hospital; Honorary Secretary of the Obstetrical Society of London.

It is with a feeling of anxiety that I see, in the *BRITISH MEDICAL JOURNAL* of August 30th, a series of abstracts of papers which were read at the annual meeting in Cork, advocating the treatment of certain uterine complaints by means of medicaments applied to the cavity of the body of the womb. This *JOURNAL* has such an enormous circulation among all classes of the profession, that the mere heading will attract the notice of many who have but little time to do more than glance at the contents of the papers; and I fear that, having ascertained that carbolic acid, iodine, etc., are useful agents in the treatment of uterine discharges, many will fly to put their efficacy to the test, without having learnt to discriminate the class of cases applicable to them, or having attained the skill necessary to employ them. There is much in fashion. A century ago, every one was bled; now, venesection is hardly ever practised. We have passed from one extreme to the other. A few years since, in gynaecological practice, division of the cervix by the metrotope was constantly performed by some physicians; now, this operation is but rarely undertaken; indeed, we hear, at least in one quarter, that sewing up the gaping os uteri is all the rage. Lately, there has been a discussion on the use of the forceps and its alternatives in lingering labour, before the Obstetrical Society of London. Many eloquent discourses have been contributed, and the subject has been ably discussed by several of our eminent leaders in the school of midwifery. But, to sum up the whole, at what have we arrived? The general impression gained, is that the forceps is used far less frequently than it might be with advantage to both patient and practitioner. The result of this will be, unless I am much mistaken, that an authority has been granted for the frequent use of the instrument, and that we shall find the evil effects of this from its indiscriminate and unskilful employment. In the out-patient female department of St. Bartholomew's Hospital, where I see from fifty to seventy new patients per week, I have unusual opportunities of observation. After having made an examination, I am frequently led to inquire: Were you delivered with instruments in your confinement? And, in nine cases out of ten, the answer is in the affirmative. The reason why I put the question was that I found the cervix deeply fissured up to its junction with the vaginal wall, along which extended a cicatrix. This is a clinical fact, which testifies to the truth of the remark made by Dr. Henry Bennet in his speech, and objected to by Dr. Barnes in his reply, at the Obstetrical Society's discussion.

But, with respect to uterine medication, the results are likely to be much more serious; and I have already met with instances. If an application to the cavity of the body of the womb be required, there should be an amount of *endometritis* calling for it; but, in how many cases of discharges from the os uteri does the exudation come from above the canal of the cervix—the so-called internal os? I venture to say, in very few. The probe, covered with cotton-wool saturated with carbolic acid, is passed along the cervical canal up to the fundus uteri, and in many cases good results are obtained—not because the fundus uteri has been reached, but *because the application has been made to the canal of the cervix*; and, had a zinc-alum point been placed in this situation, and the uterine cavity have not been reached, the result would have been equally satisfactory, without the risk of danger possible to occur from probing the cavity of the womb. This brings me to the nature of the evil consequences.

One which I have met with is abortion. The possibility of existing pregnancy has not been thought of. The patient may have gone but a few days over her time for menstruation, and, not having been very regular previously, little heed is paid to this. Nevertheless, she may have conceived immediately after her last period, and, if even she have not quite reached her next menstrual epoch, she may yet be pregnant when the application is made. Another danger is that of inflammatory mischief in and around the uterus. It is well known how the simple passage of the ordinary uterine sound is liable to give rise to this. Surely, then, a

probe saturated with carbolic acid or other agent is equally if not more likely to produce such results.

The points, therefore, that I would urge upon my professional brethren are these. Before deciding to make applications to the cavity of the body of the womb, be satisfied that this part is affected by disease, and that it is not the cervical canal only that is involved. Without having determined this, it would be just as irrational to pass a medicated bougie into a man's bladder for every purulent discharge escaping from the meatus urinarius.

The papers to which I have referred, as published in the *JOURNAL*, fail to my mind to show sufficiently clearly the symptoms which give proof of the existence of endometritis. Then, the application should never be made to a sensitive uterus in which inflammation is likely to ensue, or pelvic inflammation or abscess may result; and a probe should never be passed into the uterus of a childbearing woman about the time of an impending menstrual period; indeed, it would be well to make a rule not to probe the cavity within a week of the catamenia, but to limit the time for intra-uterine medication to the fortnight subsequent to the cessation of the flow.

OÖPHORECTOMY PERFORMED SUCCESSFULLY FOR MENSTRUAL EPILEPSY.

By LAWSON TAIT, F.R.C.S.Ed.,

Surgeon to the Birmingham Hospital for Women.

THE case of E. E. has already been published in the *Obstetrical Journal* for May 1873, and in my book on *Diseases of Women*, as a typical case of ovarian atrophy arising from exanthematic oöphoritis: a disease to which I was the first to call attention many years ago, as occurring very frequently as a sequel of scarlet fever and small-pox.

Her history after May 1873 is, that the efforts to induce the return of the menstrual flow by means of galvanic pessaries were persevered in, after several relapses, till at last everybody concerned was wearied of them. The patient ceased her attendance at the hospital, and I heard of her only occasionally to the effect that her epilepsy was becoming worse and worse. One of my critics, in alluding to this case, broadly stated that I had mistaken a bad case of hysteria for one of epilepsy. I wish he had seen the poor woman as I saw her on February 5th, 1877, in an attack of epileptic mania, covered with bruises of all kinds. I urged her friends to commit her to an asylum; but to this I could not persuade them, as for at least one week in every month she was quite rational and comparatively free from fits. During the other three she had fits almost every day; and during the menstrual week, when a faint show generally appeared for one or two days, her fits were incessant and her delirium uninterrupted.

This state of matters lasted till August of this year, when her friends, completely worn out by her, seemed at last to agree that she should be placed in an asylum. Before doing that, however, I urged them to let me cut out the ovaries, in the hope that this might arrest the disease by removing what I believed to be the cause. To this they readily consented, and I performed the operation on August 11th. There was no particular difficulty about the operation, save that the patient was very fat. The ovaries when removed presented all the characters of the atrophy peculiar to inflammatory change of their true gland-structure. (*Diseases of Women*, p. 215.) Mr. Alban Doran has favoured me with a report upon the ovaries removed, which completely confirms the views of the pathology of this case which I have expressed in my former publication of it. "The elongated cells of the stroma are larger than in normal ovaries; and there are few vessels, the hypertrophy of the walls of those that remain, and the bundles of fibrous tissue, point to a cirrhotic

Ovaries showing naked eye appearances of atrophy resulting from exanthematic oöphoritis, removed from case of menstrual epilepsy.

change following the exanthematic oöphoritis." One of the ovaries is now in the Museum of the College of Surgeons.

The result of the case so far is very satisfactory, for the fits have entirely ceased. Three days after the operation, there came on the