REMARKS ON ANIMAL VACCINATION.

A Paper read at the Conference of the Parliamentary Bills Committee of the British Medical Association, on Animal Vaccination, December 4th, 1879.

By E. WARLOMONT, M.D.,

Member of the Royal Academy of Medicine of Belgium; Director of the State Vaccine Institute at Brussels, etc.

GENTLEMEN,—In rising to speak before this assembly, I feel that my temerity is great in discoursing to you on vaccination in the country of its birth, under the sky which has seen the growth of that to which we owe the most stupendous benefit with which the genius of man has ever endowed mankind. Permit me, before commencing my subject, to fulfil a religious duty in at once saluting the great name of Jenner. Jenner was struck with an idea which doubtless would have passed through a commonplace mind, but which was destined to fructify in his intellect. What were, in fact, the elements of this discovery which was to protect the world from the most frightful of scourges? They were of the simplest. It was known that an attack of small-pox generally preserved from an attack of the same disease; that, in short, it only once invaded the same person. Now, Jenner had by chance discovered another malady having in its fundamental external characteristic the closest analogy to the small-pox pustule. The distinctions between this affection and that of which it was the type, and which it so strikingly resembled in its outward aspect, were the almost complete absence of general reaction, and its entire powerlessness to spread the infection itself. To give intentionally this mild disease, which in the animal economy was equal to an attack of true small-pox, and thus to create in it an immunity during a certain duration of this specific process—such was the idea which inspired Jenner, and from which he evolved the marvellous fact of vaccination. The history of this discovery is too well known to need to be here related. I only wish to point out—and this will not at all diminish the honour which will eternally attach to a venerated name—that Jenner was only able, notwithstanding the trials made during his lifetime, to lay down the bases of his method, and to form hypotheses as to the extent of its application, the duration of the acquired immunity, and the necessity of renewing the inoculation after a certain interval; in a word, he was not able to lay down the conditions of revaccination. This task was left for future generations. To enlarge the application of the method, to make it benefit the greatest possible number of persons, to create a legislation for the purpose of procuring this extension, to multiply the sources able to furnish the preservative matter—such were the duties left to us. In accomplishing this task, in completing the work, the successors of Jenner do nothing for their own glory; the little that they add to the edifice with which this commanding genius has enriched the world will only serve to still further increase his renown. For myself, in coming here to speak of vaccination, I have no other ambition.

Primary vaccinia is the product of an affection capable of being developed spontaneously in the horse or in the cow (in both, perhaps, but this point is not quite settled), whence the name of horse- or cow-pox which has been given to it. Transferred to the human subject, it develops in a regular fashion; and after a lapse of time, varying between six and seven times twenty-four hours, the vesicle which it has caused furnishes in its turn inoculable lymph. The exceptions to this rule are very rare; few subjects are insusceptible to a first inoculation, if the lymph be of good quality and of proper age. The rule, on the contrary, is that the power of the lymph is weakened the further we get from the moment that the vesicle begins to enclose it. If it be inserted from a vesicle of more than eight days from the date of vaccination, failure is to be feared. Thus, by proceeding with care (and Jenner had already given some instructions with regard to this which have never been altered) each subject can be made a factory of the material for protecting others as it has just been itself protected. Thus the process maintains itself, following a course which carelessness alone can arrest. This is, in fact, what is known as vaccination from arm to arm.

In speaking of this for the first time, I wish to make a profession of faith, which will be sincere, as will be everything that I shall have the honour of saying to you. Although I have come to speak to you on the method of so-called animal vaccination, I am anxious, above all things, to tell you the whole of my ideas as to the part which it is ex-

pedient to assign to each of the two methods. In my view, animal vaccination should have no tendency to forcibly supplant vaccination from arm to arm. The latter ought, on account especially of the facilities afforded by it, principally in the countries where the population is much scattered, to be scrupulously observed. They are, in fact, two sisters, which must not be separated.

This proposition is based upon a fact without which it could not be maintained—the perfect identity between the lymph of the child and of the calf, so far as regards their active principle. The vaccine matter is composed of two principal elements, a vehicle, serum, and some special granulations which represent the vaccinal power. This last fact has been quite recently placed beyond doubt by the experiments made at the vaccination establishment of Amsterdam by MM. Carsten and Coert, with the assistance of M. A. H. Parcau. These gentlemen report as follows.

“Some animal lymph” “was filtered through filter-paper, and through some small plates of porous porcelain, by the help of the pneumatic machine. Microscopical examination showed us that the filter-paper retained the epithelium-cells, the fibrinous clots and other solid particles, but that it allowed to pass globular molecules, which are found in great quantities in animal lymph, and which are supposed to be schizomycetes. The porcelain plate does not allow either these molecules or any other morphological particle to pass. We repeated these filtrations in different ways, and we were able to ascertain on each occasion that the inoculations practised on calves with the serum, deprived by filtration of its globular molecules, only gave negative results. When, on the contrary, these molecules had been preserved in their vehicle, the inoculations were constantly successful. We have mixed and washed the lymph with plenty of water, to separate these molecules from other solid constituents that might adhere to them, and which might have been considered as equally endowed with some virulent property; and we have then passed through the porcelain all that the latter allowed to pass through. Inoculation with the matter remaining on the filter was again followed by positive results. Thus it is evident that the virulence is inherent in the globular molecules.”*

These results fully confirm those which M. Chauveau had already obtained by other means with regard to human lymph.

The identity is therefore perfect as to the nature of the active principle of the lymph, whether it is derived from the calf or the child. It is in both cases a figurate granulation, perhaps a microbium, suspended in serum. This serum, however, differs a little; it is more plastic in the calf, perhaps because the lymph yielded by the latter has had to be expressed by a force which expels more fibrine; hence a tendency to coagulation, which renders more difficult its keeping in the liquid state, or rather its expulsion from the capillary tubes into which it has been put.

This identity is established, if possible still more completely, by my own personal experience. During six months, I successively transmitted the same lymph from the calf to the child, from the child to the calf; and, after twenty-five transmissions thus crossed, I found the lymph still to possess all its original qualities. This fact, well established in the way that I have just related, enables us to formulate the proposition that human lymph and calf-lymph can render each other mutual aid and assistance.

But, it will be asked, if the two lymphs are of equal value, why call to the aid of humanised lymph, the supply of which never fails, the assistance of animal lymph? The answer is that this help is especially necessary to satisfy doubts, fears, imputations, and perhaps prejudices. It has been said that lymph degenerates by passing through the human organism; so that what was introduced by Jenner has now lost a part of its power. This is only a simple assertion, and its truth has not been proved. There is perhaps some merit in my acknowledging it, but I myself, rashly I fear, but honestly, brought forward this idea some years ago. At the present time, nothing proves to me, nothing tells me, that lymph can degenerate. As long as its globular molecule—its microbium (?)—is preserved and is healthy, there is no ground for asserting that it has degenerated. Have small-pox and syphilis lost their vigour by lapse of time? However this may be, the feeling exists, many physicians share it, and I think it must be respected. If to their mind animal lymph promises to those vaccinated with it a more certain and more durable immunity, what right have we to refuse it to them? Again, it has been said, and this is much more serious, that human lymph taken from a syphilitic subject can transmit syphilis to the vaccinee. The fact is only too true, and certainly it is not in England that it is necessary to strive to establish it. Vainly is it maintained

* *Animal Vaccination in Holland: Some Provisional Experiments made at the Vaccinating Station at the Hague on the particular qualities of Vaccine Virus.* By Carsten and Coert (Medical Congress at Amsterdam, 1879).

that lymph from a syphilitic subject is innocuous, if it be taken with the necessary care, and without the admixture of blood. But who of us, gentlemen, would like to expose his children or his grandchildren to such a danger?

The physician ought, then, to guard himself against such a possibility; and he can indeed only do this by imposing upon himself the obligation only to make use of the lymph from infants he has under his own eyes, and of the state of whose health, as well as that of their parents, he can be positively assured. We will not insist too strongly upon the necessities in this direction; but we believe that the practitioner can in no other way ensure complete security. This being so, he ought not to be allowed on his own responsibility to make use of any human lymph collected by any other person than himself; and he should impose upon himself as a line of conduct that, when a family requests him to perform vaccinations or revaccinations, the family should itself provide him with the inoculating matter. In this way, and in this way alone, can the physician protect himself from future responsibilities and claims. I am fully aware that this method of procedure faithfully followed would have the result of increasing and exaggerating the mistrust already existing in the public mind. But what is to be done? Physicians are by no means anxious to offer themselves as a holocaust to the ill-feeling of families, always disposed to attribute to the lymph employed all the maladies which may subsequently attack their children; and the prudent course which I recommend to them is the only one which can at once really cover their responsibility, and at the same time screen them from unjust reproaches.

I know full well that I shall be accused of an excess of precaution. At vaccination establishments, I shall be told, the collection of lymph from the arms of infants is done under conditions calculated to offer every security; the persons charged with this office quite appreciate the importance of the duty confided to them, and not a tube of lymph escapes their hands which has been taken from a doubtful source. What a mistake is this assertion! Does not the moment always arrive when the most attentive, the most scrupulous of men, relaxes his vigilance? So, when the question is of human lymph, this rule must always be rigorously observed: Do not vaccinate any subject unless you can take lymph for its vaccination from the arm of a child present, whom you have yourself carefully examined. Now, this condition cannot be always observed. Too often stocks can only be kept up by means of preserved lymph collected from subjects intended to serve as vaccinifers. These conditions apply indiscriminately to all countries in which vaccination is voluntary, but there recourse to it can be dispensed with until opportunity serves. In those countries in which vaccination is compulsory, it is not the same; it is necessary to submit within a fixed period to the lancet of the vaccinator, one's children or grandchildren, without, in the majority of cases, having the time—we are speaking especially of the poor—the power, or the means to inquire if the matter to be used for vaccinating fulfils all the conditions of harmlessness which one has the right to demand of it. I do not hesitate to repeat—I say repeat, because I have already said and written many times—that this is not right. In the countries where vaccination is compulsory by law, the duty of the State is to put in the hands of the public a lymph which should be free—I will not say from all adulteration—but from all suspicion of diathetic adulteration. Like Caesar's wife, it ought always to be above suspicion. I know that nothing is so blind as the prejudices of the public; but, here, are these prejudices inconsiderate? We see a little pink and white baby, the idol of all who surround it; it is radiant with health and beauty. You take its arm and instil into it a strange principle, of which you, the physician, are doubtless certain, but of which the family knows neither the source nor the purity. And you do not like that the family should ask for every possible guarantee; and that, not obtaining such, it should rebel and resist. But it is in human nature; and you all know it well—you who have witnessed for some years a rising tide of opposition to the Vaccination Acts, which you either disregard or despise. If these prejudices be not unreasonable, then give to the public the satisfaction which it demands. Give it the calf and its lymph; give it, at the same time, that confidence, that security which it has the right to exact from you, and you will see, from day to day, this opposition give place to the most perfect submission; for confidence in vaccination is a matter of conscience for every mother.

But are we, then, in the right in saying to families, "Come to the calf, it will not betray you?" The reply cannot be for a moment doubtful. We answer with the most entire confidence: "No, the calf will not betray you; you can go to it *cito, tuto, et jucunde*." It is not, however, that contrary opinions have not been expressed. It has been said, amongst other enormities—and the allegation has come from those very men who scout the possibility of vaccino-syphilis, which makes the matter more absurd still—it has been said by those who do not believe

it themselves, that lymph taken from a tuberculous calf can transmit tuberculosis. They did not take the trouble to ascertain whether there had ever been found a microscope powerful enough to reveal the presence of a tuberculous element in the vaccine vesicle; and, in default of such an authentication, whether tuberculosis could be transmitted otherwise than by the engrafting of that disease itself. All this mattered little; it was necessary to cast a doubt on the new method. They did not stop to think that this suspicion, if it had any foundation, must fall with all its weight upon humanised lymph itself. Let us say at once that the argument was not long maintained, and that those who had put it forward now practise animal vaccination.

Vaccinal glanders has also been talked of. The following article appeared recently in the *Lyon Medical*, June 22nd, 1879, and was written by an author otherwise sensible, but sufficiently careless and thoughtless to publish it, without the least proof in support of its assertions.

"EPIDEMIC OF VACCINE-GLANDERS.

"The following fact has seemed to us too serious not to be brought immediately under the notice of our readers, for it is of a nature to singularly damp the zeal of the apostles of animal vaccination. On April 22nd last, the authorities of San Quirico d'Orceia asked a semi-official vaccination committee at Rome for some tubes of lymph, which arrived on the 24th. On the 26th and 28th, the local doctors vaccinated with this lymph thirty-eight children, all aged less than twenty months. Whilst they wereawaiting the incubation of the vaccine pustules, they soon perceived that they had inoculated one of the most horrible of maladies, and that they were the involuntary authors of a real massacre of the innocents. The gentleman who sent these particulars to the *Gazzetta d'Italia*, betook himself to San Quirico. He saw the victims. He observed vast phlegmons, laying bare the muscles and penetrating into the joints, accompanied by eclamptic symptoms. To him it appeared to be very probably an epidemic of glanders. If this really was the case, all the responsibility must rest upon the Roman committee. One cannot excuse men of skill who use a glandered calf as a vaccinifer. If it were a question of another malady, which it was not possible to diagnose, animal vaccination does not appear less compromised." This is what it was wanted to bring out. The article ends thus: "In any case, we shall endeavour to make known to our readers the results of the inquiry, which is being actively pursued."

This inquiry has, in fact, been made, and it has resulted—so runs a letter, dated June 15th, addressed to Senator Professor Palasciano by Commander Constanzo Mazzoni, member of the Board of Health sitting at Rome—that the facts above reported have been audaciously misrepresented: the inquiry having established the perfect state of health of the calf that had furnished the inculpated lymph. This is what happened. Certain institutes in Italy still deliver—(though it is a detestable practice, against which we have always protested), pustules cut off the calf, and send these scraps of skin to larger or smaller distances without any care about the state in which the pustules may arrive, trusting no doubt to the sense of smell of their correspondents. In the case of San Quirico, such pustules, and not tubes, as stated in the *Lyon Medical*, were sent to the local doctors, who did not use them for five or six days after they had been cut off the animal. And this in the height of summer! The vaccinators, indeed, admitted that from the pustule was exhaled a musty smell (*mucedo*), but this did not deter them from using it. As a matter of fact, if they had wished to purposely propagate septicæmia, would they have proceeded otherwise? The practice, we repeat, is deplorable, and it is inconceivable that responsible persons still consent to adopt it. We know that those who practised it were indeed the first promoters of animal vaccination, and as such, claim indulgence at our hands; but we are amongst those who think that, from however exalted a source an idea springs, it is not on that account infallible. If the method of Negri had not been freed by us from its original conditions, it would never have been able to acclimatise itself far from its place of origin. It is in reducing it little by little to its most simple elements, and in approaching in its operations to the traditional method, vaccination from arm to arm, that we have made it a simple proceeding by a logical and easy application. The cutting off of the pustules, which has a long time encumbered it, has never troubled us. We have never put the method in practice. The occurrences at San Quirico d'Orceia prove sufficiently well that we are right in prohibiting it. A great thinker and justly celebrated author, Alexandre Dumas *fil.*, has said that, to spare himself illusions, he had taken the part of considering men as being naturally bad. If he met a good man, it was an agreeable emotion that he felt. This emotion the *Lyon Medical* has not given us. We still await, as we have done since June 29th last, its publication of the result of the inquiry.

But may we not ourselves be ranked amongst the mischievous persons

of whom we have just spoken, and have we not perhaps sometimes given proof of it, in admitting too readily, and spreading abroad without sufficient proofs the statements as to vaccino-syphilis, of which we ourselves believe only half? We are not quite sure of our perfect innocence in this respect, and we ask ourselves if, in echoing the stories—often untrue and nearly always exaggerated, we have not made ourselves culpable accessories to this discredit. If we have been attacked by such a mania, let us cure ourselves of it as soon as possible.

The possibility of the transmission of anthracoid affections by animal vaccination has also been spoken of; but, besides that nothing is rarer than these affections in very young bovine animals (and it is these only that our method makes use of), besides that such an affection could not escape the supervision of the experts who inspect the markets, it has been perfectly proved to us that proper vaccine pustules cannot be developed upon animals suffering from this generalised disease. This is, in fact, one of the most characteristic features of the question. In the vaccination of cows, the appearance of the pustules alone shows the state of health of the vaccinifer calf. From the moment that the animal suffers, the vaccine pustule suffers, and it is doubtful whether a vesicle could ever prosper upon an anthracoid subject. As a general rule then, one ought to be able to utilise without fear lymph taken from good pustules, because proper pustules never rise on sick animals. We know that it is not the same with regard to the infant; a subject highly syphylised can show vaccine vesicles perfectly irreproachable in appearance. Let us add that, less favoured than the advocates of septicæmia, the champions of anthrax have yet to find a single fact to bring against us, and that their objection is up to the present time purely speculative.

Syphilis remains, and of that, and all the world is in accord on this point, we have no fear. Civilisation has not yet penetrated so far into stables as to leave there its most characteristic mark; and to the humble and pure calf may equally be applied the old adage: "*La plus jolie fille du monde ne peut donner que ce qu'elle a.*" But the objectors to animal vaccination are not content with this adage. They have wanted to prove that the innocent calf can give syphilis, although not having it, and to do that they have only to delicately pervert facts. We have read somewhere an article entitled "Transmission of Syphilis by Animal Lymph". It was a sensational title. Now, nothing can be more easily explained than this transmission. It is sufficient for the contamination of the lymph that the vaccinator, inoculating a series of children, transfers blood from a syphilitic subject by means of his instrument to the animal pustule into which he plunges it to get a fresh charge. But the poor animal is perfectly innocent of the mishap; the author of the mischief is the vaccinator, who has not taken the needful care with his instrument, and who has not cleansed it, as he ought, after each vaccination.

[To be concluded.]

INTRA-UTERINE VACCINATION AND SMALL-POX.—In a note on this subject in the JOURNAL for November 22nd (page 824), reference was made to the case of intra-uterine vaccination quoted by Dr. Bollinger as recorded by Dr. A. S. Underhill of Great Bridge. It may be of interest to state that this case was one in which a lady, revaccinated when eight months pregnant, the resulting pocks being very large and full of lymph, was delivered of a child which was vaccinated at the age of three months with fresh carefully taken lymph. No effect whatever was produced on the arm; and, about a month afterwards, the child was again vaccinated, particular care being taken to well rub in the lymph; but the result was again unsatisfactory. Dr. Underhill's explanation was that the vaccine virus, thoroughly pervading the system of the mother, had directly influenced the child, and made it not susceptible of taking again the virus. To this view Mr. Burton of Birmingham objected (JOURNAL, vol. i, 1875, p. 44), observing that, when he had charge of the Birmingham Small-pox Hospital, he practised vaccination in several instances upon women at various stages of pregnancy, and he had not a single case of unsuccessful vaccination amongst the children.

An analogous case of a somewhat striking kind, though relating to variola instead of vaccinia, is recorded by Dr. Cory in the last volume of *St. Thomas's Hospital Reports*. A woman gave birth, on December 14th, 1877, to a child at full time. On June 13th previously, and consequently when about four months pregnant, she had been admitted into the Small-pox Hospital at Hampstead, suffering from an unmodified attack of small-pox. On May 21st, 1878, she brought her child to the Surrey Chapel station for vaccination. She was then deeply pitted, and had lost most of her hair. The child showed no evidence of having had the disease. It was vaccinated, with the result of having five good characteristic vesicles, which went through a regular course, except that the areola on the eighth day was well developed.

LATENT ALBUMINURIA: ITS ETIOLOGY AND PATHOLOGY.

By GEORGE JOHNSON, M.D., F.R.S.,

Professor of Clinical Medicine; Senior Physician to King's College Hospital.

IT is a matter of common and in fact every-day observation, that the urine of persons apparently in perfect health is often found to contain more or less albumen. The discovery of what had been a latent and unsuspected albuminuria is often made by accident. Within the last twenty years, at least half a dozen members of our profession have come to me with essentially the same statement, which was to this effect: "I was testing some albuminous urine, when it occurred to me to compare the action of the tests upon a presumed healthy specimen. I therefore tested my own urine; and, to my dismay, I found it highly albuminous." During the last few months, I have been consulted by three men, each of whom, believing himself to be in good health, and having proposed an insurance on his life, had been found to have albuminuria by the medical adviser of the insurance-office. Quite recently, an apparently healthy young man told me that, seeing his medical brother testing some albuminous urine, he said to him jokingly, "You may as well test mine"; and the result was that it was found to contain an abundance of albumen. About the same time, a surgeon called on me and said that he had been made very anxious by the accidental discovery that one of his children had albuminuria. The eldest son, who was studying practical chemistry, had busied himself in testing the urine of every inmate of the house, when he discovered that a younger brother, apparently in good health and attending a day-school, had albuminuria.

These are a few instances out of a large number that have come under my observation in which the existence of albuminuria, heretofore latent and unsuspected, has been discovered as it were by accident.

The attention of the profession has recently been especially directed to the subject, of what I propose to call latent albuminuria by several physicians, especially by Dr. Moxon (*Guy's Hospital Reports*, 3rd series, vol. xxviii), by Dr. Clement Dukes (*BRITISH MEDICAL JOURNAL*, November 30th, 1878) and by Dr. Saundby (*BRITISH MEDICAL JOURNAL*, May 10th, 1879; also *Birmingham Medical Review*, July and October 1879). My main object in this communication is to maintain, first, that this latent albuminuria—albuminuria, that is, unassociated with any other evidence of functional disorder or structural disease—may, by a careful inquiry, be traced back, in a very large proportion of cases, to some probable exciting cause; secondly, that the presence of even the smallest trace of albumen in the urine is always pathological, never physiological, as I have somewhere seen it suggested that it may be; and that the neglect of this indication of a pathological condition and tendency, especially such negligence as involves repeated exposure to the exciting cause, may convert a temporary and occasional into a persistent albuminuria, which sooner or later, though it may be after many years, will result in a fatal disorganisation of the kidney.

In prosecuting the important practical inquiry into the probable cause of a recently discovered albuminuria, we shall not unfrequently come upon a history more or less clear and definite of an acute nephritis, either with or without dropsy, dating back for a period varying from a few months to several years. The acute renal disease may have been the result of exposure to cold and wet, or it may have occurred in connection with one or other of the following diseases: scarlet fever, measles, diphtheria, erysipelas, typhus or typhoid, pyæmia, rheumatic fever, etc.; or it may have been associated with the puerperal state, either before or after parturition. The acute symptoms having passed away, convalescence has appeared to be established; but, in some cases, the patient having been kept under careful observation, the albuminuria has been found to continue for months and years. In other cases, there has been a less careful and prolonged observation, and the history, therefore, is less complete. Convalescence having been apparently established, no further examination of the urine had been made until after an interval of months or years, when albuminuria is found to be present, and there is great reason to believe that it has continued from the time of the previous illness. The practical lessons to be deduced from facts such as these, which are matters of every-day observation, are: 1. To test the urine for albumen repeatedly during the progress of all febrile and inflammatory diseases until convalescence has been completely established; 2. When, in the circumstances referred to, albuminuria has been found to exist, to keep the patient under