this view advised her to go to the Royal Infirmary, whither he himself conducted her in a cab. On admission, she was in the following condition. She had an anxious pained look; her respirations were laboured, crowing, and much impeded. The saliva trickled from her mouth, as she could not swallow, and very often a spasmodic cough took place, ending in bringing up some mucus mixed with saliva and slightly tinged with blood. She had aphonia; attempts to whisper evidently gave pain, so she curtailed them to monosyllables, or substituted a sign, such as a shake or nod of the head. She hesitated about making any attempt at deglutition; she did try, however, and apparently took a spoonful of milk, but after four or five seconds it was expelled during a fit of coughing. This was repeated with a like result. On examining the throat, the orifice of the larynx was found to be very much narrowed—œdematous tissue surrounding it on all sides and extending backwards in the form of a tumour, which completely blocked the pharynx. The tip of the forefinger could occlude the portion of the laryngeal orifice that remained patent. The epiglottis was not active, and neither it nor the surrounding parts exhibited the usual sensitiveness to touch. In consequence of the tumour-like extension of the œdema behind and to the sides, the larynx was pushed well forward and fixed in that position. Her pulse was rapid and weak; her skin was hot and clammy. She was about three months pregnant. It was just such a case as would have been operated upon by tracheotomy under ordinary circumstances.

A fresh No. 12 catheter was passed into the trachea by way of exploration, and then the smallest sized tracheal tube that was at the time obtainable. This was found to be just on the thick side, but it passed. A fit of coughing, which lasted more or less for a couple of minutes, ensued; after which she had comparative ease. The tube was inserted at 6.20 P.M. on December 8th, and was retained till 8 A.M. on December 10th—it being removed at intervals of about twelve hours to be cleaned and to permit an œsophageal tube to be inserted for feeding purposes. In all, it was in the larynx thirty-five hours. At the end of that time, the œdema had greatly subsided; she then breathed easily. She could swallow and speak a little, and from this time she made an uninterrupted recovery.

The following points are particularly noted. The introduction of the tube was easy, as the laryngeal orifice was thrown well forward and fixed, owing to the great amount of œdema; and from the same cause the parts had an impaired sensibility, which rendered the manipulation tolerable. The first tube seemed to fill the entire free portion of the larynx above the cords. The next tube was very slightly narrower, at the same time thinner in its walls, thereby affording a large internal diameter relatively to the size of the tube. The ease which this gave was great, and, after it was introduced, the patient marked her appreciation of it by grasping and clapping the hands of the operator by way of thanks. She turned on her side afterwards, and soon fell asleep. The two last times the tube was inserted, it did not cause any cough or spasm.

In the former case, it was noticed that there was a distinct cough; in this case, the expiratory explosion of air was not what could come clearly under the physiological term cough, except toward the end of treatment, when there were sounds which were more purely explosive in character. Again: at first, after the introduction of the tube, the patient could not give the monosyllables as noted in the former case. It must be remembered, however, that she had complete aphonia on admission. Toward the end of the first twenty-four hours, she began to say "Yes" and "No"; and, after the thirtieth hour, the monosyllables were sounded distinctly.

The expectoration was profuse, semi-purulent, fluid, and streaked or mixed with blood; but it afterwards, towards the end of the first twenty-four hours, began to be thick and tenacious. The expectoration was thrown to the outside of the tube, sometimes by a distinct cough, when a quantity of the mucus had partly occluded the tube for a moment. On several occasions, when the mucus began to be tenacious, a little brush, mounted on a wire of definite length, was pushed into the tube and the expectoration removed. Twice, when the breathing was not quite so free, the tube was removed, and a very thin coating of mucus was found in the interior; this was washed off and the instrument returned.

Remembering the state of the pharynx, rendering deglutition impossible, and the loss of sensibility of the larynx prior to the introduction of the tubes, she was not asked to swallow during the first twenty-four hours while the tube was in position. Several times, a catheter was introduced into the œsophagus, and beef-tea, eggs, milk, and brandy were injected. At the end of the first day she began to swallow a little, and about the thirtieth hour she took several mouthfuls with the tube *in situ*. Several times she slept for hours with the tube in the trachea, and often she had snatches of sleep besides.

In this case, the œdema glottidis, mechanically blocking the pharynx, was so great as to exercise pressure on the superior laryngeal nerves,

exhibited in the non-sensibility of the epiglottis and the parts of the larynx above the true cords. This constituted a danger, as was shewn when she attempted to swallow. The impaired laryngeal sensibility, coupled with the fixity of the structure from the œdema, prevented the epiglottis and the neighbouring parts from closing the laryngeal orifice, when an attempt at deglutition was made. As a consequence of this and the pharyngeal obstruction, the fluid entered the larynx, passing down into the trachea before it produced the irritation necessary for its expulsion. A few seconds of quiescence ensued after the attempt to swallow the fluid before it was again brought up by coughing, and it is likely that the time might have been extended if persistent efforts to feed the patient were made. From cases which I have seen and heard of, this is not an unfrequent cause of at least hastening a fatal issue in children affected with œdema of the glottis. It is satisfactory to note that the patient, who was pregnant at the time when she was suffering from the throat-affection, afterwards had a well-developed strong child at full time.

[To be continued.]

ON A NEW METHOD OF ARRESTING GONORRHŒA.

By W. WATSON CHEYNE, M.B., F.R.C.S.,

Assistant-Surgeon to King's College Hospital.

HAVING been for some time past occupied with the problem of the infective diseases of wounds, the subject of gonorrhœa, as an affection probably belonging to the same class of diseases, has occupied my attention. The extreme contagiousness of this disease, the existence of a distinct period of incubation, and the steady spread of the inflammation from a given spot, all point strongly to a parasitic origin. Acting on this idea, I made, in the spring of 1879, a number of inoculations of gonorrhœal pus, under certain precautions, into flasks containing infusion of meat or infusion of cucumber. In these flasks micrococci grew in large numbers, and also sometimes bacteria, showing that these organisms were present in the gonorrhœal pus. Circumstances prevented me from pursuing this subject further at that time. In the meantime, Dr. Neisser published an elaborate research on this subject, in which he showed the presence of enormous numbers of micrococci in gonorrhœal pus, and in the pus from contagious ophthalmia. He further asserted that these organisms were always of a definite size, and that they differed in respect of size from the micrococci found in wounds. The presence of large numbers of micrococci in gonorrhœal pus has since been confirmed by several observers. Whether these micrococci are the cause of the gonorrhœal inflammation or not, I do not attempt to say, but the general history of the disease, taken together with these facts, points strongly to the idea that its essence consists in the growth of these or allied organisms.

If this disease be due to the spread of organisms, where are they situated? Several facts lead to the supposition that they are not only free in the urethral canal, but that they are also present in the substance of the inflamed mucous membrane. Thus, in the case of erysipelas, it has been demonstrated that the skin at the margin of the inflammatory redness is full of micrococci. Koch found, in his case of erysipelas in rabbits, that bacilli were present throughout the inflamed part, and coextensive with the inflammation. The same writer obtained a progressive gangrene of the tissues in mice by the injection of putrid blood, and he has demonstrated conclusively that this gangrene is due to an organism—streptococcus—which is present in large numbers around the limits of the gangrenous part. A similar observation was made by him in a case of spreading suppuration in rabbits. Mr. Lister has long held the opinion that, in the case of putrid sinuses, the organisms were present, not only in the canal of the sinus, but also in the substance of the unhealthy granulation-tissue lining them. This view has been justified by the fact that, though formerly, by the simple injection of the sinuses with antiseptics, he did not often succeed in eradicating the septic element, yet, since he has adopted the use of Volkmann's sharp spoon, and has removed the layer of granulation-tissue lining them, success is by no means uncommon. And, lastly, I have demonstrated that, though many forms of organisms will not survive if introduced into a healthy animal, yet, if an animal be previously in a state of ill health, these forms of organisms are not destroyed, but may be found alive in the blood or tissues.

In the case of gonorrhœa, then, I suppose that, at the time of infection, a small number of the specific organisms, which in all probability possess a considerable resisting power to the destroying action of the healthy living tissues, are retained in the urethra, that these go on developing, that the products of their growth irritate and weaken the

mucous membrane in their vicinity, that the organisms can then penetrate into and live in that weakened tissue, and that the extension of this process over a portion of the mucous membrane of the urethra is the cause of the inflammatory symptoms.

Now, granting that this view, which I think must be admitted to be very probable, were proved, the problem to be solved for the cure of gonorrhoea would be, how to destroy these organisms without at the same time injuring the inflamed and highly sensitive mucous membrane. If they were destroyed, one would expect the extension of the disease to cease, and the inflamed mucous membrane to return more or less rapidly to a normal state. On thinking this matter over, two substances appeared to me suitable for this purpose, being both powerfully antiseptic, and at the same time but little irritating. These are iodoform and oil of eucalyptus.

The next question was, how to apply them. It is quite clear that, if used as an injection, there would be no certainty that they would be brought into contact with the whole of the inflamed surface, partly because the swollen mucous membrane would interfere with the passage of the fluid, and partly because the patient would not in many cases apply it effectually. At the same time, an injection could not be expected to do much good, for it would flow out very quickly, and the antiseptic would not have sufficient time to act. I therefore use these antiseptics mixed with cacao butter, and made into bougies of various lengths. These bougies are introduced well into the urethra, and a strap and pad over and around the orifice retain them. The bougie rapidly melts, and the mucous membrane of the urethra remains bathed in the antiseptic material for any length of time desired. These bougies possess an additional advantage over injections in that from their size (they have a diameter of a No. 9 or 10 catheter, tapering at the point), they, so to speak, unfold the swollen mucous membrane, and thus cause the antiseptic to be more thoroughly applied.

I have tried the two antiseptics separately and also combined, and I find that they are most effectual when used in combination (possibly because iodoform is soluble to a considerable extent in oil of eucalyptus, and is thus brought into more perfect contact with the mucous membrane). The formula which seems best is five grains of iodoform* and ten minims of oil of eucalyptus in a bougie of forty grains. These bougies have been made for me by Mr. Martindale, of New Cavendish Street.

The specific cause of the disease being eradicated by this means, the question of further treatment arises. It seems to me that, although the development of the gonorrhoea is arrested, yet, if the discharge be allowed to become septic and irritating, urethritis might be kept up for some time. I, therefore, order an injection of boracic lotion (saturated aqueous solution of boracic acid), or an emulsion of eucalyptus oil (one ounce of eucalyptus oil, one ounce of gum acacia, water to forty or twenty ounces) to be used for two or three days. At the end of that time, injections of sulphate of zinc, two grains to the ounce, may be begun. At the same time, the great tendency of the urethral mucous membrane, when once inflamed, to remain in a state of inflammation, must be kept in mind, and everything which might tend to keep up the inflamed state must be removed. Notably, the patient must be cautioned against drinking, and it is well to order diluents and alkalies.

The method may be summed up as follows. The patient is first told to empty his bladder, partly to clear out his urethra, and partly to prevent the necessity of expelling the antiseptic from the canal for several hours. He then lies down on his back, and a bougie from four to six inches long is introduced, and the orifice of the urethra closed by strapping. The bougie ought to be dipped in eucalyptus oil, or in carbolic oil (1—20) before insertion. The patient is instructed to refrain from passing water, if possible, for the next four or five hours. If the case be severe and advanced, he takes another bougie home, and is instructed to introduce it in the same manner after he next passes urine. On that evening, or on the following day, he commences the antiseptic injection, which he uses four or five times daily. On the third or fourth day, when the symptoms have entirely subsided, an injection of sulphate of zinc, two grains to the ounce, is begun.† At the same time, the other points mentioned are attended to.

I have now used this method in about forty cases, and in all the result has been the arrest of the progress of the gonorrhoea. For a day or two the purulent discharge continues; but afterwards it steadily diminishes in amount, becoming in four or five days mucous, and ceasing altogether in a week or ten days. At the same time, the

scalding and pain and the symptoms of inflammation rapidly diminish, and disappear completely in about thirty-six to forty-eight hours. In fact, the case becomes no longer one of virulent gonorrhoea, but one of simple urethritis, rapidly progressing towards recovery, if properly treated.*

I have used this treatment only in the early stages of the disease, from the first to the seventh day after the commencement of the symptoms; but it has answered equally well in all. Thus the following is the case in which it was used seven days after the commencement of the symptoms. The patient presented himself on June 19th, stating that the symptoms of gonorrhoea had existed for seven days. There was a profuse purulent discharge from the urethra; the penis was somewhat swollen and red; there was intense scalding when urine was passed, and a constant feeling of heat and uneasiness; no chordee. A bougie, containing ten grains of iodoform and ten minims of eucalyptus oil, was passed down, and the orifice closed in the usual manner. The patient was also ordered an injection of an ounce of oil of eucalyptus and an ounce of gum acacia in a pint of water, to be commenced in the evening, and to be used four or five times daily. On the 19th, he again presented himself, and stated that he had not passed water till five hours after the introduction of the bougie; that the scalding and feeling of uneasiness rapidly subsided, and had completely ceased in forty-eight hours; and that the discharge had steadily decreased from the second day, and was now very small in quantity. He was ordered the sulphate of zinc injection, which completed the cure in three days.

In one case, there was a recurrence of the symptoms. The patient, a hospital patient, first presented himself on June 5th, stating that on June 2nd, five days after connection, a discharge had commenced, which had steadily increased, and was now profuse and accompanied with considerable uneasiness and scalding in passing urine. A bougie containing ten minims of oil of eucalyptus alone was inserted; no other treatment was ordered. On June 9th, he returned, stating that, after the introduction of the bougie, the scalding and uneasiness had diminished, and had almost disappeared on the evening of the 6th; but that on the afternoon of the 7th they began to return, and were now more severe than on the 5th. I introduced a bougie containing ten grains of iodoform and ten minims of eucalyptus oil, and gave the patient another to insert at bedtime. At the same time, I ordered the injection of boracic lotion to be commenced on the following day. When seen again on the 16th, he stated that this time the treatment had been successful, and that now the discharge was very slight. An injection of sulphate of zinc and a mixture containing copaiba were ordered, and the discharge ceased entirely on the 20th.

In two or three cases, there has been slight increase in the scalding on the first or second occasion on which the patient passed urine after the introduction of the bougies; but this has only been temporary, and these cases were as rapid as the others. In four instances, however, there has been considerable increase in the symptoms for twenty-four or thirty-six hours. In three of these, the bougies had been made with bees' wax; and they did not melt properly, and consequently came out of the urethra at various periods as small cakes. Further, it seems that some iodine had been set free from the iodoform, probably during their manufacture. In the fourth case, four bougies, each containing 10 grains of iodoform were introduced in succession. In all these, however, the symptoms passed off in about three days; and then the gonorrhoea was found to be checked, just as in the other instances.

Such are the results as yet obtained by this method. I do not claim any specific power for the two substances I have mentioned. It may be that there are other antiseptics which would be more suitable, and I intend to test any which seem likely to yield good results. Whatever substance be used, however, I venture to think that the results already obtained show that the principle on which it ought to be applied, and on which it will prove most satisfactory, is that which I have attempted to indicate in this paper.

My thanks are due to my colleague, Mr. Royes Bell, for having allowed me to use some of his cases; and to Mr. Farmer, the assistant house-surgeon, for carrying out the treatment in these instances.

* The course described here is that usually followed when boracic lotion has been employed as the injection; but since I have begun the use of the eucalyptus emulsion, the cessation of the discharge has, as a rule, been more rapid. Thus, to give an example, a patient came to the hospital on July 3rd with symptoms of gonorrhoea, which had lasted four days. He was suffering from a very acute attack, having severe scalding and commencing chordee. He had not previously suffered from gonorrhoea. A bougie containing five grains of iodoform and ten minims of eucalyptus oil was introduced; and he was ordered to begin an injection of the eucalyptus emulsion (1 in 40) in the evening. The patient showed himself again on July 7th, and stated that in twenty-four hours the painful symptoms had entirely disappeared, and that the discharge diminished rapidly, and ceased altogether on July 6th. I have since that time had several nearly as rapid cases. I have tried in three cases injections of eucalyptus emulsion without previous introduction of a bougie, but without any appreciable effect on the progress of the disease.

* A considerable number of the cases have been treated with bougies containing ten grains of iodoform; but Mr. Martindale informs me that during the warm weather it is almost impossible to make them. I find, however, that bougies containing five grains are quite satisfactory, and I have had no symptoms of irritation following their use.

† In hospital practice, where the patient is only seen once a week, and where there is no great necessity for arresting the discharge quickly, I do not order the sulphate of zinc injection till the week following the introduction of the bougies.