asserted that ovarian cysts have been cured by these means; but, without attaching too much importance to this statement, it is at least possible that ovarian tumours may thus be prevented in the earliest stage. Locally, douches and injections are freely used, and with the best results. In overcoming sterility, the effects are so remarkable as to have given to one of the springs the name of *Source des Garçons*.

For scrofulous affections of the skin, the arsenical waters of La Bourboule are in high favour. Eczema, pityriasis, psoriasis, all yield to their curative influences, particularly the scrofulous and rheumatic forms of these diseases.

As in the case of the Mont Dore and other similar watering-places, some of the beneficial results experienced by bathers are no doubt attributable to the change of air and scene, to the beautiful scenery and the pleasant outdoor life; but, after due allowance has been made for all of these concomitant advantages, I think I have said enough to show that the course of treatment at Saint Nectaire and La Bourboule is specially valuable in rheumatic affections, and in the treatment of scrofula in its various manifestations in the lymphatics, the bones and joints, the sense-organs, the skin, and the uterus and its annexes.

NOTES ON THE MINERAL WATERS AND CLIMATE OF SPA.*

By LITTON FORBES, M.D., Resident Physician.

SINCE the middle of the sixteenth century, the chalybeate springs of Spa have enjoyed the reputation of being therapeutically some of the most valuable in Europe. In the eighteenth century, they were more frequented than perhaps any others, and formed a centre round which a large share of the wealth and fashion of the Continent used to meet. Of later years, this has, to a certain extent, ceased to be the case. The abolition of gambling has considerably diminished the number, while it has improved the quality, of the visitors who annually resort to Spa. Those who come now, come, as a rule, mainly in search of health. They live temperately and moderately, and rarely fail to derive considerable and permanent benefit from their sojourn at these springs. No mineral waters, perhaps, in suitable and properly selected cases, are capable of effecting so much good in such a short space of time as those of Spa. But to do this the cases must, in the first place, be such as require a tonic course of treatment; and, in the second, that treatment must be carried out, both on the part of the patient and the physician, with judgment and discrimination. The physiological action of the various chalybeate springs of Spa is by no means identical. No greater error, indeed, can be made, than that of treating in a routine way, and without attention to individual peculiarities, even those cases in which Spa waters are pre-eminently suitable. Each case must be treated on its own merits; and it is in the choice of the particular spring, and in the mode in which its waters are prescribed, that the special skill and local knowledge of the practitioners is chiefly shown.

Comparing Spa with two well known, and at present fashionable, watering-places—Schwalbach and St. Moritz—we are at once struck by the much larger amount of iron which is held in solution in the springs of Spa. Thus the two Pouhon springs of Spa contain respectively three times more iron than the Stahlbrunnen and Weinbrunnen of Schwalbach, and from six to eight times as much as the Paracelsus spring of St. Moritz. But, while thus rich in the actual amount of chalybeate matter which they contain, they are still more rich in the quantity they contain proportionately to other ingredients, such as the earthy and alkaline salts. Thus, taking the series of the alkaline bicarbonates, we find that, representing the amount contained in the two Pouhon springs just mentioned by the numbers 1.8 and 5.9, the amount contained in the Weinbrunnen of Schwalbach would be represented by 12.1, and that in the Paracelsus of St. Moritz by 16.9. On the other hand, the amount of free carbonic acid is slightly in favour of Schwalbach, and St. Moritz as compared with Spa, viz., for the first, 27.1; for the second, 33.6; and for the Pouhon of Spa, 25.5. It is this richness in iron, both actually and relatively, which constitutes at once the chief virtue and the chief drawback in the Spa waters. It is this which, in suitable cases, makes their action rapid and beneficial, and which in unsuitable cases rapidly aggravates the patient's symptoms. Consequently, there are few waters which require more care in their administration than those of Spa, and few also in which a satisfactory result can be so speedily and confidently promised when once they have been found to be in harmony with the patient's requirements and temperament.

It would be impossible to give, even briefly, a sketch of the various

morbid conditions for which the waters of Spa have from time to time been vaunted as almost specifics. Such indiscriminate praise is entirely at variance with the severe and scientific methods of modern medicine; and, moreover, utterly defeats its own object. It causes annoyance to the physician, intense disappointment to the patient, and brings unmerited censure on a really valuable therapeutic agent. I will, therefore, state at the outset what I venture to consider as distinct contraindications to the use of the Spa waters. Every year, patients come to these springs, frequently without having obtained any medical advice whatever as regards their choice of a spa, stay here some days, and depart considerably worse than when they arrived. As a rule, such patients belong to one of two classes; they are either plethoric subjects, with a tendency to attacks of acute gout, and a threatening of cerebral or other congestions; or they are phthisical patients, with abundant hæmoptysis and a well marked hectic condition. In the first class of invalids, the state of chronic congestion in which many of their internal organs are increased; in the second, the stomach is quite unable to absorb or assimilate the amount of iron in the waters, the heart's action is deranged and accelerated, and a fresh burst of hæmorrhage is often the result. Invalids belonging to either of these two categories should not visit Spa for the purpose of drinking its waters, although they may, with perfect safety and even benefit, resort to the mineral baths, or enjoy the bracing mountain-air of the high table-land of the Ardennes, on which Spa is situated. There are, however, in addition to these two classes, a large number of invalids who come hither, to whom, on their first arrival, the waters cannot be at once ordered with advantage. Among such, I may mention women at the epoch of the menopause, who suffer from more or less abdominal congestion, with palpitation of the heart or frequent headaches. In these cases, a week or so of preliminary exercise and careful dieting, together with the use of baths, will frequently be sufficient to enable them to assimilate the waters, from which they will eventually derive much benefit. There is also another set of cases which require very careful treatment. They are met with chiefly, though not entirely, in girls or young married women in whom the nervous system is in extreme disorder, and in a state of unstable equilibrium. Such persons are the victims of intense irritability, in which every organ of the body, the stomach included, shares. These patients frequently find it impossible, during the early days of their sojourn at Spa, to assimilate the waters; and it is worse than useless in such cases to press them. Tepid mineral baths, with or without graduated douches, often prove very beneficial. They calm the peculiar condition of nervous irritability present, and thereby prove quite as efficacious as the chalybeate waters, which, moreover, can generally be had recourse to with still further advantage at a later period.

On the other hand, the mineral waters of Spa are pre-eminently indicated in that large group of diseases which may be classed under the heading of *anæmia*. There may be *anæmia* from nervous exhaustion, *anæmia* as a direct result of the physiological insufficiency of any organ or group of organs, and *anæmia* from excessive or deficient secretion or excretion, either locally or generally. This *anæmic* or *spanæmic* type of disease is one which has assumed increased importance in recent times, and is more or less intimately connected with the growth and development of modern civilisation, especially in the great centres of population and industry. *Anæmia* is a symptom daily becoming more common, at least among the patients who resort to these springs. Tonic treatment is now undoubtedly often required where, one or two generations ago, depletion would have been as imperatively called for. It is in these cases of what I will venture to call "*vital weakness*," with or without localised or functional lesions, that the Spa waters are eminently beneficial. It matters not, apparently, whether the *spanæmia* present be due to a disordered condition of the chylipoietic viscera or of the uterus, of the lungs or of the nervous system, or whether it be dependent on some subtle change in the constitution of the vital fluids themselves, apart from any appreciable local manifestations. Whatever the cause be, the result of the use of these waters in this great subclass of disease is apparently always the same, namely, temporary or permanent improvement.

Among the various forms of *anæmia*, there is one to which I would wish to draw especial attention; I mean climatic *anæmia*. Perhaps no class of cases derive so much benefit, and in so short a time, from the ferruginous waters of Spa as this. I have lately had under my care an English lady, whose health had been completely broken by a prolonged residence in the moist and relaxing climate of Brazil. After taking the waters for one week, a marked improvement was visible. In this case, they agreed better than they sometimes do; nevertheless, in the form of *anæmia* referred to, their action would appear almost that of a specific. I have seen the greatest benefit from their use among persons whose health had been undermined by long residence in tropical and

* Read in the Section of Medicine at the Annual Meeting of the British Medical Association in Cambridge, August 1880.

unhealthy climates, such as parts of India, South America, and China. In women, this climatic anæmia is frequently associated with disordered uterine function, and it is just in such cases that the waters of Spa are particularly applicable.

As regards the climate of Spa, it is essentially a mountain climate. The town itself is situated at an altitude of about 1,000 feet above the sea, while the neighbouring hills range in height from 500 to 800 feet more. The season commences in May, and terminates in October. The best time for persons with delicate chests to arrive at Spa is between June and September. Before and after that date, there is a chilliness in the morning and evening air which is not without danger. At times, the rainfall is considerable, even in summer, and is generally accompanied with wind and more or less piercing cold. During the continuance of such weather, invalids should not venture out to any of the more distant springs, but either cease the use of the waters altogether for a day or two, or confine themselves to those in the town. The general salubrity of Spa is now beyond question. The town is singularly free from all diseases of a zymotic type. Whatever its sanitary condition may have been in the past, it is now at least beyond reproach. The natural position of the town on a hill facilitates drainage, while the civic authorities have adopted, so far as appeared called for, all the most modern hygienic improvements. Typhoid fever has been for several years unknown; while, during the cholera epidemic of 1866, when the neighbouring towns of Verviers and Liège suffered severely, not even a sporadic case occurred in Spa.

CLINICAL MEMORANDA.

TREMORS IN A CHILD RESEMBLING PARALYSIS AGITANS.

THE following case may possibly prove of interest to some of your readers.

Mrs. C. consulted me in the early part of this year about her son, aged eight years, who, she informed me, was troubled with "shivers", after which he became hot and flushed, and perspired; the whole attack did not last longer than half an hour, but returned almost daily, and the child seemed perfectly well in the intervals. I could detect nothing wrong with the child, except that he had an occasional nervous twitching of the mouth and eyelids when spoken to, which he was supposed to have copied from a sister, two years older, who had recently been under my care for similar twitchings of the face and slight chorea. I was also informed that the lad had complained of pains about his body like rheumatism.

The "shivers" lasted so short a time that I had no opportunities of observing the lad in one of his attacks until some three months had elapsed, during which time they returned very regularly, except for a period of four weeks which he spent away from home at the sea-side. On his return the old symptoms reappeared, and soon became so severe that the parents were enabled to call me in time to see for myself. I found that the so-called "shiver" was a tremor of the muscles of the arms and legs, exactly resembling the movements of paralysis agitans, attended with slight subsultus tendinum. The lad was perfectly conscious, and kept his fingers clasped together to try and restrain the to-and-fro movements of his arms that occurred when his hands were loosed. When asked to carry either hand to his mouth, or to pick up any small article, he did so without any of the jerking, irregular movements observed in chorea, though the tremor still continued. His legs were kept a little apart, apparently to keep his knees from knocking together. There was no perceptible loss of muscular power in his grasp. The lad was flushed and perspiring, possibly from the muscular exertion. The tremors continued for four days almost without intermission, except when sleep was induced by chloral hydrate. There was no perceptibly disordered state of the primæ viæ, and anthelmintics brought to light no worms as a cause of reflex irritation. I ordered a bland nutritious diet, interdicting tea and coffee, prescribed the saccharated carbonate of iron, five grains to be taken twice daily after food, and five grains of chloral hydrate at bedtime, to induce sleep. I also recommended the use of the constant current, by means of a Pulvermacher's chain of 120 links for an hour daily. After the expiration of four days, the child went without his daily tremor. Since then he has had no return, though the slight twitch of the eyelid is still noticeable at times. I am still continuing the treatment, for fear of a relapse, with the exception of the chloral at night, though the child has been free from any sign of a tremor for some weeks. I might add that the only noticeable point in the family history is gout, with which the father is much troubled.

HENRY Y. PITTS, L.R.C.P. Lond.,
Bank Villa, Tue Brook, Liverpool.

REPORTS

OF

MEDICAL AND SURGICAL PRACTICE IN THE HOSPITALS AND ASYLUMS OF GREAT BRITAIN AND IRELAND.

HOSPITAL FOR CONSUMPTION AND DISEASES OF THE CHEST, BROMPTON.

TWO CASES OF THORACIC ANEURISM.

(Under the care of Dr. THEODORE WILLIAMS.)

Reported by F. P. WIGHTWICK, M.R.C.S., Clinical Assistant.

ALFRED J., aged 40, married, an engine-fitter, was admitted into Wallace Ward on July 20th, 1880. He had had good health, with the exception of measles and small-pox, until sixteen months previous to admission. From that date his health had gradually been failing, and he found it more and more difficult to continue his usual occupation, which necessitated considerable physical exertion. He could not attribute his illness to any distinct cause. His chief complaint had been shortness of breath on exertion. On examination, there was moderate dulness in the front of the thorax, limited above by the first interspace, and continuous below with the cardiac dulness, extending to the left as far as a line drawn vertically from the centre of the clavicle, and bounded to the right by a similar vertical line, distant a quarter of an inch beyond the middle of the sternum. No pulsation or prominence was visible over this area, but a marked heaving impulse could be distinguished at the thin end of the stethoscope when the broad end was applied to the chest. Between the second and third ribs, near to the sternum, the respiratory murmur was of a tubular character. Posteriorly, over a space lying between the left scapula and vertebral column, and corresponding to the fourth, fifth, and sixth vertebræ, the respiration was also tubular. No *bruit* was audible. He complained of severe pain passing from the ensiform cartilage to the angle of the left scapula, of a heavy burning nature; and had slight cough, without expectoration. There was some hoarseness and loss of the musical quality of the voice. There was no dysphagia. The pupils were both greatly but equally contracted. No difference could be distinguished between the character of the two pulses, the number being 72; temperature, 98.8° Fahr. His urine was normal. He was treated by means of rest, and with ten-grain doses of iodide of potassium three times a day. He continued much in the same condition till August 11th, when he desired to go home, feeling, he said, "quite well".

August 19th, 7.30 A.M. While getting out of bed, he suddenly fell back, and immediately brought up about two and a half pints of blood. This he distinctly vomited, and in a manner so different from that usual in fatal hæmoptysis, that it led to the suspicion, on the part of Mr. Wightwick, that the aneurism had opened into the œsophagus. There were marked symptoms of collapse. 11.30 A.M. He had much rallied, but shortly after he vomited nearly another pint of blood, consisting mainly of large clots, and smelling strongly of the gastric juice. He lingered for three days, complaining only of nausea and thirst, and taking considerable quantities of beef-tea and milk, without the slightest dysphagia.

August 22nd, 2 P.M. A small quantity of bright blood welled up in his mouth; he seemed to try to vomit without having the strength to do so, and rapidly sank.

On *post mortem* examination, the stomach and upper part of the intestine contained blood; in the stomach the blood had formed a large clot, continued upwards into the œsophagus. Both lungs were fully inflated and anæmic. A good deal of œdema was found in the posterior portions. Nothing abnormal was noticed in the air-tubes, with the exception of a small patch of greenish discoloration, apparently due to incipient sloughing in the mucous surface of the left bronchus, slightly below the bifurcation. The walls of the heart were thin in proportion to the cavities, which were rather enlarged. The valves were normal. The first portion of the aorta was uniformly dilated, and showed numerous raised patches of soft atheroma. The pulmonary artery was dilated, but apparently not diseased. At the junction of the third portion of the arch with the descending aorta, a large orifice led the way into a good sized oblong aneurism, which lay parallel to, and to the right of, the thoracic aorta. The length of the aneurism was between four and five inches; its diameter between two and a half and three inches. Six of the dorsal vertebræ were deeply eroded, and their cancellous tissue lay bare at the bottom of the sac; the remaining sides of the sac possessed true arterial walls. No coagulum was found in the