same properties now as then.....Pancreatine is a very wonderful thing, and not half understood or appreciated at present."—(Signed) Julius Schweitzer.

HEPATIC ABSCESS OPENED ANTISEPTICALLY: DEATH.

By NEIL MACLEOD, M.B. Edin., Shanghai.

W. M., a merchant's clerk aged 39, complained, on November 1st, 1879, of a severe stitch-like pain in the right infra-axillary region, and of inability to lie on either side. He had resided in Shanghai without a change for fourteen years, had been a free-liver, had taken little or no exercise for some years, and had a specific history dating ten years ago. In July 1879, he had a severe attack of diarrhoea; in the end of August, an acute attack of dysentery; and in the middle of September he went to Japan, the stools being solid, and one daily. He continued well until October 22nd, when the dysentery returned slightly; and, on his return to Shanghai in the end of October, the motions were liquid, two or three a day, light yellow, foetid, and occasionally contained a little blood and mucus.

From the 1st till the 13th of November, in the morning, the temperature varied from 97.8 to 100.3 deg.; in the evening, from 101.8 to 103.8 deg., once reaching 104.6 deg. The pulse varied from 80 in the morning to 104 in the evening, and latterly he had had night-sweats. There was no rigor or shivering at any time. The motions were sometimes solid, sometimes fluid—one to three daily; and there was occasional vomiting. The hepatic dulness was increased, but not markedly so. On November 11th, he was removed from the country house where he was residing into the settlement, when abscess of the liver was suspected, from the increased liver-dulness, fever, sweats, tenderness and pain on the right side, and slight bulging just behind the anterior axillary line. Fifteen to twenty grains of quinine, given frequently, had no permanent effect on the temperature; and opium and chloral were given to procure sleep. Mustard, a blister, and heat in various forms, were applied to the side.

November 13th. I attempted a thorough systematic examination (but had to desist, as the patient seemed unable to bear it), with this result. The tongue was dry, with a white fur at the back part; no appetite, and nausea. Slight bulging was to be seen over the lower ribs in the mid-axillary line and behind it; and in this region there were flattening of the intercostal spaces, slight oedema, and a tender spot. In the nipple-line, comparative dulness stretched six inches from the fifth rib to half an inch below the costal margin; absolute dulness four and a half inches from the sixth rib in the same line. On tracing the line of dulness backwards, nothing abnormal could be made out in its direction. Breath-sounds were inaudible over the bulging area, where friction could be distinctly heard on deep respiration, which caused a stitch-like pain and a sudden check in the respiratory act. Pulse 90, feeble; face flushed slightly; apex-beat not displaced. Morning temperature 99 deg.; evening pulse 104; temperature 102.8 deg.

November 14th. Pulse 101; temperature 98.3 deg. At 9 A.M., with Dr. Little's assistance, I performed the following operation under chloroform, with antiseptic precautions. I passed the largest-sized Matthieu's aspirator-needle between the seventh and eighth ribs in the medio-axillary line, at the tender spot before mentioned. At a depth of little more than an inch, yellow pus flowed through the tube. Aspiration was immediately stopped; and, the cannula being left *in situ* as a guide, I made an incision on each side of it, and cut down until pus began to well up by the side of the cannula. Having removed the latter and enlarged the opening with a probe-pointed bistoury, fully a pint of thick yellow odourless pus flowed from it, with the aid of pressure in the epigastrium. A drainage-tube four inches and a half long, with a calibre of half an inch, and stout silk guards, was introduced; and the usual dressings applied, and secured by gauze and elastic web bandages. No attempt was made to prevent the entrance of air into the cavity, which allowed a probe to pass five inches in a direction right across the body, at right angles to the surface. At noon, the pulse was 80; at 4 P.M., 78, and compressible; temperature 97.2 deg. At 9 P.M., pulse 90; temperature 98 deg. The patient said he was very comfortable. The discharge, under the microscope, showed pus-corpuscles full of granules, much free granular matter, and many large cells about two or three times the diameter of a pus-cell, free and in masses, and also very granular.

November 15th. Pulse 90; temperature 97.8 deg. He passed an uncomfortable night, serum having soaked through in large quantity, at the edge of the dressing behind, and the bandages having been too tight. He had no pain. There was about an ounce, of thick shreddy pus in the dressing. The expression was no longer anxious. He had

one solid motion. Evening: pulse 108; temperature 100.8 deg. He passed a very comfortable day, and the appetite was very good.

November 16th. Pulse 98; temperature 99.4 deg. He slept well. The dressing contained about half an ounce of pus in shreds. Round the wound there was a little bulging and tenderness. Temperature at noon, 97.8 deg. Evening: pulse 108; temperature 101.4 deg. He had one motion.

November 17th. Pulse 96; temperature 98.4 deg. The discharge was bile-stained. There was quite a deep hollow in the right hypochondrium. Evening: pulse 114; temperature 101.8 deg. He had two motions.

November 18th. Pulse 98; temperature 97.8 deg.; discharge less, but enough to necessitate daily dressing. Bulging and tenderness were still present, chiefly behind the wound over the seventh rib. Evening: pulse 108; temperature 102.1 deg. He had two motions. For three days, the patient had taken ten grains of iodide of potassium thrice daily, peritonitis of the rib being suspected.

November 19th. Pulse 108; temperature 99.7 deg. I gave the patient chloroform, and slit up the bulging side of the wound backwards. Pus had burrowed backwards under the skin; and the rib could be felt thickened, but not bare. Evening temperature 103 deg.

November 20th. Pulse 108; temperature 97.6 deg. The wound looked healthy and filled with clot. The discharge was as yesterday, and still bile-stained, very thick. Evening: pulse 114; temperature 103 deg. Twenty grains of quinine were given in the afternoon.

November 21st. Temperature 98 deg. I suspected tension from accumulation of matter. By means of a glass funnel, an India-rubber tube attached to it, and a long glass nozzle, having a calibre of a quarter of an inch, the whole being filled with a saturated solution of boracic acid (antiseptic and unirritating), I drew out of the abscess an ounce and a half of thick curdy pus. The air was first expelled from the tube by keeping the funnel a little above the abscess-level; then, on introducing the tube into the abscess, and lowering the funnel beneath the level of the abscess, the apparatus acted as a syphon. Evening: pulse 114; temperature 102.4 deg.

From this date to December 1st, the pulse continued to rise gradually. The temperature in the morning varied from 97.7 to 99 deg.; in the evening, from 99.4 to 101.7 deg., being once 102.4 deg. The motions became loose and frequent, but were kept under control by means of small doses of Dover's powder and bismuth subnitrate. The dressing was changed daily; and the discharge soon ceased to be bile-stained, remained odourless, and varied from a quarter to half an ounce in twenty-four hours, latterly becoming of a livery colour. It contained aseptically. On the 28th, the probe passed three inches obliquely backwards and downwards. Twenty grains of quinine were given frequently, without appreciable effect on the evening temperature. The patient sat up all day, sometimes walked in the verandah, and did not seem to lose strength.

From the 1st to the 20th of December, the temperature in the evening varied from 99 to 102.5 deg., being usually about 101 deg.; in the morning, as a whole, below normal. Two severe rigors, with an interval of a week between, were each followed by a change from the liver-coloured discharge to a fresh yellow-pus, lasting for two or three days, and then becoming once more dark-coloured, remaining aseptic throughout. Strength began to fail, and the appetite was lost; and, finally, severe diarrhoea set in. All action ceased in the wound, and the walls of the abscess could be felt soft and rotten by means of the probe. Death occurred on the 20th. I may add that there were never any lung-symptoms present.

POST MORTEM EXAMINATION, twenty hours after death.—There was no smell in the wound. The interior of the body was warm. The organs were anæmic. The liver was very fatty, the left lobe being about twice its natural size. The back part of the right lobe, opposite and behind the wound, was occupied by a cavity nearly twice the size of a hen's egg, having what appeared to be prolongations backwards and inwards; one of these, nearly two inches long, admitting the finger; and all containing pus of a lighter colour than the cavity. The liver-tissue round the abscess was soft and extremely friable, and bands of fibrous tissue ran through the cavity. There was no abscess outside of this softened area. Firm adhesions were present opposite the abscess, over a space of about six square inches from the wound backwards. The peritoneal cavity was normal. From the friability of the liver-tissue in the neighbourhood of the abscess, and the firm adhesions on removal of the organ (which I took the precaution to slice from left to right *in situ*, removing each slice), it was impossible to determine whether it was a case of multiple or of single burrowing abscess. From the clinical history and the result, I am inclined to think it was a multiple abscess of the liver.

REMARKS.—This is the first and only case treated by the method

above described which has been unsuccessful. I have had two cases in my own immediate charge; this one, and another which was published in the *Lancet* of June 29th, 1878. I have seen in consultation, and assisted in operating on, four cases under the care of two of my colleagues in Shanghai, all of which recovered; and Dr. Johnston of Shanghai treated a case in a Chinaman in the summer of 1879, with the same result. In all seven cases thus treated, one has died. The number of cases is as yet small; but I think that the results so far are encouraging, and better than those obtained by any other method of treatment yet published, being 85.71 per cent. cured. The other five cases that have been treated in Shanghai are, I am informed, to be published shortly.

CHAULMUGRA OIL IN PHTHISIS.

By WILLIAM MURRELL, M.D., F.R.C.P.,

Senior Assistant-Physician to the Royal Hospital for Diseases of the Chest; Lecturer on Practical Physiology at the Westminster Hospital.

DURING the last two and a half years, I have used chaulmugra oil in fifty-nine cases of phthisis—forty-two being men, fourteen women, and three children. Of the adults, all, with the exception of five, were between the ages of twenty and fifty. In sixteen, there was consolidation; in twenty-eight, there was softening; and in fifteen, there were signs of cavities.

My first endeavour was to find out what dose could be given with safety. In twenty-two cases, the chaulmugra was given in cod-liver oil; and in thirty-one cases in milk. The latter was decidedly the better vehicle. The plan usually adopted was to begin with three drops, three or four times a day; and then gradually to increase the dose, week by week, till the patient could take no more. In about half the cases, ten minims four times a day was the limit; and this frequently gave rise to vomiting and purging. Sometimes the diarrhoea would come on after only a few ten-minim doses had been taken, but more commonly not till the expiration of a week. Often enough the attack was a sharp one—the patient having five or six watery motions during the day, and as many at night. The stools were in most cases described as being liquid, and were unattended with griping or straining. There was usually a temporary disrelish for food, the patient feeling altogether out-of-sorts. These symptoms quickly subsided on discontinuing the medicine. In a few cases, they were produced by smaller doses—even two or three drops upsetting the stomach. On the other hand, some patients were able to take considerably larger quantities. Thus a man, aged 26, took chaulmugra oil four times a day, for six weeks, in doses of from six to twenty-four minims, without experiencing the slightest difficulty. He then tried half-drachm doses, but this made him violently sick, and he had to discontinue it. Most patients found that they could take it better after meals than on an empty stomach.

An attempt was made to estimate the value of chaulmugra oil by giving, alternately, for periods of six weeks, cod-liver oil, and the same dose of cod-liver oil with the addition of the chaulmugra. This plan presented so many difficulties that it had to be abandoned; and the conclusions here given are derived almost exclusively from those cases in which the oil was given in milk.

In twenty-four of the thirty-three cases, I was satisfied that benefit was derived from the treatment, although in some instances the results were not very striking. The chaulmugra seemed to act first as an expectorant, and “the phlegm came up more easily”; then “the cough was less troublesome”; and, finally, the patient “felt better in himself”. I was unable to discover any improvement in the physical signs, even after the oil had been taken for many weeks. Sometimes the loss of weight appeared to be temporarily arrested, but there was rarely any absolute gain. Several of the patients took the chaulmugra with benefit after they had ceased to improve on cod-liver oil.

I have had very little experience of the chaulmugra oil *perles*, as they are too expensive for hospital use. In one case, however, the results were very satisfactory. In twenty-three cases of phthisis, I used chaulmugra oil as an external application—the patient rubbing into the chest from two to four ounces weekly. This is an excellent mode of administration, and yields the best results. I have never known it to upset the stomach, and it certainly eases the cough and loosens the phlegm. In some instances, there has been a marked gain in weight. As a rule, I have employed it alone, experimentally; but ordinarily it would be used simply as an adjunct to other treatment. The smell is not very pleasant, but can be readily covered by using a sweet-scented violet-powder. In some cases of marasmus, I have employed inunctions of chaulmugra oil with marked benefit.

SURGICAL MEMORANDA.

EXTENSIVE CARBUNCLE.

IN reference to the remarkable case of carbuncle reported by Mr. Walter, in the *JOURNAL* of October 23rd, I would like to call attention to the extraordinary extent to which the disease spread, in special relation to the fact that such a continuous extension almost proves that a carbuncle is something more than it is generally reputed to be, viz., a sloughing inflammation, or destruction of cellular tissue, induced by a faulty condition of the blood or general-health. Such a course and progress, as is described, goes very far to confirm the opinion formerly expressed by Mr. Startin; and since several times repeated by me, in the medical journals (*BRITISH MEDICAL JOURNAL*, 1876; *Lancet*, 1874 and 1869), that carbuncle is essentially not a mere destructive inflammation, but the growth and life of a distinct superadded skin-disease. Whether in it there be a parasitic vegetable cause, or a special growth of animal cells, the whole appearance, course, and almost definite duration of the disease, point strongly to its specificity; whilst microscopical examination has certainly, now and again, shown the presence of apparently cryptogamic germs, in addition to the hyperplastic cell elements of the growing disease.

Further: in view of the fatal issue of the case, I venture to recall attention to the marvellous efficacy of carbolic acid in staying the spread and progress of a carbuncle, when applied by the “insertion” method.

This method, as detailed in the above papers, has now been largely and successfully employed by myself and many others; and consists in inserting into the diseased spot a small quantity of a concentrated solution of the acid, so as to destroy, especially, the centre—the presumed root or stem of the disease—from which it spreads circularly. In the earliest stages of both boil and carbuncle, almost any caustic will, as is well known, commonly abort the disease. After the first periods, there is nothing like the carbolic acid: for it can be used concentrated, and yet freely, as it has the valuable property of only mildly attacking healthy organised tissue; whilst it destroys the life and vitality of free cells and germs. The carbolic acid is of little avail when merely applied externally, but it at once arrests the growth when inserted into the disease, so as to come into direct contact with the morbid germs or cell-elements; and its effect, when visible, upon the colour and aspect of the sloughing tissue, is very remarkable. The acid appears to have but little influence in removing the hard infiltration which may already exist; but, when properly applied, it almost certainly prevents its spread and increase, and so nearly limits the size of the sore to that which it has already attained.

In Mr. Walter's case, a slight local improvement appears to have followed even the superficial use of the carbolic acid in the form of lotion. PETER EADE, M.D., F.R.C.P., Norwich.

FRACTURE OF THE INTERNAL TABLE OF THE SKULL ALONE.

As this accident is of rare occurrence, I think the following may be worthy of record.

C. C., aged 35, was brought to the Sheffield General Infirmary, in a cab, on December 22nd, 1879, suffering from a scalp-wound. He was said to have been struck by the handle of a crane, heavily weighted, over which he had lost control.

On examination, I found a contused wound of the scalp, longitudinal in direction, and situated a little to the left of the median line in the fronto-parietal region. The pericranium was severed, for about half an inch, in a direction corresponding to the wound; but, upon careful examination by finger and probe, no fracture could be detected. The left edge of the wound was undermined, and detached from the pericranium beneath, as if the force of the blow had been partly expended in separating these structures. Statements made by eye-witnesses confirmed this view, leading me to suppose that the patient had been brushed, not directly felled, by the crane-handle.

The subject of the accident was a healthy man: he had not lost consciousness, and could walk, but was somewhat collapsed; his pulse, however, being good. On this account, he was advised to remain; but, having a good home, and a cab at the door, he preferred to become an out-patient. The friends having been warned to bring him at once should any untoward symptoms arise, his wound was dressed with carbolic oil, and the patient allowed to go.

He returned eight times during the course of the next fortnight, during which the wound was healing kindly. He walked on each of these visits, and complained of no symptoms. Five weeks after the