Published in final edited form as:

Matrix Biol. 2007 October ; 26(8): 587–596.

Control of Matrix Metalloproteinase Catalytic Activity

Hyun-Jeong Ra and William C. Parks

Center for Lung Biology, University of Washington, Seattle WA

Summary

As their name implies, MMPs were first described as proteases that degrade extracellular matrix proteins, such as collagens, elastin, proteoglycans, and laminins. However, studies of MMP function *in vivo* have revealed that these proteinases act on a variety of extracellular protein substrates, often to activate latent forms of effector proteins, such as antimicrobial peptides and cytokines, or to alter protein function, such as shedding of cell-surface proteins. Because their substrates are diverse, MMPs are involved in variety of homeostatic functions, such as bone remodeling, wound healing, and several aspects of immunity. However, MMPs are also involved in a number of pathological processes, such as tumor progression, fibrosis, chronic inflammation, tissue destruction, and more. A key step in regulating MMP proteolysis is the conversion of the zymogen into an active proteinase. Several proMMPs are activated in the secretion pathway by furin proprotein convertases, but for most the activation mechanisms are largely not known. In this review, we discuss both authentic and potential mechanisms of proMMP activation.

Keywords

Matrix metalloproteinase; zymogen; proteolysis; adaptor molecules

The Matrix Metalloproteinase Family

Metalloproteases, which are found in all organisms, are endopeptidases that contain an active site Zn^{2+} (hence, the prefix “metallo”) and are divided into subfamilies or clans based on evolutionary relationships and structure of the catalytic domain. The metzincin subfamily of metalloproteases are characterized by 3-histidine zinc-binding motif and a conserved methionine turn following the active site (Bode et al., 1993). The members of metzincin family are the repolysins or ADAMs, (A Disintegrin And Metalloproteinase), serralysins, astacins, and matrixins (aka: matrix metalloproteinases; MMPs).

MMPs comprise a family of currently 25 related, yet distinct vertebrate gene products, of which 24 are found in mammals (Table 1). MMPs are translated as a zymogen (i.e., an inactive enzyme) and contain a signal sequence peptide for targeting to secretory vesicles. Hence, MMPs are secreted or anchored to the cell surface, thereby confining their catalytic activity to membrane proteins or proteins within the secretory pathway or extracellular space. To be classified as an MMP, a protein must have at least the conserved pro and catalytic domains (Fig. 1). The prodomain of a typical MMP is about 80 amino acids and contains the consensus sequence PRCXXPD. The exception to this rule is MMP-23, in which the critical cysteine is

Corresponding Author: William C. Parks, Center for Lung Biology, 815 Mercer Street, Seattle, WA 98109 USA, Phone: 206-897-1303, Fax: 206-897-1546, Email: parksw@u.washington.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

found within a distinct run of amino acids (Velasco et al., 1999). The catalytic domain contains three conserved histidines in the sequence HEXHXXGXXH, which ligate the active site Zn^{2+} . The glutamate residue within the catalytic motif activates a zinc-bound H_2O molecule providing the nucleophile that cleaves peptide bonds. The cysteine-thiol and zinc ion interaction keeps proMMPs in a latent state.

With the exceptions of MMP-7, -23, and -26, MMPs have a flexible proline-rich hinge region and a hemopexin-like C-terminal domain, which functions in substrate recognition (Fig. 1). Other domains are restricted to subgroups of enzymes. For example, four membrane-type MMPs (MMP-14, -15, -16, -24) have transmembrane and cytosolic domains, whereas MMP-17 and -25 have C-terminal hydrophobic extensions that act as a glycosylphosphatidylinositol (GPI) anchoring signal. The two gelatinases (MMP-2 and MMP-9) have gelatin-binding domains that resemble similar motifs in fibronectin. In addition to a remarkably common 3-D structure (Massova et al., 1998), MMPs share a similar gene arrangement suggesting that they arose by duplications of an ancestor gene. At least eight of the known human MMP genes (MMP-1, -3, -7, -8, -10, -12, -13, and -20) are clustered on chromosome 11 at 11q21-23. Other MMP genes are scattered among chromosomes 1, 8, 12, 14, 16, 20, and 22 (Shapiro, 1998).

Regulation of MMP Activity

As their name implies, MMPs are thought to be responsible for the turnover and degradation of extracellular matrix, a function that is indeed performed by some family members, most notably MMP-14, a demonstrated physiologic collagenase (Holmbeck et al., 2004). However, matrix degradation is neither the shared nor predominant function of these enzymes. Several reports in recent years have demonstrated that various MMPs act on non-matrix proteins, such as cytokines, chemokines, receptors, antimicrobial peptides, and more (Parks et al., 2004). Thus, MMPs should not be viewed solely as proteinases of matrix catalysis, but rather as extracellular processing enzymes involved in regulating cell-cell and cell-matrix signaling events, quite typically, gain-of-function processing of latent proteins (Page-McCaw et al., 2007; Parks, 1999; Vu and Werb, 2000).

As for all secreted proteinases, the catalytic activity of MMPs is regulated at four points - gene expression, compartmentalization (i.e., pericellular accumulation of enzyme), proenzyme (or zymogen) activation, and enzyme inactivation - and is further controlled by substrate availability and affinity. Some members, such as MMP-2, MMP-19, MMP-28, and several MT-MMPs, are expressed in normal tissues, implying a role in homeostasis. However, most MMPs are not expressed in resting tissue yet are induced in repair or remodeling processes and in diseased or inflamed tissues. Although the qualitative patterns and quantitative levels of MMPs vary among tissues, diseases, tumors, inflammatory conditions, and cell types, a reasonable generalization is that activated cells, whether in tissue or in a culture dish, express a variety of MMPs. For the most part, the production of MMPs is regulated at the level of transcription by specific signals that are temporally limited and spatially confined.

In this review, we focus on two points of MMP regulation: 1) the known and potential non-furin mechanisms mediating activation of proMMPs, and 2) the known and potential mechanisms affecting enzyme activity, compartmentalization, and substrate specificity. In addition, we will emphasize *in vivo* mechanisms. As we discuss below, a lot of potential mechanisms of MMP activation have been suggested by defined *in vitro* studies. However, until validated, such approaches - though invaluable in understanding enzymology, protein-protein interactions, and much more - show how a zymogen can be activated, not how it is activated.

The Cysteine Switch

As stated, proMMPs are kept in a catalytically inactive state by the interaction between the thiol of the conserved prodomain cysteine residue and the zinc ion of the catalytic site. For a proMMP to become catalytically active, the thiol-Zn²⁺ interaction must be disrupted. In truly a classic paper, Harold Van Wart and Henning Birkdahl-Hansen proposed this process as general and required step in the activation of all proMMPs (1990), and they named this mechanism the “cysteine switch”, a term that remains valid and widely accepted. In essence, the thiol-Zn²⁺ interaction can be broken - and a latent MMP can gain catalytic activity - by three mechanisms: 1) direct cleavage of the pro-domain by another proteinase; 1) reduction of the free thiol by oxidants or by nonphysiologic reagents such as alkylating agents, heavy metal ions, and disulfides (Springman et al., 1990); and 3) allosteric perturbation of zymogen (Fig. 2). Thiol reduction and allosteric controls would lead to inter or intramolecular autolytic cleavage of the prodomain (Fig. 2).

Allosteric Activation

An important component of the cysteine-switch mechanism is that the prodomain does not need to be removed for a zymogen to acquire activity; only disruption of the zinc-thiol interaction is absolutely required. This concept is best illustrated in substrate zymography in which proMMPs (activated by SDS) are fully active. However, activation in a gel is an artificial system. Regardless of how the cysteine-Zn²⁺ interaction is disrupted, proteolysis of the prodomain is considered to be the final step in MMP activation *in vivo*. As is discussed below, proMMPs can interact with non-substrate macromolecules or be post-translationally modified, both of which can cause conformational changes favoring conversion to an activated state (Fig. 2). In turn, allosteric disruption of the cysteine-Zn²⁺ forms a transitional active state allowing autolytic cleavage of the prodomain (Fig. 2).

Furin Activation

Analogous to most pro-proteins, direct proteolysis is considered to be the principal way that proMMPs are activated. About one-third of MMPs - including all membrane-bound MMPs - contain an RXKR or RRKR sequence between the pro and catalytic domains, which serves as a target sequence for proprotein convertases or furins (Fig. 1). Furin is a type 1 membrane, subtilisin-like serine protease present in the *trans*-Golgi network (Thomas, 2002), and thus, MMPs with a furin cleavage site are processed intracellularly before secretion (Illman et al., 2003; Kang et al., 2002; Pei and Weiss, 1995; Sato et al., 1996). For the other MMPs, the mode of activation is more presumed than proved, and the *in vivo* mechanism for activation of most non-furin cleaved proMMPs is unknown.

Activation of ProMMPs by MMPs

In vitro, a number of MMPs can cleave the prodomain of other MMP zymogens leading to activation, and it has been suggested that active MMPs may mediate the final proteolytic step to produce a fully active enzyme (Nagase, 1997). The initial cleavage could be done out by either a serine proteinase or an MMP. For example, in a cell model of endothelial tube regression in collagen gels, activation of proMMP-1 requires both kallikrein (or plasmin) and MMP-10 (Saunders et al., 2005), but it is not clear if the activation is mediated by direct proteolysis or by mechanisms downstream of these proteinases.

Furthermore, there are three considerations that question the MMP-mediated activation mechanism. First, it is not clear if potentially activating MMPs are co-expressed or, and quite important, co-localized with their potential substrate proMMPs. For example, MMP-3 effectively activates proMMP-1 and proMMP-7 *in vitro* (Imai et al., 1995; Ito and Nagase, 1988; Murphy et al., 1987). However, full activation of proMMP-7 requires both a molar excess

of MMP-3 and an extended incubation, suggesting MMP-3 is not an efficient activator of MMP-7 (Imai et al., 1995). Furthermore, MMP-3 is not typically co-expressed with MMP-1 (Parks, 1999; Saarialho-Kere et al., 1994) nor with MMP-7 *in vivo* (Dunsmore et al., 1998; Halpert et al., 1996; Rudolph-Owen et al., 1997). Second, if co-localized, the putative activating MMP may not be present in sufficient relative amounts to cleave the zymogen, particularly in a pericellular microenvironment that contain high concentrations of other or authentic substrates. Third, we know of no studies in MMP null mice in which an observed phenotype has been linked or demonstrated to be due to inactivation of another MMP (Parks et al., 2004).

Despite these caveats, MMP activation of proMMPs may occur for some members. Possibly the best described non-furin proteolytic activation mechanism of a proMMP is cleavage of proMMP-2 at the cell surface by active MMP-14 (Caterina et al., 2000; Hernandez-Barrantes et al., 2000; Strongin et al., 1995; Wang et al., 2000). The collective data from different laboratories demonstrated that proMMP-2 requires cooperative action of MMP-14 and TIMP-2, which is typically considered to a natural MMP inhibitor, within a 1:1:1 stoichiometric ratio. Whereas the right amount of exogenous TIMP-2 enhances proMMP-2 activation (Cao et al., 1996; Strongin et al., 1995), excess amounts block activation (Cao et al., 1995; Imai et al., 1996; Itoh et al., 2001; Strongin et al., 1995; Toth et al., 2000; Ward et al., 1994; Worley et al., 2003). This activation mechanism involves the C-terminal end of TIMP-2 interacting with both the C-terminus of proMMP-2 and the N-terminus of activated MMP-14 (Cao et al., 1996; Cao et al., 1995; Sato et al., 1996; Strongin et al., 1993). Within the ternary complex, TIMP-2 brings zymogen and active proteinase in close proximity such that MMP-14 can cleave the pro-domain of proMMP-2.

Although the role of TIMP-2 in proMMP-2 activation is supported by a lack of zymogen activation in TIMP-2 null mice (Wang et al., 2000), a similar defect in MMP-14 knock-out mice has not been reported and activation of proMMP-2 in fibroblasts is not markedly, if at all affected by deficiency of MMP-14 (Ruangpanit et al., 2002). The role for MMP-14 in activation of MMP-2 is further questioned by a lack of phenocopy between null mouse lines. Whereas MMP-2 null mice reveal mild phenotypes, mostly related to neovascularization and inflammation (Berglin et al., 2003; Corry et al., 2002; Itoh et al., 2002; Itoh et al., 1998; Kato et al., 2001; Ohno-Matsui et al., 2003), MMP-14 knock-out mice have severe defects in skeletal development and turnover of type I collagen (Holmbeck et al., 1999; Holmbeck et al., 2003; Zhou et al., 2000). Overall, the *in vivo* data indicate that MMP-14 is not required for activation of MMP-2, but they do not rule the possibility that other membrane-type MMPs might be involved.

Activation of proMMPs by Plasmin and other Serine Proteinases

Several studies - much more than cited at the end of this sentence - have speculated that serine proteases, particularly plasmin, are *in vivo* activators of proMMPs, including some of the furin-activated MMPs (Carmeliet et al., 1997; Creemers et al., 2000; Monea et al., 2002; Okumura et al., 1997; Ramos-DeSimone et al., 1999). In fact, the concept that plasmin functions to activate latent MMPs is oft accepted as dogma (Janssens and Lijnen, 2006; Lo et al., 2003), however, the data in support of this mechanism is mostly circumstantial. For example, plasminogen-null mice (*plg*^{-/-}) present defects in wound healing after myocardial infarction, and the authors speculate that these defects are due to diminished activation of proMMP-2 and proMMP-9 in the absence of plasmin (Creemers et al., 2000). In another study, activation of proMMP-3, -9, -12, and -13 was impaired in urokinase-type plasminogen activator-null mice (*upa*^{-/-}), but not in tissue-type plasminogen activator-null mice (*tpa*^{-/-}) (Carmeliet et al., 1997). The authors conclude that u-PA activates plasminogen into plasmin, which subsequently activates this group of MMPs, however, this was demonstrated in cell lysates with exogenous

plasminogen. Other groups have reported that proMMP-9 is not activated by plasmin (Okada et al., 1992).

A rub with plasmin-mediated activation of proMMPs is fibrin. It is very clear that primary function of plasmin is to breakdown fibrin. Where deposited, the pericellular concentration of fibrin would be enormous, whereas MMPs tend to be produced at much more modest levels. Furthermore, the affinity of plasmin for fibrin is much higher than that for proMMPs. Thus, in a biologic setting, such as an inflamed tissue, where plasmin, fibrin deposition, and MMPs are all present, plasmin may not even “see” proMMPs. Indeed, Lim *et al.* (1996) reported that the regulation of plasmin and MMPs are not coordinated in HT1080 cells. Moreover, they demonstrated that plasmin does not activate proMMP-2 nor proMMP-14 directly and serpins do not prevent proMMP-2 activation in these cells. Additional evidence suggesting that plasmin does not function *in vivo* as in MMP activator comes from observations of mice lacking various components for the plasmin activation cascade. For example, in aortic extracts and conditioned medium of tissues and fibroblast from *u-pa*^{-/-}, *t-pa*^{-/-}, *plg*^{-/-} and *pai-1*^{-/-} mice, the amount of active MMP-2 and MMP-9 do not differ among genotypes (Lijnen et al., 1998), and Lund *et al.* reported that MMP-dependent activity in wound healing is not affected by a lack of plasmin or the plasminogen-activating enzyme (Lund et al., 1999). One group of serine proteases that do seem to activate proMMPs *in vivo* are mast cell chymases (Fang et al., 1997; Fang et al., 1999). Mice lacking mast cell chymase-4 (MCP-4) have diminished activation of proMMP-2 and proMMP-9 (Tchougounova et al., 2005). However, the absence of mMCP-4 does not completely abolish the activation of these two MMPs by mast cells. Thus, the limited number of MMPs affected by plasmin or mMCP-4 further indicates that other means of activation are necessary.

Oxidative Control of MMP Activity

Oxidants generated by leukocytes or other cells can both activate (via oxidation of the prodomain thiol followed by autolytic cleavage) and subsequently inactivate MMPs (via modification of amino acids critical for catalytic activity), providing a mechanism to control quantum bursts of proteolytic activity. *In vitro*, a number of proMMPs are activated by reactive oxygen species (ROS), which can modify the cysteine thiol group and activate proMMPs via autolytic cleavage. For example, hypochlorous acid (HOCl), a product of neutrophil myeloperoxidase (MPO), and hydroxyl radicals activate proMMPs-1, -7, and -9 (Fu et al., 2001; Michaelis et al., 1992; Peppin and Weiss, 1986; Weiss et al., 1985), and peroxynitrate can activate proMMPs-1, -2, and -9 (Okamoto et al., 1997). However, a role for most ROS as *in vivo* activators of MMPs has not been established. To date, the only *in vivo* evidence we are aware of is in ischemic mouse brains, where activation of proMMP-9 requires *s*-nitrosylation of the pro-domain (Gu et al., 2002). Even though MPO and MMP-7 colocalize within lipid-laden atherosclerotic plaques (Fu et al., 2001), in these collaborative studies, we have not yet established if MPO is required for proMMP-7 activation *in vivo*.

In addition to promoting autolytic activation, Fu and colleagues have demonstrated that whereas low concentrations of HOCl activate proMMP-7, higher concentrations or extended exposure potentially inactivate the active enzyme (Fu et al., 2004; Fu et al., 2001; Fu et al., 2003). Thus, it is possible that the pericellular production of HOCl by phagocytes provides a physiological mechanism for regulating both the activation and inactivation of MMPs within an inflammatory setting (where many MMPs are usually expressed). Interestingly, this model implies temporal control over enzyme activity: The cysteine-switch thiol in the prodomain is the preferred target of HOCl, and its oxygenation leads rapid activation (which may proceed before autolytic cleavage of the prodomain). Sustained HOCl production would modify additional sites, and once critical residues are hit, the enzyme would lose its catalytic activity. Thus, ROS could confine MMP activity to quantum bursts of pericellular proteolysis that are

regulated in space and time. Whether inactivation predominates over proenzyme activation would depend on ROS generation and concentration, which would correlate closely with the extent of inflammation. This mechanism provides an alternative pathway of MMP inactivation typically ascribed to protein inhibitors, such as TIMPs (Woessner, 1991; Woessner, 1998).

Compartmentalization

In vitro studies have demonstrated significant overlap in the substrates MMPs can cleave, particularly among the extracellular matrix (ECM) substrates (Sternlicht and Werb, 2001). In disease settings, like cancer and chronic inflammation, in which essentially all MMPs are present, the shared substrate potential among enzymes would seemingly permit biochemical redundancy. However, substrate selectivity can be honed by two processes: enzyme affinity and compartmentalization. Kinetic studies with model substrates have demonstrated that specific enzymes degrade some substrates more efficiently than others. For example, both MMP-2 and MMP-9 act on cleaved collagen better than other MMPs (Mackay et al., 1990) and human MMP-7 is a more potent proteoglycanase and elastase than other MMPs (Filippov et al., 2003; Halpert et al., 1996). Thus, in a tissue environment, which contains many potential substrates, selectivity of MMP catalysis, in addition to being regulated by the concentration of active enzyme, may be directed by the concentration of a preferred substrate relative to that of other potential substrates within range of a secreted MMP.

Compartmentalization, that is where and how in the pericellular environment an MMP is released and held, is equally important, if not more so, for regulating the specificity of proteolysis than is the affinity of the enzyme-substrate interaction. Analogous to MMPs with a transmembrane domain, the so-called “secreted” MMPs likely require a cellular compartment-specific localization mechanism to function properly. An important concept is that cells do not indiscriminately release proteases. Rather, proteinases, such as MMPs, would likely be anchored to the cell membrane, thereby maintaining a locally high enzyme concentration and targeting their catalytic activity to specific substrates within the pericellular space. In addition to the membrane-bound MMPs, several examples of specific cell-MMP interactions have been reported, such as the binding of MMP-2 to the $\alpha_v\beta_3$ integrin (Brooks et al., 1996), MMP-1 to the $\alpha_2\beta_1$ integrin (Dumin et al., 2001; Stricker et al., 2001), MMP-9 to CD44 (Yu and Stamenkovic, 2000), and MMP-7 to surface proteoglycans (Yu and Woessner, 2000; Yu et al., 2002), cholesterol (Yamamoto et al., 2006), and CD151 (Shiomi et al., 2005). As suggested for CD44 (Yu and Stamenkovic, 2000) and the $\alpha_2\beta_1$ integrin (Dumin et al., 2001), these membrane anchors may act as accessory factors that mediate both pro-enzyme activation and binding of both substrate and proteinase, thereby increasing the probability of proteolysis (Fig. 2). Thus, as for most protein-protein interactions, MMP specificity may be driven by a third component, if not higher order. It is likely that other MMPs are also attached to cells or matrix via similar specific interactions, and determining these anchors will be a key advance toward identifying activation mechanisms and substrates.

Anchoring “released” MMPs to the cell surface could promote zymogen activation via allosteric disruption of the cysteine-zinc bond and autolytic cleavage of the prodomain (Fig. 3). For example, Shiomi et al. (2005) reported that CD151, a tetraspanin, binds and activates proMMP-7. The authors speculate that CD151 causes a conformational change in proMMP-7, thus facilitating a spontaneous activation of proMMP-7 on a pericellular level. Interestingly, activated MMP-7 was only detected in the presence of carboxymethylated transferrin, a preferred *in vitro* substrate of MMP-7 (Shiomi et al., 2005). Thus, membrane anchors may function by facilitating the interaction between substrate and proteinase, and in the presence of substrate, the allosteric release of the cysteine switch may be more favored. Although these may be attractive ideas, they still need to be tested experimentally.

GAGs and MMPs

One class of cell surface molecules that may prove to be quite important in regulating activation and activity of MMPs are proteoglycans, which contain glycosaminoglycan (GAG) chains attached to a protein core. Most of cell surface proteoglycans contain either heparan sulfate (HS) or chondroitin sulfate (CS) GAG chains. Proteoglycans interact with a wide variety of effector proteins, such as chemokines and growth factors, via their GAG chains to elicit various processes (Park et al., 2000), and their ability to control proteinase activity, particularly of serine proteinases, is well established. For example, heparin, a mast cell GAG that is a more sulfated than HS, affects the activity of several coagulation cascade proteinases, such as antithrombin III, thrombin, and factor Xa (Capila and Linhardt, 2002; Conrad, 1998), as well as tryptase, chymase, and cathepsin B (Almeida et al., 2001; Alter et al., 1987; Hallgren et al., 2000; Pejler and Sadler, 1999).

Heparin within mast cells is associated with serglycin, which is present in secretion vesicles of many cell types and is often produced as CS proteoglycan (Stevens et al., 1988; Stevens et al., 2000). In serglycin-null mice ($sg^{-/-}$), which are otherwise normal, mast cell proteases are missorted and have reduced activity (Åbrink et al., 2004). Because of its location within the secretion pathway and its established role in interacting with and regulating serine proteinases, serglycin may also regulate MMP zymogen activation. In THP-1 cells, a macrophage cell line, proMMP-9 is covalently linked to CS proteoglycans, most likely serglycin, which induces processing of proMMP-9 at both its N- and C-termini (Winberg et al., 2003; Winberg et al., 2000). However, this activation of proMMP-9 does not seem to require CS chains. In our studies, we have found that proMMP-7 is co-packaged with serglycin in secretion vesicles, and that autolytic activation and activity toward authentic substrates are markedly accelerated by CS GAGs (unpublished observations). One important future direction for the MMP field will be to explore the relationship between proteoglycans and MMPs *in vivo*, not only to elucidate mechanisms of activation, but also to investigate how proteoglycans may function confine secreted MMPs and substrates within the same pericellular compartments.

Significance

Because they have the ability to act on extracellular matrix proteins, MMPs have been often implicated as contributing to tumor progression and metastasis, supposedly by breaking down tissue barriers that would otherwise restrain invasion (Coussens et al., 2002; Egeblad and Werb, 2002). Consequently, many pharmaceutical companies developed programs some years back to target MMPs in cancer. Several drugs, all designed to directly block MMP catalytic activity, were tested in phase III clinical trials; however, none were effective (Coussens et al., 2002). In fact, most patients in these studies, all of whom had advanced stage adenocarcinomas, complained of painful side-effects and requested to be taken off of the drug. In one study, patients taking the MMP inhibitor died at a rate higher than those on placebo.

The key shortcoming of the MMP inhibitor trials is that these drugs lack specificity (Coussens et al., 2002; Parks et al., 2004). Not only do they block the activity of all 24 MMPs, these compounds also inhibit the activity of the related ADAM and ADAMTS proteinases as well as other metalloenzymes. Despite the largely disappointing clinical trials, several studies in mice and cell models have convincingly demonstrated that *specific* MMPs are indeed essential mediators of tumor progression and invasion (Coussens et al., 2002; Hotary et al., 2003; Hotary et al., 2002; Wilson et al., 1997). In a complex tissue environment, like cancer, several cell types (resident, inflammatory, tumor) express several, different MMPs, and these proteinase can either promote or restrain disease or repair processes by affecting multiple and apparently opposing processes by the same cell at the same time (Coussens et al., 2002; Egeblad and Werb, 2002; Parks et al., 2004). For example, we have determined that epithelial MMP-7 is required

for wound closure and generation of antimicrobial activity (Dunsmore et al., 1998; Wilson et al., 1999) (clearly beneficial functions), but it also promotes neutrophil activation, which can be damaging and lethal (Li et al., 2002; McGuire et al., 2003). It is likely that to effect on different processes at the same time, a cell uses distinct mechanisms to compartmentalize an MMP to different substrates. In addition, if compartmentalization and zymogen activation mechanisms are linked, if not indeed the same, then understanding how proMMPs are activated and anchored may provide a means to blocked a potentially deleterious MMP-mediated process while sparing a beneficial function.

References

- Åbrink M, Grujic M, Pejler G. Serglycin is essential for maturation of mast cell secretory granule. *J Biol Chem* 2004;279:40897–40905. [PubMed: 15231821]
- Almeida PC I, Nantes L, Chaga JR, Rizzi CCA, Faljoni-Alario A, Carmona E, Juliano L, Nader HB, Tersariol ILS. Cathepsin B activity regulation : Heparin-like glycoaminoglycans protect human cathepsin B from alkaline pH-induced inactivation. *J Biol Chem* 2001;276:944–951. [PubMed: 11016923]
- Alter SC, Metcalfe DD, Bradford TR, Schwartz LB. Regulation of human mast cell tryptase. Effect of enzyme concentration, ionic strength and the structure and negative charge density of polysaccharides. *Biochem J* 1987;248:821–827. [PubMed: 2449172]
- Balbin M, Fueyo A, Knauper V, Lopez JM, Alvarez J, Sanchez LM, Quesada V, Bordallo J, Murphy G, Lopez-Otin C. Identification and enzymatic characterization of two diverging murine counterparts of human interstitial collagenase (MMP-1) expressed at sites of embryo implantation. *J Biol Chem* 2001;276:10253–62. [PubMed: 11113146]
- Berglin L, Sarman S, van der Ploeg I, Steen B, Ming Y, Itohara S, Seregard S, Kvanta A. Reduced choroidal neovascular membrane formation in matrix metalloproteinase-2-deficient mice. *Invest Ophthalmol Vis Sci* 2003;44:403–8. [PubMed: 12506102]
- Bode W, Gomis-Rüth FX, Stockler W. Astacins, serralytins, snake venom and matrix metalloproteinases exhibit identical zinc-binding environments (HEXXHXXGXXH and Met-turn) and topologies and should be grouped into a common family, the ‘metzincins’. *FEBS Lett* 1993;331:134–40. [PubMed: 8405391]
- Brooks PC, Stromblad S, Sanders LC, von Schalscha TL, Aimes RT, Stetler-Stevenson WG, Quigley JP, Chersesh DA. Localization of matrix metalloproteinase MMP-2 to the surface of invasive cells by interaction with integrin alpha v beta 3. *Cell* 1996;85:683–693. [PubMed: 8646777]
- Cao J, Rehemtulla A, Bahou W, Zucker S. Membrane type matrix metalloproteinase 1 activates pro-gelatinase A without furin cleavage of the N-terminal domain. *J Biol Chem* 1996;271:30174–30180. [PubMed: 8939968]
- Cao J, Sata H, Takino T, Seiki M. The C-terminal region of membrane type matrix metalloproteinase is a functional transmembrane domain required for pro-gelatinase A activation. *J Biol Chem* 1995;270:801–805. [PubMed: 7822314]
- Capila I, Linhardt RJ. Heparin-protein interactions. *Agnew Chem Int Ed* 2002;41:390–412.
- Carmeliet P, Moons L, Lijnen R, Baes M, Lemaître V, Tipping P, Drew A, Eeckhout Y, Shapiro S, Lupu F, Collen D. Urokinase-generated plasmin activates matrix metalloproteinases during aneurysm formation. *Nat Genet* 1997;17:439–444. [PubMed: 9398846]
- Caterina JJ, Yamada S, Caterina NC, Longenecker G, Holmback K, Shi J, Yermovsky AE, Engler JA, Birkedal-Hansen H. Inactivating mutation of the mouse tissue inhibitor of metalloproteinases-2 (Timp-2) gene alters proMMP-2 activation. *J Biol Chem* 2000;275:26416–26422. [PubMed: 10827176]
- Clark HF, Gurney AL, Abaya E, Baker K, Baldwin D, Brush J, Chen J, Chow B, Chui C, Crowley C, Currell B, Deuel B, Dowd P, Eaton D, Foster J, Grimaldi C, Gu Q, Hass PE, Heldens S, Huang A, Kim HS, Klimowski L, Jin Y, Johnson S, Lee J, Lewis L, Liao D, Mark M, Robbie E, Sanchez C, Schoenfeld J, Seshagiri S, Simmons L, Singh J, Smith V, Stinson J, Vagts A, Vandlen R, Watanabe C, Wieand D, Woods K, Xie MH, Yansura D, Yi S, Yu G, Yuan J, Zhang M, Zhang Z, Goddard A, Wood WI, Godowski P, Gray A. The secreted protein discovery initiative (SPDI), a large-scale effort

- to identify novel human secreted and transmembrane proteins: a bioinformatics assessment. *Genome Res* 2003;13:2265–70. [PubMed: 12975309]
- Conrad, HE. Heparin-binding proteins. Academic Press; San Diego: 1998.
- Corry DB, Rishi K, Kanellis J, Kiss A, Song LZ, Xu J, Feng L, Werb Z, Kheradmand F. Decreased allergic lung inflammatory cell egression and increased susceptibility to asphyxiation in MMP2-deficiency. *Nat Immunol* 2002;3:347–353. [PubMed: 11887181]
- Coussens LM, Fingleton B, Matrisian LM. Matrix metalloproteinase inhibitors and cancer: trials and tribulations. *Science* 2002;295:2387–2392. [PubMed: 11923519]
- Creemers E, Cleutjens J, Smits J, Heymans S, Moons L, Collen D, Daemen M, Carmeliet P. Disruption of the plasminogen gene in mice abolishes wound healing after myocardial infarction. *Am J Pathol* 2000;156:1865–1873. [PubMed: 10854210]
- Dumin JA, Dickeson SK, Stricker TP, Bhattacharyya-Pakrasi M, Roby JD, Santoro SA, Parks WC. Procollagenase-1 (matrix metalloproteinase-1) binds the integrin $\alpha 2\beta 1$ upon release from keratinocytes migrating on type I collagen. *J Biol Chem* 2001;276:29368–29374. [PubMed: 11359786]
- Dunsmore SE, Saarialho-Kere UK, Roby JD, Wilson CL, Matrisian LM, Welgus HG, Parks WC. Matrilysin expression and function in airway epithelium. *J Clin Invest* 1998;102:1321–1331. [PubMed: 9769324]
- Egeblad M, Werb Z. New functions for the matrix metalloproteinases in cancer progression. *Nat Rev Cancer* 2002;2:161–174. [PubMed: 11990853]
- Fang KC, Raymond WW, Blount JL, Caughey GH. Dog mast cell alpha-chymase activates progelatinase B by cleaving the Phe88-Gln89 and Phe91-Glu92 bonds of the catalytic domain. *J Biol Chem* 1997;272:25628–35. [PubMed: 9325284]
- Fang KC, Wolters PJ, Steinhoff M, Bidgol A, Blount JL, Caughey GH. Mast cell expression of gelatinases A and B is regulated by kit ligand and TGF-beta. *J Immunol* 1999;162:5528–35. [PubMed: 10228034]
- Filippov S, Caras I, Murray R, Matrisian LM, Chapman HA Jr, Shapiro S, Weiss SJ. Matrilysin-dependent elastolysis by human macrophages. *J Exp Med* 2003;198:925–35. [PubMed: 12963695]
- Fu X, Kao JL, Bergt C, Kassim SY, Huq NP, d'Avignon A, Parks WC, Mecham RP, Heinecke JW. Oxidative cross-linking of tryptophan to glycine restrains matrix metalloproteinase activity: specific structural motifs control protein oxidation. *J Biol Chem* 2004;279:6209–12. [PubMed: 14670964]
- Fu X, Kassim SY, Parks WC, Heinecke JW. Hypochlorous acid oxygenates the cysteine switch domain of pro-matrilysin (MMP-7). A mechanism for matrix metalloproteinase activation and atherosclerotic plaque rupture by myeloperoxidase. *J Biol Chem* 2001;276:41279–41287. [PubMed: 11533038]
- Fu X, Kassim SY, Parks WC, Heinecke JW. Hypochlorous acid generated by myeloperoxidase modifies adjacent tryptophan and glycine residues in the catalytic domain of matrix metalloproteinase-7 (matrilysin): an oxidative mechanism for restraining proteolytic activity during inflammation. *J Biol Chem* 2003;278:28403–28409. [PubMed: 12759346]
- Gu Z, Kaul M, Yan B, Kridel SJ, Cui J, Strongin A, Smith JW, Liddington RC, Lipton SA. S-nitrosylation of matrix metalloproteinases: signaling pathway to neuronal cell death. *Science* 2002;297:1186–1190. [PubMed: 12183632]
- Hallgren J, Karlson U, Poorafshar M, Hellman L, Pejler G. Mechanism for activation of mouse mast cell tryptase: Dependence of heparin and acidic pH for formation of active tetramers of mouse mast cello protease 6. *Biochemistry* 2000;39:13068–13077. [PubMed: 11041873]
- Halpert I, Roby JD, Sires UI, Potter-Perigo S, Wight TN, Welgus HG, Shapiro SD, Wickline SA, Parks WC. Matrilysin is expressed by lipid-laden macrophages at sites of potential rupture in atherosclerotic lesions and localizes to areas of versican deposition, a proteoglycan substrate for the enzyme. *Proc Natl Acad Sci USA* 1996;93:9748–9753. [PubMed: 8790402]
- Hernandez-Barrantes S, Toth M, Bernardo MM, Yurkova M, Gervasi DC, Raz Y, Sang QA, Fridman R. Binding of active (57 kDa) membrane type 1-matrix metalloproteinase (MT1-MMP) to tissue inhibitor of metalloproteinase (TIMP)-2 regulates MT1-MMP processing and pro-MMP-2 activation. *J Biol Chem* 2000;275:12080–12089. [PubMed: 10766841]
- Holmbeck K, Bianco P, Caterina J, Yamada S, Kromer M, Kuznetsov SA, Mankani M, Robey PG, Poole AR, Pidoux I, Ward JM, Birkedal-Hansen H. MT1-MMP-deficient mice develop dwarfism,

- osteopenia, arthritis, and connective tissue disease due to inadequate collagen turnover. *Cell* 1999;99:81–92. [PubMed: 10520996]
- Holmbeck K, Bianco P, Chrysovergis K, Yamada S, Birkedal-Hansen H. MT1-MMP-dependent, apoptotic remodeling of unmineralized cartilage: a critical process in skeletal growth. *J Cell Biol* 2003;163:661–71. [PubMed: 14610065]
- Holmbeck K, Bianco P, Yamada S, Birkedal-Hansen H. MT1-MMP: a tethered collagenase. *J Cell Physiol* 2004;200:11–9. [PubMed: 15137053]
- Hotary KB, Allen ED, Brooks PC, Datta NS, Long MW, Weiss SJ. Membrane type I matrix metalloproteinase usurps tumor growth control imposed by the three-dimensional extracellular matrix. *Cell* 2003;114:33–45. [PubMed: 12859896]
- Hotary KB, Yana I, Sabeh F, Li XY, Holmbeck K, Birkedal-Hansen H, Allen ED, Hiraoka N, Weiss SJ. Matrix metalloproteinases (MMPs) regulate fibrin-invasive activity via MT1-MMP-dependent and -independent processes. *J Exp Med* 2002;195:295–308. [PubMed: 11828004]
- Illman SA, Keski-Oja J, Parks WC, Lohi J. Mouse epilysin (MMP-28) is alternatively spliced and processed by a furin-like pro-protein convertase. *Biochemistry* 2003;375:191–197.
- Imai K, Ohuchi E, Aoki T, Nomura H, Fujii Y, Sato H, Seiki M, Okada Y. Membrane-type matrix metalloproteinase 1 is a gelatinolytic enzyme and is secreted in a complex with tissue inhibitor of metalloproteinases 2. *Cancer Res* 1996;56:2707–2710. [PubMed: 8665498]
- Imai K, Yokohama Y, Nakanishi I, Ohuchi E, Fuji Y, Nakai N, Okada Y. Matrix Metalloproteinase 7 (Matrilysin) from human rectal carcinoma cells: activation of the precursor, interaction with other matrix metalloproteinases and enzyme properties. *J Biol Chem* 1995;270:6691–6697. [PubMed: 7896811]
- Ito A, Nagase H. Evidence that human rheumatoid synovial matrix metalloproteinase 3 is an endogenous activator of procollagenase. *Arch Biochem Biophys* 1988;267:211–6. [PubMed: 2848449]
- Itoh T, Matsuda H, Tanioka M, Kuwabara K, Itohara S, Suzuki R. The role of matrix metalloproteinase-2 and matrix metalloproteinase-9 in antibody-induced arthritis. *J Immunol* 2002;169:2643–7. [PubMed: 12193736]
- Itoh T, Tanioka M, Yoshida H, Yoshioka T, Nishimoto H, Itohara S. Reduced angiogenesis and tumor progression in gelatinase A-deficient mice. *Cancer Res* 1998;58:1048–1051. [PubMed: 9500469]
- Itoh Y, Takamura A, Ito N, Maru Y, Sato H, Suenaga N, Aoki T, Seiki M. Homophilic complex formation of MT1-MMP facilitates proMMP-2 activation on the cell surface and promotes tumor cell invasion. *EMBO J* 2001;20:4782–4793. [PubMed: 11532942]
- Janssens S, Lijnen HR. What has been learned about the cardiovascular effects of matrix metalloproteinases from mouse models? *Cardiovasc Res* 2006;69:585–94. [PubMed: 16426591]
- Kang T, Nagase H, Pei D. Activation of membrane -type matrix metalloproteinase 3 zymogen by the proprotein convertase furin in the trans-golgi network. *Cancer Res* 2002;62:675–681. [PubMed: 11830519]
- Kato T, Kure T, Chang JH, Gabison EE, Itoh T, Itohara S, Azar DT. Diminished corneal angiogenesis in gelatinase A-deficient mice. *FEBS Lett* 2001;508:187–90. [PubMed: 11718713]
- Li Q, Park PW, Wilson CL, Parks WC. Matrilysin shedding of syndecan-1 regulates chemokine mobilization and transepithelial efflux of neutrophils in acute lung injury. *Cell* 2002;111:635–46. [PubMed: 12464176]
- Lijnen HR, Van Hoef B, Lupu F, Moons L, Carmeliet P, Collen D. Function of the plasminogen/plasmin and matrix metalloproteinase systems after vascular injury in mice with targeted inactivation of fibrinolytic system genes. *Arterioscler Thromb Vasc Biol* 1998;18:1035–1045. [PubMed: 9672063]
- Lim YT, Sugiura Y, Laug WE, Sun B, Garcia A, DeClerck YA. Independent regulation of matrix metalloproteinases and plasminogen activators in human fibrosarcoma cells. *J Cell Physiol* 1996;167:333–340. [PubMed: 8613475]
- Lo EH, Dalkara T, Moskowitz MA. Mechanisms, challenges and opportunities in stroke. *Nat Rev Neurosci* 2003;4:399–415. [PubMed: 12728267]
- Lund LR, Romer J, Bugge TH, Nielsen BS, Frandsen TL, Degen JL, Stephens RW, Dano K. Functional overlap between two classes of matrix-degrading proteases in wound healing. *Embo J* 1999;18:4645–56. [PubMed: 10469644]

- Mackay AR, Hartzler JL, Pelina MD, Thorgeirsson UP. Studies on the ability of 65-kDa and 92-kDa tumor cell gelatinases to degrade type IV collagen. *J Biol Chem* 1990;265:21929–21934. [PubMed: 2174891]
- Massova I, Kotra LP, Fridman R, Mobashery S. Matrix metalloproteinases: structures, evolution, and diversification. *FASEB J* 1998;12:1075–95. [PubMed: 9737711]
- McGuire JK, Li Q, Parks WC. Matrilysin (matrix metalloproteinase-7) mediates E-cadherin ectodomain shedding in injured lung epithelium. *Am J Pathol* 2003;162:1831–43. [PubMed: 12759241]
- Michaelis J, Vissers MC, Winterbourn CC. Different effects of hypochlorous acid on human neutrophil metalloproteinases: activation of collagenase and inactivation of collagenase and gelatinase. *Arch Biochem Biophys* 1992;292:555–62. [PubMed: 1309976]
- Monea S, Lehti K, Keski-Oja J, Mignatti P. Plasmin activates pro-Matrix Metalloproteinase-2 with a Membrane-Type 1 Matrix Metalloproteinase-Dependent mechanism. *J Cell Physiol* 2002;192:160–170. [PubMed: 12115722]
- Murphy G, Cockett MI, Stephens PE, Smith BJ, Docherty AJ. Stromelysin is an activator of procollagenase. A study with natural and recombinant enzymes. *Biochem J* 1987;248:265–8. [PubMed: 2829822]
- Nagase H. Activation mechanisms of Matrix Metalloproteinases. *Biol Chem* 1997;378:151–160. [PubMed: 9165065]
- Ohno-Matsui K, Uetama T, Yoshida T, Hayano M, Itoh T, Morita I, Mochizuki M. Reduced retinal angiogenesis in MMP-2-deficient mice. *Invest Ophthalmol Vis Sci* 2003;44:5370–5. [PubMed: 14638740]
- Okada Y, Gonoji Y, Naka K, Tomita K, Nakanishi I, Iwata K, Yamashita K, Hayakawa T. Matrix metalloproteinase 9 (92-kDa gelatinase/type IV collagenase) from HT 1080 human fibrosarcoma cells. Purification and activation of the precursor and enzymic properties. *J Biol Chem* 1992;267:21712–9. [PubMed: 1400481]
- Okamoto T, Akaike T, Nagano T, Miyajima S, Suga M, Ando M, Ichimori K, Maeda H. Activation of human neutrophil procollagenase by nitrogen dioxide and peroxynitrite: A novel mechanism for procollagenase activation involving nitric oxide. *Arch Biochem Biophys* 1997;342:261–274. [PubMed: 9186487]
- Okumura Y, Sato H, Seiki M, Kido H. Proteolytic activation of the precursor of membrane type 1 matrix metalloproteinase by human plasmin- a possible cell surface activator. *FEBS Lett* 1997;402:181–184. [PubMed: 9037191]
- Page-McCaw A, Ewald AJ, Werb Z. Matrix metalloproteinases and the regulation of tissue remodelling. *Nat Rev Mol Cell Biol* 2007;8:221–33. [PubMed: 17318226]
- Park PW, Reizes O, Bernfield M. Cell surface heparan sulfate proteoglycans: selective regulators of ligand-receptor encounters. *J Biol Chem* 2000;275:29923–29926. [PubMed: 10931855]
- Parks WC. Matrix metalloproteinases in repair. *Wound Repair Regen* 1999;7:423–432. [PubMed: 10633001]
- Parks WC, Wilson CL, Lopez-Boado YS. Matrix metalloproteinases as modulators of inflammation and innate immunity. *Nat Rev Immunol* 2004;4:617–629. [PubMed: 15286728]
- Pei D, Weiss SJ. Furin-dependent intracellular activation of the human stromelysin-3 zymogen. *Nature* 1995;375:244–247. [PubMed: 7746327]
- Pejler G, Sadler JE. Mechanism by which heparin proteoglycan modulates mast cell chymase activity. *Biochemistry* 1999;38:12187–12195. [PubMed: 10508424]
- Peppin GJ, Weiss SJ. Activation of the endogenous metalloproteinase, gelatinase, by triggered human neutrophils. *Proc Natl Acad Sci USA* 1986;83:4322–6. [PubMed: 3012563]
- Ramos-DeSimone N, Hahn-Dantona E, Siple J, Nagase H, French DL, Quigley JP. Activation of matrix metalloproteinase-9 (MMP-9) via a converging plasmin/stromelysin-1 cascade enhances tumor cell invasion. *J Biol Chem* 1999;274:13066–13076. [PubMed: 10224058]
- Ruangpanit N, Price JT, Holmbeck K, Birkedal-Hansen H, Guenzler V, Huang X, Chan D, Bateman JF, Thompson EW. MT1-MMP-dependent and -independent regulation of gelatinase A activation in long-term, ascorbate-treated fibroblast cultures: regulation by fibrillar collagen. *Exp Cell Res* 2002;272:109–18. [PubMed: 11777335]

- Rudolph-Owen LA, Hulboy DL, Wilson CL, Mudgett J, Matrisian LM. Coordinate expression of matrix metalloproteinase family members in the uterus of normal, matrilysin-deficient, and stromelysin-1-deficient mice. *Endocrinology* 1997;138:4902–4911. [PubMed: 9348221]
- Saarialho-Kere UK, Kovacs SO, Pentland AP, Parks WC, Welgus HG. Distinct populations of keratinocytes express stromelysin-1 and -2 in chronic wounds. *J Clin Invest* 1994;94:79–88. [PubMed: 8040294]
- Sato H, Kinoshita T, Takino T, Nakayama K, Seiki M. Activation of a recombinant membrane type 1-matrix metalloproteinase (MT1-MMP) by furin and its interaction with tissue inhibitor of metalloproteinases (TIMP-2). *FEBS Lett* 1996;393:101–104. [PubMed: 8804434]
- Saunders WB, Bayless KJ, Davis GE. MMP-1 activation by serine proteases and MMP-10 induces human capillary tubular network collapse and regression in 3D collagen matrices. *J Cell Sci* 2005;118:2325–40. [PubMed: 15870107]
- Shapiro SD. Matrix metalloproteinase degradation of extracellular matrix: biological consequences. *Curr Opin Cell Biol* 1998;10:602–608. [PubMed: 9818170]
- Shiomi T, Inoki I, Kataoka F, Ohtsuka T, Hashimoto G, Nemori R, Okada Y. Pericellular activation of proMMP-7 (promatrilysin-1) through interaction with CD151. *Lab Invest* 2005;85:1489–1506. [PubMed: 16200075]
- Springman EB, Angleton EL, Birkedal-Hansen H, Van Wart HE. Multiple modes of activation of latent human fibroblast collagenase: Evidence for the role of a Cys 73 active-site zinc complex in latency and a “cysteine switch” mechanism for activation. *Proc Natl Acad Sci USA* 1990;87:364–368. [PubMed: 2153297]
- Sternlicht MD, Werb Z. How matrix metalloproteinases regulate cell behavior. *Annu Rev Cell Dev Biol* 2001;17:463–516. [PubMed: 11687497]
- Stevens RL, Fox CC, Lichtenstein LM, Austen KF. Identification of chondroitin sulfate E proteoglycans and heparin proteoglycans in the secretory granules of human lung mast cells. *Proc Natl Acad Sci USA* 1988;85:2284–2287. [PubMed: 3353378]
- Stevens, RL.; Wong, GW.; Humphries, DE. Serglycin Proteoglycans: The family of proteoglycans stored in the secretory granules of certain effector cells of the immune system. In: Iozzo, RV., editor. *Proteoglycans Structure, Biology and Molecular interactions*. Marcel Dekker; New York: 2000. p. 177-199.
- Stricker TP, Dumin JA, Dickeson SK, Chung L, Nagase H, Parks WC, Santoro SA. Structural analysis of the α_2 integrin I domain:pro:collagenase-1 (MMP-1) interaction. *J Biol Chem* 2001;276:29375–29381. [PubMed: 11359774]
- Strongin A, Marmer BL, Grant GA, Goldberg GI. Plasma membrane-dependent activation of the 72-kDa type IV collagenase is prevented by complex formation with TIMP-2. *J Biol Chem* 1993;268:14033–14039. [PubMed: 8314771]
- Strongin AY, Collier I, Bannikov G, Marmer BL, Grant GA, Goldberg GI. Mechanism of cell surface activation of 72-kDa type IV collagenase. Isolation of the activated form of the membrane metalloprotease. *J Biol Chem* 1995;270:5331–5338. [PubMed: 7890645]
- Tchougounova E, Lundquist A, Fajardo I, Winberg JO, Abrink M, Pejler G. A key role for mast cell chymase in the activation of pro-matrix metalloprotease-9 and pro-matrix metalloprotease-2. *J Biol Chem* 2005;280:9291–9296. [PubMed: 15615702]
- Thomas G. Furin at the cutting edge: from protein traffic to embryogenesis and disease. *Nat Rev Mol Cell Biol* 2002;3:753–66. [PubMed: 12360192]
- Toth M, Bernardo MM, Gervasi DC, Soloway PD, Wang Z, Bigg HF, Overall CM, DeClerck YA, Tschesche H, Cher ML, Brown S, Mobashery S, Fridman R. Tissue inhibitor of metalloproteinase (TIMP)-2 acts synergistically with synthetic matrix metalloproteinase (MMP) inhibitors but not with TIMP-4 to enhance the (Membrane type 1)-MMP-dependent activation of pro-MMP-2. *J Biol Chem* 2000;275:41415–23. [PubMed: 10998420]
- Van Wart HE, Birkedal-Hansen H. The cysteine switch: A principle of regulation of metalloproteinase activity with potential applicability to the entire matrix metalloproteinase gene family. *Proc Natl Acad Sci USA* 1990;87:5578–5582. [PubMed: 2164689]
- Velasco G, Pendas AM, Fueyo A, Knauper V, Murphy G, Lopez-Otin C. Cloning and characterization of human MMP-23, a new matrix metalloproteinase predominantly expressed in reproductive tissues

- and lacking conserved domains in other family members. *J Biol Chem* 1999;274:4570–4576. [PubMed: 9988691]
- Vu TH, Werb Z. Matrix metalloproteinases: effectors of development and normal physiology. *Genes Dev* 2000;14:2123–2133. [PubMed: 10970876]
- Wang Z, Juttermann R, Soloway PD. TIMP-2 is required for efficient activation of proMMP-2 in vivo. *J Biol Chem* 2000;275:26411–26415. [PubMed: 10827175]
- Ward RV, Atkinson SJ, Reynolds JJ, Murphy G. Cell Surface -mediated activation of progelatinase A: Demonstration of the involvement of the C-terminal domain of progelatinase A in cell surface binding and activation of progelatinase A by primary fibroblasts. *Biochem J* 1994;304:263–269. [PubMed: 7998943]
- Weiss SJ, Peppin G, Ortiz X, Ragsdale C, Test ST. Oxidative autoactivation of latent collagenase by human neutrophils. *Science* 1985;227:747–9. [PubMed: 2982211]
- Wilson CL, Heppner KJ, Labosky PA, Hogan BLM, Matrisian LM. Intestinal tumorigenesis is suppressed in mice lacking the metalloproteinase matrilysin. *Proc Natl Acad Sci USA* 1997;94:1402–1407. [PubMed: 9037065]
- Wilson CL, Ouellette AJ, Satchell DP, Ayabe T, López-Boado YS, Stratman JL, Hultgren SJ, Matrisian LM, Parks WC. Regulation of intestinal α -defensin activation by the metalloproteinase matrilysin in innate host defense. *Science* 1999;286:113–117. [PubMed: 10506557]
- Winberg JO, Berg E, Kolset SO, Uhlin-Hansen L. Calcium-induced activation and truncation of promatrix metalloproteinase-9 linked to the core protein of chondroitin sulfate proteoglycans. *Eur J Biochem* 2003;270:3996–4007. [PubMed: 14511382]
- Winberg JO, Kolset SO, Berg E, Uhlin-Hansen L. Macrophages secrete matrix metalloproteinase 9 covalently linked to the core protein of chondroitin sulphate proteoglycans. *J Mol Biol* 2000;304:669–680. [PubMed: 11099388]
- Woessner JF. Matrix metalloproteinases and their inhibitors in connective tissue remodeling. *FASEB J* 1991;5:2145–2154. [PubMed: 1850705]
- Woessner, JF, Jr. The matrix metalloproteinase family. In: Parks, WC.; Mecham, RP., editors. *Matrix Metalloproteinases*. Academic Press, Inc; San Diego: 1998. p. 1-14.
- Worley JR, Thompkins PB, Lee MH, Hutton M, Soloway P, Edwards DR, Murphy G, Knauper V. Sequence motifs of tissue inhibitor of metalloproteinases 2 (TIMP-2) determining progelatinase A (proMMP-2) binding and activation by membrane-type metalloproteinase 1 (MT1-MMP). *Biochem J* 2003;372:799–809. [PubMed: 12630911]
- Yamamoto K, Higashi S, Kioi M, Tsunazumi J, Honke K, Miyazaki K. Binding of active matrilysin to cell surface cholesterol sulfate is essential for its membrane-associated proteolytic action and induction of homotypic cell adhesion. *J Biol Chem* 2006;281:9170–9180. [PubMed: 16476739]
- Yu Q, Stamenkovic I. Cell surface-localized matrix metalloproteinase-9 proteolytically activates TGF- β and promotes tumor invasion and angiogenesis. *Genes Dev* 2000;14:163–176. [PubMed: 10652271]
- Yu WH, Woessner JF Jr. Heparan sulfate proteoglycans as extracellular docking molecules for matrilysin (matrix metalloproteinase 7). *J Biol Chem* 2000;275:4183–4191. [PubMed: 10660581]
- Yu WH, Woessner JF Jr, McNeish JD, Stamenkovic I. CD44 anchors the assembly of matrilysin/MMP-7 with heparin-binding epidermal growth factor precursor and ErbB4 and regulates female reproductive organ remodeling. *Genes Dev* 2002;16:307–23. [PubMed: 11825873]
- Zhou Z, Apte SS, Soininen R, Cao R, Baaklini GY, Rauser RW, Wang J, Cao Y, Tryggvason K. Impaired endochondral ossification and angiogenesis in mice deficient in membrane-type matrix metalloproteinase I. *Proc Natl Acad Sci USA* 2000;97:4052–7. [PubMed: 10737763]

Fig. 1. Domain Structure of Mammalian MMPs

Although MMPs are often subdivided into groups based on differences in domain composition, there is little consensus in the field as to how such subdivisions should be assigned. Domain structure alone does not predict function nor substrate preference. However, two clear divisions can be made: 1) MMPs that are secreted or anchored *in cis* to the cell surface; and 2) MMPs that contain a furin-recognition motif and those that do not. Nine MMPs, including all the membrane anchored enzymes, have a furin-recognition domain; how the others (17 in human; 18 in mice) are activated is the focus this review. The anchored MMPs link to the cell surface via a transmembrane domain (TM; MMP-14, -15, -16, and -24), a GPI linkage (MMP-17 and -25), or an N-terminal signal anchor (SA; MMP-23). As discussed in the text, the so-called “secreted” MMPs may still be confined to the cell surface via interactions with specific adaptor macromolecules.

Fig. 2. Mechanisms of ProMMP Activation

Latency of MMPs (i.e., proMMPs) is maintained by an electrostatic interaction between the free thiol of a conserved cysteine in the prodomain with the HIS-ligated zinc atom in the catalytic pocket. In this state, the prodomain covers the catalytic cleft thereby barring an interaction with a protein substrate. Proteolytic cleavage of the prodomain by furin or other proteases removes the thiol constraint. The thiol-Zn interaction can also be disrupted by non-proteolytic means, and in the lab, this is easily with organomurcials, such as APMA, or SDS. In cells and tissues, proMMPs can anchored to other macromolecules, such as integrins and proteoglycans, and these interactions can lead to allosteric disruption of the thiol-Zn bond. Even though the pro-domain may not be cleaved, the MMP will be active. However, final activation of MMP seems to involve prodomain cleavage, which following an conformational perturbation of the cysteine switch, seems to be controlled by autolysis.

Fig. 3. Control of MMP Activation and Activity by Adaptors

In vitro, proteolysis requires only a proteinase and a substrate, but *in vivo*, at least one additional component is typically included to augment, if not define, specificity, as well as perhaps catalytic rate. These accessory factors could (and do) come in various flavors: protein or glycosaminoglycans, membrane-associated or extracellular matrix. As suggested by this diagram, a transmembrane accessory factor would simultaneously interact with the proteinase and substrate, bringing both together at an effective concentration. In addition, the activation of some proMMPs, autolytic or otherwise, may be mediated by interaction with these factors, among other potential functions.

Table 1**Mammalian Matrix Metalloproteinases**

Designation*	Common Name	Other Name(s)	Activation Mechanism
1. MMP-1	Collagenase-1	Fibroblast collagenase, Interstitial collagenase	Unknown
	Mcol-A, Mcol-B [‡]		Unknown
3. MMP-2	Gelatinase-A	72-kD gelatinase, 72-kD type IV collagenase	MT1-MMP/TIMP-2
4. MMP-3	Stromelysin-1	Transin-1	Unknown
5. MMP-7	Matrilysin	PUMP	Unknown
6. MMP-8	Collagenase-2	Neutrophil collagenase	Unknown
7. MMP-9	Gelatinase-B	92-kD gelatinase, 92-kD type IV collagenase	Unknown
8. MMP-10	Stromelysin-2	Transin-2	Unknown
9. MMP-11	Stromelysin-3		Furin
10. MMP-12	Metalloelastase		Unknown
11. MMP-13	Collagenase-3	Rat collagenase	MT1-MMP/TIMP-2
12. MMP-14	MT1-MMP	Membrane-type MMP	Furin
13. MMP-15	MT2-MMP		Furin
14. MMP-16	MT3-MMP		Furin
15. MMP-17	MT4-MMP		Furin
16. MMP-19	RASI-1		Unknown
17. MMP-20	Enamelysin		Unknown
18. MMP-21			Unknown
19. MMP-22			Unknown
20. MMP-23	CA-MMP		Furin
21. MMP-24	MT5-MMP		Furin
22. MMP-25	Leukolysin	MT6-MMP	Furin
23. MMP-26	Endometase	Matrilysin-2	Unknown
24. MMP-27 [§]			Unknown
25. MMP-28	Epilysin		Furin

* MMP-4, -5, and -6 turned out to be either MMP-2 or MMP-3 and, hence, were not unique MMPs. MMP-18 (collagenase-4) has only been cloned from *Xenopus*. A mammalian homologue has not been found.

[‡] Mcol-A and -B are likely the murine homologues of MMP-1 (Balbin et al., 2001). Based on its chromosome position and enzymatic activity, Mcol-A is a strong candidate to be the murine orthologue of human MMP-1.

[§] In addition to being identified in a large scale bioinformatic search of secreted proteins (Clark et al., 2003), full-length cDNAs for human, rat, and northern tree shrew MMP-27 have been submitted to GenBank (access no. AF195192, XM_235795, AF281673, respectively).