Lectures

ON CERTAIN VIEWS ON THE NATURE AND TREATMENT OF PHTHISIS BY PHYSICIANS.

DELIVERED AT THE

BROMPTON HOSPITAL FOR CONSUMPTION AND DISEASES OF THE CHEST, IN THE SUMMER OF 1856.

By EDWARD SMITH, M.D., LL.B., B.A., Licentiate of the Royal College of Physicians; Assistant-Physician to the Hospital, etc.

LECTURE I (concluded).

RESEARCHES AS TO THE QUANTITY OF THE INGESTA AND EGESTA, SHEWING THAT PHTHISIS IS NOT ESSENTIALLY DUE TO DEFICIENCY OF FOOD, DEFECT IN DIGESTION, OR WASTE OF FOOD.

We will now proceed to consider, with some detail, the state of the general system which is co-ordinate with the attack of phthisis, at least at or about the period of origin, and during a large part of the progress of the disease. This may be most conveniently effected by discussing the subjects which relate to the two classes of ingesta, viz., the food and the air; for upon these two processes mainly the health of the system must be dependent. The former will occupy the remaining part of this lecture, and the latter the next lecture.

Food. The appetite does not in general differ materially from that of health; it is said to be somewhat increased, but I have not found in the majority of patients any state which would warrant me in affirming that there is any decisive change induced by the disease. This will of course imply both that the enjoyment of food continues, and also that the quantity of the food taken is, in ordinary language, sufficient. I cannot adduce any other facts to corroborate this inference respecting the degree of enjoyment (and none are needed); but I can prove that the quantity of food actually taken by phthisical patients is equal to the healthy standard. (See *British and Foreign Medico-Chirurgical Review* for April 1856, with diagrams in the number for July 1856.) During the past year, I prosecuted an inquiry into this subject on fifteen male in-patients of the hospital, and weighed with great accuracy, I believe, all the fluid and solid food which they took during a whole month. The patients were of different ages, and in various stages of the disease; and a few of them were not only in the advanced stage, but had the general system much emaciated and enfeebled. The returns thus obtained are very voluminous, but a simple statement of the average results will suffice.

The subjoined table shows the average daily quantity (in ounces avoirdupois) of solid food, and of meat and bread; also the average daily quantity of fluid, and of ale and porter, used by fourteen patients for a month during May and June; also the average daily weight (in ounces) of undiseccated faeces passed by them during the same period. The cases are arranged in the order of quantity of solids in the urine, beginning with the highest.

No. of case.	Solid food.	Meat and fish.	Bread.	Total fluids.	Ale and porter.	Fæces.
69	40.1	6.8	21.7	103.0	15.1	7.0
95	32.3	6.2	18.4	72.2	8.2	7.5
107	21.1	5.2	9.1	83.3	14.7	7.8
88	22.3	5.4	8.7	53.0	19.0	2.41
79	30.1	4.1	14.0	63.3	8.7	9.6
75	21.6	4.3	9.1	60.0	8.0	5.65
46	26.6	3.8	15.6	67.0	10.0	5.21
80	28.8	4.7	14.0	62.6	7.4	4.75
51	14.7	5.0	6.8	92.8	8.0	5.85
90	21.7	4.2	8.6	43.6	2.4	6.14
58	31.4	5.2	16.6	82.6	8.0	7.55
73	15.2	3.1	8.4	62.0	9.1	2.6
85	19.7	3.7	7.1	54.8	8.0	4.82
77	19.0	4.0	8.2	51.3	10.0	2.34

It thus appears that, on the total average of fourteen patients taken for the period of a month in May and June, the quantity of solid food per day was 24½ ounces, and that of this quantity were 4½ ounces of cooked meat without bone, and ¾ pound of

bread. These average quantities differed much in the several cases, as the quantity of food in health differs, according to the stature and other peculiarities of each person, and were the least in the most advanced cases of phthisis. The highest average occurred in a soldier, 6 feet 1½ inches high and 13½ stones in weight, and was to the extent of 40 ounces per day, and of that nearly 7 ounces were cooked meat, and nearly 22 ounces were bread. The lowest average was in an advanced and shattered case, of short stature and light weight, and was 19 ounces per day, of which were 4 ounces of cooked meat, and 8½ ounces of bread. The average of the seven less advanced cases was 27½ ounces of solid food per day, including 5.1 ounces of cooked meat, and 15½ ounces of bread; whilst, in the seven more advanced cases, the average quantity was 21½ ounces of solids, including 3.6 ounces of meat and 10 ounces of bread.

In estimating the value of these figures, we must not forget that, even in the least advanced case, the duration of the disease had been for a long period, and the amount of deposit or destruction, or both, considerable; and also that all the cases had become emaciated to a greater or a less extent. In referring to the results of these experiments, I therefore take the strongest argument against myself; for, as my aim is to prove whatever may be causative of phthisis, I at least go far enough when I take cases of the earliest deposition, and too far when the disease, by long continuance, has reacted upon the system. If, therefore, in the latter case, it can be shown that the quantity of food taken into the system is equal to that in health, *à fortiori*, the argument would be strengthened if we could refer to conditions existing at an earlier period of the disease.

Now, if we refer to our own experience, or to the dietaries of public institutions, we cannot doubt that the quantities of solid food thus given, on the averages, in early phthisis, are equivalent to that which is taken in health. But, since the appetite, whether in health or disease, varies much in the long period of a month, and since in phthisis it is hardly possible that a whole month should pass without some complication having appeared which would doubtless modify the appetite, we may gain a truer representation of the amount of food taken by considering the daily returns of the cases. This I cannot do in detail; but I shall refer to the case of the Life Guardsman, No. 69, who took the largest amount of food. From May 15 to 31, the largest quantity on any one day was 49 ounces, and the smallest 40½ ounces; and, on the average, the quantity was 45 ounces, or nearly 3 pounds in weight, against 40 ounces on the average of the month. After May 31st, his appetite failed from an inflammatory complication; so that, on one day, he took no food whatever, and on the remaining days until June 11th, the quantity varied from 20 ounces to 38½ ounces. In this case, the quantity of meat per day, from May 15th to 31st, varied from 6 to 11 ounces, and subsequently from nothing to 9 ounces; whilst that of bread varied in the former period from 22 to 30½ ounces; and in the latter period, from nothing to 20 ounces.

This case forms one end of the table; but if we take one at the other end, which was under observation for a month, and was in a most shattered condition, we shall find great variation from day to day. Thus, in case No. 85, the quantity of solid food varied per day from 13 ounces to 23½ ounces, and of meat from 2½ ounces to 6 ounces, and of bread from 5 to 10 ounces.

A careful perusal of the tables in each case cannot fail to teach that, in uncomplicated phthisis, in the earlier stage, the quantity of solid food is equal to that taken in health.

Now, if we turn from such facts as those now given to the impression which is left upon our mind by careful inquiry from numbers of patients who come under our observation, the same result is obtained. It is my practice amongst the out-patients to note the state of the appetite; and I affirm, with every conviction of truth, that in but a small minority of the cases is the appetite not good, or the quantity of food taken or desired less than their ordinary standard. That, in some instances, the appetite should be insufficient or unsteady, either for a longer or a shorter period, would not be by any means peculiar to disease, much less to phthisis, and has no weight in opposition to the general results.

The quantity of fluid food which was taken in the long inquiry before referred to is not so important to our subject, since that might depend upon other conditions than a desire for nutrition. It consisted, however, in part, of nutritive materials, such as milk, beer, wine, and beef-tea; but the larger quantity was of tea, coffee, and water. It may, however, render the argument less incomplete, if we refer to it; and it may also be of interest, if considered without much detail. (See Table.)

On the total average, the quantity of fluid per day in each case was 70 ounces, or 3½ imperial pints, which quantity in-

cluded half an ounce of wine and 10 ounces of ale and porter. It was the least in the most advanced, and the greatest in the least advanced cases; and when the fourteen cases are divided into these two classes of equal numbers, the quantity is 64 ounces and 72 ounces. The case which took the largest quantity of solid food took also the largest quantity of liquids; and in that the latter was, on the average, 103 ounces, or more than 5 imperial pints. The least average was found in one of the advanced cases, No. 90, and was 43½ ounces. There was much variation in this quantity from day to day. In the cases which took the largest quantity, it varied from 66 ounces to 127 ounces; and in that which took the least, the variation was from 34½ ounces to 54 ounces.

Thus, on the average, the proportion which the fluids bore to the solids in the food was somewhat less than as 3 to 1, and it varied in the different cases from as 2 to 1 to as 4 to 1. The total quantity of ingesta on the whole average was 93 ounces, or nearly 6 avoirdupois pounds weight per day. In case No. 69, it was as high as 143 ounces, or 9 pounds in weight; and in the case which took the least food, it was 70½ ounces, or one-half of that taken by case 69. The total average in the seven cases the least advanced was 99½ ounces, and in the remaining seven cases it was 85½ ounces. Case 69 took 170½ and 172½ ounces of ingesta in two succeeding days, or a total of 21½ pounds by weight; whilst the largest quantity taken in one day by the most advanced case was 90 ounces, only one-quarter of which was solid.

It is not my present purpose to show the effect which phthisis has in its ultimate progress in lessening the quantity of the ingesta; but it is evident from the above mentioned facts, that the quantity of both solid and fluid food is less in advanced than in early cases. It is, however, clear that, in every stage of the disease, upon the average, there is no inconsiderable quantity of ingesta, especially of a nitrogenous kind; and that the diminution in quantity is less than the wasting and exhaustion of system would lead us to expect. In the early stages, and for a long period after the first deposition of tubercle, there is no lessening of the quantity or the enjoyment of food; and, as we cannot prove the state of the appetite at the moment of first deposition, since that moment is never known, we are left to the inference, that, as in health, viz., before the first deposit of tubercle, the appetite was good; and, as it was so long after the deposit, it must have been the same at the moment of the deposition. I would, therefore, affirm the rule, that phthisis is not necessarily dependent upon loss of appetite, nor upon an insufficient quantity of food, where it can be obtained.

Having, then, established this fact, the next step will be to consider if the food thus abundantly introduced into the system is duly digested and prepared for the sustentation of the body. Now, the fact of the appetite being continuously good, and the quantity of food being normal, would, *a priori*, show that the food must be duly disposed of according to the usual course observed by the system. Our own experience, as that of our patients, leads to the belief that, wherever the organs concerned in digestion become deranged in their action, we suffer a diminution or loss of appetite, and consequently greatly diminish the quantity of solid ingesta. Then, again, when, from overfeeding or lessened wear and tear of the system, the body experiences repletion, the appetite and quantity of food commonly lessen. Both of these results are quite in keeping with what has long been proved to be a law of nature in the system, viz., that, whenever she cannot use or has no need, she ceases to desire. It is quite true that, in one or two diseases, this law may appear to be set at nought; but, upon the whole, the rule would be universally acknowledged. Hence, on this ground, we should be prepared to believe that the food taken into the system in phthisis is duly digested. If it were not so, it would depend either upon an altered action, and therefore an altered state of the organs of digestion, or upon its being hurried too rapidly through the alimentary canal.

If it depended upon derangement of the digestive organs, it would certainly evince itself by the well known signs of indigestion. We have already inferred that no such state of things exists, by the fact of the continuance of the appetite and supply of food, and we get abundant confirmation of this in our medical practice. I am of opinion that in no stage of the disease is dyspepsia more commonly observed in phthisis than in any other chronic disease of vital organs; but on the contrary, that it is much less marked than in long continued disease of the liver. Much less is it a necessary concomitant of the early stage of phthisis. As a rule, which may certainly be proved to exist amongst the out-patients, the tongue is clean and is remarkable only from a deficiency of blood in its structure, and there is

neither tenderness at the epigastrium on pressure, nor pain or inconvenience after meals. Of course, as dyspepsia is the most frequent disease affecting humanity, and the most frequent in hospital practice, it is not unfrequent amongst phthical patients, who constitute so large a proportion of all cases of disease; but in them as in others, it is only temporary, and is as amenable to treatment as it is in persons otherwise healthy. Pressure over one or both lobes of the liver will not unfrequently give a little pain, and indicate the existence of a condition which otherwise would be most readily referred to the lungs themselves, but which a little blue pill and the ordinary treatment for inactive or congested liver speedily removes. Then, again, I would affirm that in the early stages of phthisis, dyspepsia is not by any means a constant concomitant, and consequently that the latter is not even a necessary effect of, much less can it be a common cause of the former. But all this rather refers to the effect than to the cause; yet if dyspepsia were a common cause, it would surely commonly attend the early stages of the disease. But regarding it at an anterior period and as a cause, I do not think that there is any evidence to show that phthical persons were especially subject to dyspepsia before the period of earliest deposit, or that dyspeptic persons are peculiarly prone to phthisis. Hence, whilst dyspepsia may occur to phthical patients either before or after the deposit, it is only to the extent in which it attacks persons otherwise healthy, and is not necessarily or commonly causative of phthisis; and so far it is fair to infer that in phthical cases, as in health, the abundant food is properly digested.

The other alternative of want of due digestion, viz., the *hurrying* of the food again from the system, can be due only to the acts of vomiting and purging. Vomiting is not by any means common in the early stage of phthisis, nor indeed in any stage of the disease; and we know that it is not frequent in health, or in that condition which immediately precedes the earliest deposition of tubercle. In the later stage of the disease it is sometimes found, and may then in part account for the wasting; but it is then rather the effect of a general irritability attending upon an exhausted system, and of a distended stomach impeding the very limited action of the lungs. In the investigation of 15 patients for a month, only two during that period had vomiting. In one it was considerable, and produced on the average of the month 1·82 ounces per day. This case exhibited great nervous excitability in his temperament, and in the large quantity of colourless urine of very low specific gravity, and he subsequently died of brain disease. He had been intemperate. The other had vomiting to the extent of an average of one-fifth of an ounce per day, and at the same time had diarrhoea and albuminuria. The former was an advanced, the latter an earlier stage of the disease. In the former there were at the same time only 21 ounces of solid food taken per day, although the system had not greatly suffered; a circumstance probably attending upon or due to the vomiting. So also purging is not common in health, neither is it frequent nor peculiar to the early stage of phthisis. I do not know that in the early stage of phthisis there is much change in the frequency of the alvine evacuations; but if there be, it is rather on the side of a deficiency of action. Hence by neither mode is the food once taken into the system unfairly hurried out of it.

But there is yet a third source of evidence in reference to this perfect or imperfect digestion of the food to which reference should be made, viz., the amount of nutritive matter still left in the faeces when they have passed through the alimentary tract. It would not be difficult to determine this fact with accuracy, but my present inquiries have stopped short at this point. In the inquiry to which reference has been made, the quantity of most of the egesta as well as the ingesta was determined; and, so far as that may be a guide, it has been ascertained with a fair amount of accuracy. (See Table.)

On the total average of all the cases on each day for a month, the weight of the faeces was 6·3 ounces avoirdupois, and varied from 2·34 ounces to 9·6 ounces per day; whilst in health, on Lehmann's authority, the faeces weigh from 5 ounces to 7½ ounces. In the seven most advanced cases it was 4·9 ounces, and in the seven least advanced cases 6·4, or a difference of 1½ ounce per day on that average. Thus, in reference to the stage of the disease, it agrees with the ingesta, but the proportion of the faeces to the whole ingesta is even less in the advanced than in the early stage, viz., 1 in 17½ in the former, and 1 in 15½ in the latter; and the proportion of the faeces to the solids only in the ingesta is precisely the same in both classes, viz. 1 in 4·3. So far, then, as quantity is evidence of nutritive quality in the faeces, it is clear that there is no greater waste in the advanced than in the early stage of the disease.

As the feces in these experiments were not desiccated before they were weighed, they may have been more or less solid, and would thus vary in weight with the occurrence of diarrhoea or with the frequency of the stools. Hence there would not necessarily be a constant correspondence between the weight of the feces and of the whole ingesta, or of the solid or fluid parts of the ingesta taken separately. The case (No. 69) which took the largest quantity of both solid and fluid food did not have the greatest weight of feces, but the greatest weight of feces was found in a case in the early stage (No. 79), in which there were the complications of diarrhoea and excessive albuminuria.

The case which appeared to closely resemble a state of health was a ruddy looking farmer (No. 95), who had one lung sound. On the total average, the weight of his feces per day was $7\frac{1}{2}$ ounces, whilst that of the total ingesta was $105\frac{1}{2}$; so that the former was one-fourteenth of the latter, and as 1 to 4.2 of the solids in the ingesta. This is somewhat more than the proportion which the total average of the cases gives, and consequently shows that as we approach to a state of health the quantity of feces increases and not diminishes.

It has been stated that when the stools were numerous, the total weight of the feces was increased. This was seen in case 79, in whom it amounted to 20 ounces on one day, and $16\frac{1}{2}$ on another day. In case 69 also to 19 ounces in one day, and in case 95 (the healthy looking man) to 19, 19, and $21\frac{1}{2}$ ounces on succeeding days. The same fact, however, occurs in some instances where there is deficient action, or when an evacuation occurs only once in two or three days. The largest amount passed in any of the cases was in one of very sluggish temperament, who, after passing three days without an evacuation, had one to the extent of 32 ounces, and again another of 20 ounces at a further interval of two days. In this case there was a considerable quantity of food taken, viz., $31\frac{1}{2}$ ounce average. There was, however, an instance of very costive habit, in which, after an interval of two or three days, the evacuation varied only from $5\frac{1}{4}$ to $8\frac{3}{4}$ ounces, and produced on an average of a month only 2.41 ounces per day. In this instance, the average quantity of solid food was $22\frac{1}{2}$ ounces. Thus it is evident that it would be fruitless to expect the feces to bear the same proportion to the ingesta from day to day, and there is only a very general proportion between them in the long interval of a month.

From the foregoing statements, we learn that, in the cases nearest to health, the quantity of feces was above the average of the whole cases combined, and was greater in proportion to the total ingesta than in the more advanced cases; and also that, as the disease advanced, the proportion of the feces became less. Moreover, as, on the total average, the weight of the feces in phthisis is much greater than in health, there are some grounds for the inference, that the food does not in that disease pass off in the feces, undigested and unappropriated by the system, at least not to a greater extent than probably occurs in health—a probability infinitely greater when we limit the attention to the condition precedent of the deposition of tubercle. Hence we have not as yet any evidence by which to ascribe the origin of phthisis to the absence of abundant nutriment in the body.

Now, if, in addition to the previously mentioned facts, we add the not insignificant one, that phthisis occurs in all classes of the community, and therefore, in a very large proportion of instances, in persons who must, throughout their whole period of existence, have had abundant and proper nourishment, we must acquit the whole of the organs of digestion, as well as the quality and quantity of food, of any necessary share in the production of phthisis, and, for all purposes of practice, may exclude this subject from our attention.

[The researches into the quantity and chemical constituents of the urine were not completed when the above lecture was delivered; hence they are not inserted in this place.]

ALLEGED CURE FOR SEA SICKNESS. Dr. Landerer, at Athens, announces that he has discovered a sovereign specific against sea sickness. His remedy is to give from ten to twelve drops of chloroform in water. The chloroform in most cases removes nausea, and persons who have taken the remedy soon become able to stand up, and get accustomed to the movements of the vessel. Should the sickness return, a fresh dose is to be taken. It was tried on twenty passengers on a very rough voyage from Zea to Athens, and all, with the exception of two, were cured by one dose. The minority, two ladies, were able to resist the feeling of illness on taking a second dose. (*Galvani's Messenger*.)

Reviews and Notices.

A FEW REMARKS ON THE PERFORATING ULCER OF THE STOMACH AND BOWELS. By J. BOWER HARRISON, M.D., F.R.C.S., etc. pp. 68. London: Churchill. 1856.

In this little *brochure*, Dr. BOWER HARRISON presents us with some very sensible observations, written in the same easy conversational strain which characterises all his other medical productions. After narrating several very interesting cases, he discusses in succession the diagnosis, treatment, and morbid anatomy of the fatal lesion under consideration; scattering here and there, as he proceeds, a short but highly useful practical remark. For example, he very truly observes that—

“Many cases are recorded with great minuteness which still omit points of the utmost importance in diagnosis. In order to report a case well, we must have some previous knowledge of what is chiefly worthy of observation.” (pp. 9, 10.)

The first of Dr. Harrison's cases is an interesting illustration of the practical acumen of the late Mr. Robert Thorpe, of whom Sir James Bardsley so justly observed at the Manchester anniversary meeting, that his knowledge far exceeded his power of expression. The case was that of a female presenting the ordinary symptoms of strangulated hernia, with “a small tumour in the bend of the right thigh, as nearly as possible in the situation of the crural ring.” The woman was conveyed to the Manchester Infirmary, where every one thought the case to be one of hernia, with the exception of Mr. Thorpe. After death, the tumour was found to be merely a softened gland; and there was peritonitis from perforating ulcers of the small intestine.

In another case of fatal perforation occurring in a gentleman who, “though only thirty-seven years of age, had a much older look, and might easily have been taken for fifty”, the immediate occasion of the fatal rupture of the stomach by perforating ulcer, was an overloaded condition of the organ. The patient “was a man of very abstemious habits, and accustomed to long fasts; but on the morning of the attack he had made a hearty breakfast, and partaken of mutton chops.” The author also remarks that—

“Many cases are recorded in which a hearty meal has been the precursor of death by perforation; and others are mentioned in which an emetic has been the occasion of a fatal rupture.” (pp. 42, 3.)

As signs of perforation having taken place, it is stated that—

“The pain which ensues rarely or never abates, but becomes more and more severe, until death occurs. Exceptions have indeed been mentioned, but they are certainly rare. The pulse is not always affected at the commencement, and sometimes it remains steady for many hours, and fails only with the approaching collapse. The intellectual faculties are unfortunately too clear, and remain unimpaired to the end. I do not think that there is any peculiarity in the tongue, nor should we expect any. The abdomen mostly becomes tympanitic after perforation has taken place, though, before that occurrence, it is often hard and contracted. If the patient vomit, he does not vomit much; it is an abortive effort, attended with aggravation of pain and flatulence. If fluids be taken—at any rate, when the perforation is in the stomach—there is a dreadful accession of suffering after a brief interval.

“The pain which attends perforation is sudden and violent. It is violent at once, and does not become so by degrees. This in itself is striking, and unlike the more gradual invasion of ordinary inflammation. Again, it is more than commonly violent. The patient says he cannot bear it. He speaks authoritatively, and says he must have relief, or he shall die.” (pp. 22-4.)

Speaking of the liability of anæmic young women to this disease, the author observes:

“It would appear that the changes in the molecular constitution of the stomach are more rapid than those of other parts of the body, and that it is the first to suffer when aliment is