GUY'S HOSPITAL.

TREATMENT OF OPEN CANCER BY THE APPLICATION OF CHLORIDE OF ZINC.

Under the care of J. BIRKETT, Esq.

[From Notes by T. BRYANT, Esq., Surgical Registrar.]

We have been favoured with notes of two cases in which the above escharotic was used in ulcerating cancer of the breast—the only two which have been so treated in Guy's Hospital, at least for a considerable time. We record these cases as bearing on the question which has of late acquired such interest, relative to the value of different local applications, and as a continuation of those which we have already published of a series which will embrace all the cases which we can discover to have occurred in the London hospitals. Such a series will furnish some (though we fear, from its limited extent, not very perfect) data for comparison with the results of the treatment of Dr. Fell and other plans for the cure of cancer, which will shortly be published. The fault of such cases is that they comprise only the least important of the points of treatment, viz., the local action of the application. The far more important question of the progress of the constitutional disease, and the respite obtained for the patient, is not settled by them, nor under the present hospital system do we see how (unless by accident) particulars bearing thereon can be obtained. Our readers must be content to accept from us all that our opportunities enable us to offer.

These cases, being necessarily somewhat protracted, cannot be reported quickly. We hope, however, to continue the series regularly, and to sum up its results at some future period. All that can be said of the two patients referred to below is, that the treatment appears to have temporarily succeeded in the first case, and to have been less successful in the second. It should be added, that the chloride was applied in powder mixed up with starch.

CASE I. E. Bland, a female, aged 55, was admitted with a small carcinomatous tubercle (of the size of a walnut) of the skin over the left breast, of nine months growth, which had been discharging two weeks. On March 6th, 1855, the chloride of zinc was applied, and repeated on April 5th. On the 24th the whole tumour appeared to have sloughed away, and on May 6th she left with the skin cicatrising.

CASE II. M. A. Morris, aged 52, was admitted into Guy's Hospital, under Mr. Birkett, on Oct. 29th, 1856, with open carcinomatous ulcerating tubercles over left breast, of five years and a half duration. On November 3rd, the chloride of zinc was applied; the application was followed by acute pain. On the 16th, the slough came away; and on the 17th, the chloride was again applied, from which a large portion of the tumour was discharged. On November 23rd, she left relieved.

ST. MARY'S HOSPITAL.

CASE OF DISEASE OF THE AORTA : ANEURISMS AT THE AORTIC SINUSES : CLOTS IN THE ARTERIA INNOMINATA AND LEFT MIDDLE CEREBRAL ARTERY.

By W. O. MARKHAM, M.D., F.R.C.P., Assistant-Physician to the Hospital.

A FEMALE, aged 50, the subject of the following history, came into St. Mary's Hospital on the 4th instant. She walked into the Hospital, and gave the following account of her previous condition; thereby showing, that at this time there existed no paralysis either of sensation or of motion. She stated, that when very young she had suffered from some severe affection of the chest, and had ever since from time to time been in an ailing condition. The chief symptoms which troubled her were palpitations, cough, and a feeling of constriction of the chest. From the circumstance that old organised adhesions of the pericardial surfaces were found after death, and from the nature of the symptoms here mentioned, we may presume that this attack was one of pericarditis. Latterly these symptoms had increased in severity, and had incapacitated her for performing any hard work; in addition, she had also frequently had attacks of faintness and giddiness, and particularly in cold weather and on exertion. Occasionally, indeed, when out walking in the street she had been forced to cling to the walls or railings for support; and on account of her staggering gait she had been several times unjustly accused of being drunk. These symptoms increasing in severity, and her breathing in particular being very difficult, she sought relief in the Hospital.

On her admission, her lips were blue, her breathing very hurried, and the veins of her neck were much swollen. It was also remarked that the pulse at the right wrist was imperceptible. At this time she was perfectly sensible, and had perfect command over her muscles. She took her dinner, and went to bed; and in about an hour afterwards was found lying low in the bed on her back and insensible. Her right arm and leg were completely paralysed, presenting not the smallest rigidity when raised or moved. The pupils of both eyes were firmly contracted and immovable. The pulse at the right wrist was not perceptible, neither was any pulse to be felt in the arteries on the right side of the head and neck; I fancied, however, that I felt a very slight pulsation in the upper part of the right brachial artery. Occasionally, during her heavy breathing, the right cheek was puffed out.

Loud râles were heard over the front of the thorax; the breathing was very noisy, so that it was not easy to make a very careful stethoscopic examination of the heart's sounds. I satisfied myself, however, of the existence of a diastolic aortic murmur—a bruit, coincident with the diastole of the heart, being audible from the middle of the sternum down to its lower edge.*

The arrest of the circulation through the vessels supplied by the arteria innominata, the diseased condition of the aortic valves indicated by the bruit, and the suddenness of the paralysis caused me at this time to hazard the conjecture that a clot of fibrin had suddenly arrested the circulation through some one of the cerebral arteries—a conjecture which was fully justified subsequently.

The difficulty of breathing gradually increased, the paralysis and other symptoms remaining the same, so that the patient died about sixty hours after the first occurrence of the paralysis.

Post mortem examination. When the body was examined, the limbs of both sides were found of equal rigidity; there was no œdema, and but little fat on the surface; the superficial tissues were remarkably dry and bloodless. In both pleurae there was a considerable quantity of serous effusion; the lower lobe of the left lung was thereby compressed and rendered nearly solid; the right lung was adherent at its apex, which contained a few old, small, tubercular nodules. The lungs were highly congested with bloody serum. The surfaces of the pericardium were everywhere firmly adherent by old and organised attachments. The roots of the aorta and pulmonary artery were matted together; and so likewise were the auricles. The heart was slightly dilated and hypertrophied; its aortic valves were thickened and slightly shrunken, and thereby rendered incompetent; water, when poured into the aorta, escaped freely through them into the left ventricle. The other valves were quite healthy. The arch of the aorta was dilated, and its internal surface much diseased, exhibiting everywhere signs of fibroid and atheromatous degenerations. In the aortic sinuses were found the openings of three small aneurisms. One, situated above the left coronary valve, admitted the little finger, and led into a pouch which passed behind the pulmonary artery towards the appendix of the left auricle. The openings of the two other aneurisms were at the right of the aorta, and situated above the right coronary valve; one was of the size of a filbert; the other was much larger, and pressed considerably upon the descending cava and the right auricle; the descending cava would not admit the entrance of the little finger.

The arteria innominata was found completely blocked up, for the last inch of its course, by a firm, whitish-red, hardish coagulum, strongly adherent to the walls of the artery, and extending up the whole length of the common right carotid. The opening of the left carotid artery at the aortic arch was not more than one-fourth of its natural calibre; being contracted by hard fibroid exudation into its coats. The left subclavian artery was healthy.

When the dura mater was opened, blood flowed rather freely from the venous sinuses; many red points appeared on the cut surfaces of the brain. A considerable quantity of clear serous fluid was present in the lateral sinuses of the brain. The left carotid artery, where it emerges from the sella Turcica and passes to the brain, was found completely blocked up by a firm reddish coagulum, which extended for an inch along the middle cerebral artery. The portion of the brain to which the middle

* I mention the direction of the murmur, because the fact (a fact, at least, as far as my observation goes) seems not sufficiently recognised—viz., that diastolic aortic murmurs are heard much more frequently and distinctly in a direction down along the sternum, than towards the heart's apex. The reason of this is not very clear; and the fact is evidently opposed to what might be conjectured *a priori*.

cerebral artery—the chief branch of the internal carotid—is directed was much softened, readily breaking down into a pulp, and particularly so was the left corpus striatum. The other parts of the brain, and likewise the vessels at its base, were found quite healthy in appearance.

There was nothing worthy of note to be mentioned concerning the other organs of the body.

REMARKS. This case presents some points of interest both in a physiological and in a pathological sense. It appears clear that the arteria innominata was blocked up at the time of the admission of the patient into the Hospital, and for the reasons that the pulse was then imperceptible at the wrist, and that the clot itself was evidently not of recent formation. Consequently the functions of the brain were duly carried on, notwithstanding that its supply of blood through this artery was cut off, and that not more than one-fourth of the due amount of blood could have passed through the opening of the left common carotid. Also it is evident, that for the last sixty hours of life the only supply of blood which the brain received was that which passed through the left vertebral artery. The cause of the formation of the clots in the arteria innominata, and in the left cerebral artery, must be, to a certain extent, matter of conjecture. There were no local alterations perceptible in the vessels themselves which could give any account of clots. Both vessels, to all appearance, seemed perfectly healthy; the arterial branches at the base of the skull and around the obstructed artery were quite natural to the eye. Perhaps the existence of the aneurisms may give us a clue to the obstruction of the arteries. It is not unfair to presume that it was very possible for some broken down portions of fibrin to have been washed out of one or other of the aneurisms at the root of the aorta, and to have been carried along in the current of blood until arrested in their progress, acting then as a ligature of the artery, and causing behind it the formation of a firm fibrinous coagulum. This supposition is strengthened by the fact that the mouths of the aneurisms were large. There were no fibrinous beads around any of the heart's valves, so that the carrying away and arrest of one of these in the circulation could not have caused the clots, after the manner described by Dr. Kirkes. The softening of the brain was clearly secondary to the arrest of the circulation. In this case the supply of blood to the left middle lobe of the brain must have been wholly cut off; there could have been no compensatory supply to it through the right carotid artery.

To the practitioner, also, this case of paralysis, like so many others of its kind, offers a word of fruitful warning. What would a vigorous treatment have done here? What could any interfering treatment do?

Original Communications.

THE ATROPHY OF LIMITED GROUPS OF MUSCLES.

By F. BRITTAN, M.D., Physician to the Bristol Royal Infirmary, and Lecturer on the Practice of Medicine in the Bristol Medical School.

[Read before the Bath and Bristol Branch, February 19th, 1857.]

In the *Dublin Quarterly Journal of Medical Science*, for November, 1856, Dr. Reade, of Belfast, published a paper, entitled, "Contributions to the Pathology of the Spinal Marrow." In it he relates three cases to illustrate "a diversity of morbid action arising out of disease or injury of the same portion of the spinal marrow." The three cases are: 1. "Symmetrical muscular atrophy." 2. "Paralysis of the body beneath the phrenic nerves, from concussion." 3. "Paralysis of the body beneath the phrenic nerves during dentition." The first of these cases is adduced in illustration of the views of M. Cruveilhier on muscular atrophy, announced to the Institute of France, December 1855, and published in the *Archives Générales*, January 1856. The conclusions at which M. Cruveilhier has arrived may be thus summed up:

That there is a species of muscular paralysis, sometimes partial, sometimes general, which gradually and successively invades, fasciculus by fasciculus, and fibre by fibre, the voluntary muscles, leaving all the functions of nutrition intact, excepting that of muscular nutrition.

That this gradual muscular paralysis is the consequence of

progressive atrophy of the anterior roots of the spinal nerves, and equally progressive atrophy of the corresponding muscles.

That this "*paralysie musculaire atrophique*" is analogous to that which results from section of muscular nerves.

That the facts appertaining to this form of paralysis confirm the theory of Sir Charles Bell.

And that these observations demonstrate a previously unknown influence of the anterior roots of the spinal nerves on muscular nutrition. These conclusions, M. Cruveilhier states, are founded on a great number of clinical facts; and they are preceded by a declaration so essentially French, that I cannot refrain from quoting it in the original. "Le hasard a voulu que j'aie été le premier à qui il ait été donné de faire l'anatomie pathologique de cette maladie; c'est là tout le secret de la part que j'ai prise à sa détermination, dont je ne crains pas de le dire hautement, parceque c'est la vérité; la priorité ne saurait m'être contestée."

I have lately had two cases analogous to the first in Dr. Reade's paper, under my care at the Infirmary; but as they offer certain important differences as to the history of the patients and early conditions of the disease, I think it may be worth while to publish them, with a few remarks; more especially as Dr. Reade seems to think them rare, or at all events to have been able to find but one recorded.

Dr. Reade's case is that of a young man, aged 19, nearly six feet high in his clothes, apparently a specimen of robust health and excellent constitution. Eighteen months previous to examination he had been distinguished amongst his companions in all athletic exercises, and well proportioned in his muscular development. "The first sign of the approaching disease was a degree of stiffness or difficulty of executing the motion of putting on and removing his hat" (in other words, of raising his hand to his head); never suffered pain, and has enjoyed uninterrupted health in all other respects. "When he stripped his body to the waist, he exhibited neck, chest, and arms to the elbow-joints reduced to the most abject degree of emaciation; this was symmetrical; the greater and lesser pectorals were little more dense than the strongest brown wrapping paper." The muscles of the neck, scapulae and humeri, were reduced to the mere elementary outline of muscles; whilst from the elbows, the forearms and hands displayed the full development of a robust and vigorous man, with all the concomitant power, sensibility, and aptitude for use. He was under treatment for two years. A seton in the neck, the use of dumb-bells, and electro-magnetism were tried, but without material improvement. Six years later he was again inspected, and was found decidedly improved; the muscles more developed. The muscles of the neck have been fully restored, and the others considerably augmented and perfectly obedient to the will. The forearm and hand have lost much of their former bulk; but his occupation for the six years has been sedentary. His health has been uninterruptedly good.

The analogous case is quoted from Rokitsansky. A labourer, aged 45, much exposed to wet and in the habit of allowing his clothes to dry on him, was attacked with pain in the left shoulder, which was most severe about fourteen days after it commenced. There was no swelling, redness, numbness, nor tension; but great pain when the arm was raised with the other hand: he could not lift it without. After the first week the shoulder was found wasted; as the pain subsided the wasting advanced, till the deltoid, supraspinatus, infraspinatus, and teres muscles seemed completely absorbed. The shoulder-joint was healthy, and there was no emaciation of the forearm or hand.

Dr. Reade regards this case as one of muscular atrophy from abeyance or destitution of the function of nutrition; and he states "these three cases illustrate a diversity of morbid action, arising out of disease or injury of the same portion of the spinal cord."

He also alludes to two other cases within his own cognisance; one of atrophy proceeding from the shoulders to the feet, accompanied by paralysis, and a second confirming Cruveilhier's statement of the pathology of this paralysis, the same morbid changes having been found in the roots of the anterior spinal nerves. We thus have muscular atrophy, partial or general, according to Cruveilhier, depending on atrophy of the anterior roots of the spinal nerves. Muscular atrophy partial, according to Dr. Reade, depending on imperfect nutrition, the result of disease of the spinal cord.

My two cases more accord with that quoted from Rokitsansky, and show, I think, that muscular atrophy, with of course necessarily a quarter or less degree of loss of power, proportionate to the amount of atrophy, may sometimes depend on