BRITISH MEDICAL JOURNAL:

BEING THE JOURNAL OF THE BRITISH MEDICAL ASSOCIATION.

EDITED BY ANDREW WYNTER, M.D.

No. XXIII.]

LONDON: SATURDAY, JUNE 6, 1857.

[NEW SERIES.]

Illustrations OF HOSPITAL PRACTICE:

METROPOLITAN AND PROVINCIAL.

YORK COUNTY HOSPITAL.

THREE CASES OF AMPUTATION AT THE SHOULDER-JOINT.

By W. D. HUSBAND, Esq., F.R.C.S.Eng., Surgeon
to the Hospital.

CASE I. Extensive Injury of Upper and Lower Arm: Amputation at Shoulder-Joint: Death from Subcutaneous Mortification, from severe Bruise of Hip. Mark Robinson, a guard on the North Eastern Railway, was looking out for his own train on a foggy evening (Dec. 21st, 1854), when another train, proceeding in a contrary direction, came suddenly upon him, threw him down, and then passed over him. He was found to be seriously injured, and was carried to the hospital; there I saw him within half an hour after the accident.

The patient, a stout muscular man, in vigorous health, had sustained very extensive injury of the arm, which was extensively lacerated from the shoulder to the wrist-joint. The humerus was so fractured, or rather smashed, that only the head and condyles seemed to have escaped injury. Blood had been flowing freely from several wounds communicating with the fractured bone, and was only now restrained, as no tourniquet could be applied by direct pressure over the bleeding orifices. The bones of the lower arm were also broken in several places, and the soft parts lacerated. The man complained of some pain in the hip, but had walked along the station to a fly, and up the steps of the hospital without any difficulty.

As no other alternative remained, I proceeded at once to remove the injured limb at the shoulder-joint. No chloroform was administered, at the request of the patient, who sat upright in a chair, and bore the operation with great fortitude. An attempt was made to compress the subclavian artery; but, as I agree entirely with Mr. Ferguson that the compression of the artery may be effected in an experiment on a willing subject, but can never be depended upon in the actual operation, where the pressure may be so easily removed, I relied on the ready aid of our then intelligent and efficient house-surgeon, Mr. T. C. Paley, to grasp the axillary artery as soon as it was divided in forming the inner flap. The biceps muscle was transixed opposite the joint, and made to form the outer flap. The knife was then carried across the joint, dividing the capsular ligament and tendons around the joint, and disarticulating the bone; and it was made to form an inner flap, similar in size to the outer one, while cutting its way out, somewhat in the way recommended by Lisfranc. The artery, divided in the last movement of the knife, was immediately grasped by Mr. Paley, and tied. The patient lost little blood, and was soon comfortably in bed, expressing his surprise that he felt so well after the injury and the subsequent operation.

The humerus was found to be comminuted from head to condyles, and the soft parts lacerated and almost reduced to a pulpy state.

December 22nd. The patient has passed a favourable night, under the influence of an opiate given immediately after the operation. There is evidence of a bruise over the right hip, which is somewhat painful. It was ordered to be fomented.

December 24th. The shoulder is going on well, but the hip is very painful and swollen; and the skin is becoming dark coloured, and the subjacent tissues feel "boggy", as if broken up.

December 25th. The skin over the hip looks sloughy; and,

under the somewhat hardened and ulcerating skin, considerable effusion is evidently forming in the subcutaneous tissues. The patient is restless and feeble, with a small rapid pulse and brown tongue; and the amputation wound now presents a sloughy unhealthy appearance.

December 27th. The condition of the hip has continued to deteriorate. The surface of sloughing skin is now very extensive, and there is immense effusion in the gluteal and adjacent regions. The patient is evidently sinking fast, in spite of stimulants and support, which have been freely taken since the symptoms of gangrene developed themselves. The man sank gradually during the day, and died in the evening.

The hip alone could be examined. The skin over the gluteal, femoral, and lumbar regions, was almost black; the cuticle was removed; and the true skin was in a condition of incipient gangrene. On dividing the skin, which cut somewhat like crisp leather, an immense amount of bloody serum gushed out; the subjacent muscles and tissues were found dark coloured, and reduced to a condition of thick pulp.

CASE II. Injury of Upper and Lower Arm by Flax-dressing Machine: Amputation at Shoulder-Joint: Severe Prostration: Recovery. Bridget Brown, aged 22, was admitted into the hospital, November 1st, 1855, having had her right arm drawn into and extensively injured by a flax-dressing machine at Pocklington, whence she had been brought—a distance of thirteen miles. She appeared to be a stout healthy girl, and stated her previous health to have been good. The lower arm was extensively lacerated, and the bones fractured and crushed in several places, and exposed by the loss of tegumentary covering. The humerus was not fractured, but the soft parts covering it were most extensively lacerated, and separated from the bone up to its head; and when the bandages, which had been applied tightly round the arm at the time, were removed, blood freely gushed out.

Being anxious to save some part of the upper arm, if possible, as soon as the patient was fairly under the influence of chloroform, I determined, as the force had been applied to the anterior and posterior aspect of the limb, to ascertain the condition of the soft parts at the outer and inner aspects, with a view to amputate immediately below the head of the humerus by lateral flaps; but, as soon as the muscular covering was cut into, it was found to be so completely broken up, that no sound covering could be obtained for the bone if amputation lower than the joint were attempted. I therefore at once amputated at the shoulder-joint, in the manner described in the last case; and the axillary artery was secured immediately by our present house-surgeon, Mr. Hanbury. At this moment, the patient's pulse became almost extinct, her countenance blanched, and she appeared almost lifeless. A large cold sponge was thrust into the gaping stump, and held firmly to the exposed surface, to prevent hæmorrhage; and stimulants were freely administered. She rallied a little, when the sponge was slowly removed, and any bleeding vessel secured as soon as exposed.

The girl continued in a most precarious state for many hours. Strong stimulants, including an ounce of brandy, were given every ten minutes; and the interval could not be extended to twelve minutes until eight hours after the operation, without very evident increased prostration.

Nov. 2nd. The patient continues very feeble, but the pulse is somewhat fuller, the extremities warmer, and the countenance improved. The stimulants are required less frequently.

After this date, the girl continued to improve; the stump healed slowly, but progressively; and she left the hospital in about six weeks, in her usual health.

CASE III. Scrofulous Caries of the Shaft and Internal Condyle of the Humerus: Amputation at the Shoulder-Joint: Recovery. Stephen Weldrum, aged 15, a strumous looking boy, was admitted into the hospital March 29th, 1856, with extensive disease of the upper arm. He was stated to have enjoyed fair health until, about five or six months previously, he was attacked with what was supposed to be rheumatic inflammation

of the knee and elbow. The affection of the lower limb was completely relieved, but the arm continued swollen and painful; and, about three weeks ago, an abscess broke about the middle of the upper arm, and continued to discharge freely. The arm above the elbow was much thickened from extensive bony deposit, which evidently extended from the head to the condyles of the humerus; and the skin of the entire upper arm was discoloured with tortuous veins running over it in various directions. A probe, passed into the opening through the skin, detected bone denuded, but not loose. There was also considerable deposit around and under the clavicle, and about the shoulder-joint. There was also some dulness on percussion under the clavicles. He was ordered to take twenty drops of syrup of iodide of iron three times a day, and a tablespoonful of cod-liver oil twice daily.

April 24th. Under the influence of the iodide of iron and the cod oil, the patient has acquired strength and weight; but matter has formed in several places along the humerus; and sinuses now exist in connexion with denuded bone, from the head of the bone to its condyles; and, at the lowest opening, a probe can be passed through the external condyle, apparently into the elbow-joint. A consultation with my colleagues was now held, to determine whether any and what operative proceedings should be adopted, when, on a careful examination of the chest, the pulmonary symptoms were found to be so suspicious that we determined to defer any operation at present; and he was made an out-patient, and continued the iron and oil treatment until August 30th, when, his health having improved most satisfactorily, and the chest symptoms being much relieved, it was resolved in consultation, as no minor operation offered any prospect of relief, to amputate at the joint.

September 10th. With the aid of my colleague, Mr. R. Hey, and in the presence of Professor Erichsen (who accidentally visited us on the day) and several of my medical friends, the boy being under the influence of chloroform, I removed the arm at the shoulder-joint, in the manner described in the other cases, with the only exception of having to make the outer flap partly by transfixion, and partly by cutting from without inwards, on account of the very large bony deposit extending to the head of the bone, which interfered with the free sweep of the catline. The axillary artery was dexterously secured by my colleague; very little blood was lost; and the boy recovered without a bad symptom. He is now (eight months after the operation) in good health.

The bone, cleaned after maceration, is now before me, and is nearly three times the size of a healthy humerus; it is exceedingly rough and uneven, with numerous stalactite-looking processes along its surface, and several cloacæ communicating with the original shaft of the bone, which is partly detached, and partly attached to the new case. The head of the bone, as also the condyles, are carious; and a large opening passes through the internal condyle into the fossa, which receives the olecranon process. The greatest deposit of new bone is at the point of insertion of the deltoid, and along the bicipital ridges.

REMARKS. Amputation at the shoulder-joint is by no means the serious or difficult operation which the directions of some surgical writers would lead us to imagine, if we were to be guided by the minute details given by some, and the many modes recommended by others for performing this amputation. By transfixion of the deltoid opposite the joint, an ample flap can be formed from that muscle; and the joint, if not opened then, may be readily found, the bone disarticulated by division of the ligaments and tendons, and an inner flap readily formed to correspond with the outer one. In the case No. 1, I had some slight difficulty in getting the humerus to act as a lever, and thus draw its head from the glenoid cavity, from its head being separated from the shaft; but, by pressing my finger into the joint after the partial division of the capsular ligament, I was enabled to depress the head of the humerus, and thus readily to divide the remaining capsular and tendinous attachments, and to form the inner flap. In all three cases, the axillary artery was readily seized, and very little blood was lost—less, certainly, than at ordinary amputations, with a careful adjustment of the tourniquet.

The first patient died from what all surgeons who are conversant with railway accidents have reason to dread—a severe contusion, crushing the subcutaneous tissues, while the skin, from its greater elasticity, suffered little, and hence masked the subjacent mischief, until, its nutrition having been cut off by the destruction of the subcutaneous vessels and nerves, it became involved in the general ruin. How often have we to regret the loss of a patient who comes into hospital with a crushed limb,

which is amputated, and would do well, but the blow which has knocked him from his engine, or the contusion received on some part of his body, while his limb has been under the wheel, has reduced every resisting tissue to a pulp; and the death of an extensive portion of the body is soon followed by that of the whole. I say no more on this interesting subject, as I purpose soon to supply the JOURNAL with several interesting illustrations of this class of cases.

The second case occasioned some anxiety, in the attempt, as the humerus was not fractured, to save some portion of that bone; but, even after amputation at the joint, a proper covering was obtained with difficulty, and some of that procured sloughed away in a few days. Whether the syncope during the operation arose from the slight hæmorrhage, acting upon a patient who may have lost more blood at the time than we were led to believe, or, as Dr. James Arnott would probably assert, from chloroform, I do not pretend to decide; but her condition was certainly one of most frightful collapse, and she was only rescued by the constant and intelligent attention of Mr. Hanbury, who never left her for many hours after the operation, and adjusted the stimulation to the effect required to be produced. The amount of brandy, chloric ether, ammonia, and beef-tea, taken for several hours after the operation, was very large; and any attempt to withhold the stimulants for even two minutes longer than the usual period could not be successfully accomplished until many hours after the operation had elapsed.

The third case differs from the other two in the very material fact of being required for disease. The improvement in the condition of the lungs, which at first determined us to delay the operation, under appropriate treatment, was very marked; and the removal of such a serious source of exhausting influence as the diseased limb has no doubt tended to assist Nature in her subsequent conservative efforts. No operative proceeding short of the entire removal of the limb at the shoulder-joint was, after careful consideration, thought to be sufficient to remove the whole disease; and the examination of the diseased bone after its removal showed the soundness of the decision. The carious head and condyles, and diseased elbow-joint, would have rendered any attempt at removal of the sequetrum unsuccessful, and amputation at the shoulder must have been ultimately required.

ADDENBROOKE'S HOSPITAL, CAMBRIDGE.

REMARKABLE CASES OF FRACTURE.

Under the care of G. M. HUMPHRY, Esq., Surgeon to the Hospital.

CASE I. *Fracture of the Thigh for the fourth time at the same part.* Sarah Hales, aged 60, a stout, rather pale, but not unhealthy woman, in 1847 caught her foot in the carpet and fell, breaking her left thigh bone a little above the knee. In seven weeks she was allowed to get up, and in twelve weeks could walk again. In 1850 her foot slipped on a potatoe, and she again fell, breaking the bone at the same part. In six weeks it was united, and in eleven weeks she was able to get about. In 1852, when raising herself from the floor, her hand slipped, and falling upon that knee the bone was broken at the same place for the third time. She was able to walk in about the same time as before. In January, 1856, when walking along the road her foot slipped and she fell, the thigh-bone giving way for the fourth time at the same part. Whether the bone gave way before she fell, or whether it was broken by the fall she could not tell. She had not much pain, but knew the limb was broken because she had lost the use of it. She was brought to Cambridge the next day; and, when examined, the fractured portions of the bone, which were thickened a little, could be rubbed together as though the ends were rounded and smooth. This gave scarcely any pain, and there was very little swelling or bruising. Mr. Humphry was apprehensive, from this condition of the part, that union would not take place. To give it every chance, the limb was carefully splinted, and perfect rest secured by gum-chalk bandages, etc., which were kept on for eighteen weeks, when, firm union having taken place, she was allowed to get up and go on crutches. These have been some time laid aside, and she can walk without them. The bone is firmly united. There is a little overlapping, but not much thickening.

CASE II. *Spontaneous fracture of both Thigh-Bones.* Sophia Low, aged 56, was admitted April 23rd, 1857. The following is