little pus, and there was discovered disease of the bodies of the seventh, eighth, and ninth dorsal vertebrae. On the thoracic surface of these bones was found a sarcomatous mass, which, on section, looked something like "hard encephaloid" deposit. On microscopic examination, nothing could be seen distinctive of cancer, but the structure seemed to consist of fibrous tissue, mixed with small nuclear bodies. The posterior part of the bodies of the vertebrae was completely hollowed out by ulceration, and infiltrated with pus in the neighbourhood of the ulcerated part. The spinal cord corresponding to this part was slightly softer than the rest. The intervertebral substance between the sixth and seventh vertebrae was quite gone, and every movement of the spine caused these vertebrae to glide on each other, to such an extent as was evidently capable of producing considerable pressure on the cord.

There was no visceral disease except in the bladder, which was soft and thin, and tore with the slightest effort. The urethra was much congested, and in many places almost pulpy. Its lining was completely absent at the commencement of the spongy portion, and there was the cavity of an abscess in the surrounding tissues. Numerous false passages existed here. The scrotum and perineum were livid, and infiltrated with urine.

REMARKS. This case is published as a companion to one which we reported a few weeks ago, and which was also under Dr. Todd's care. There seems little reason to doubt the cancerous nature of the affection in this case also, although the microscope, as is so often the case, failed to give any certain information on the point; nor did the microscopic examination of the cord shew any malignant deposit. The disease in the latter was apparently confined to slight softening, and seemed hardly enough to account for the almost total loss of function which it had sustained. It is interesting to notice the great amount of destruction which the spinal column may sustain without dislocation occurring. In this case the patient had been frequently placed in the sitting and standing posture without accident, although after death the slightest movement appeared sufficient to displace the vertebrae on to the cord.

ST. GEORGE'S HOSPITAL.

CASES OF ANEURISM IN THE MEDICAL WARDS.

By G. GODDARD ROGERS, M.D., Medical Registrar.

[Continued from page 143.]

WE have already reported ten of these cases.

CASE XI. That which would form the eleventh of this series will be found in the JOURNAL for October 31st, 1857. It was introduced there as forming a contrast to a case of pulsating encephaloid tumour of the thorax. At the same time, a few remarks were added on the comparatively few instances in which phthisis is found to accompany aortic aneurism.

CASE XII. *Aneurism of the Left Middle Cerebral Artery.* The case occurred in a man, aged 53, who had been subject to epileptic fits for many years. They usually lasted but a very short time; but on the day of his admission, having made himself unusually comfortable at dinner (according to his friends' account), he was seized with a fit soon after the meal, and never recovered his consciousness. He died three or four hours after he was brought into the Hospital; and on the following day the body was examined.

The pupils were dilated, and there was well marked arcus senilis. The lungs were full of dark frothy fluid; and old pleural adhesions existed. On the surface of the heart were one or two large white patches; both ventricles were empty, and the left one was very firmly contracted. The kidneys were granular, and the bladder enormously distended, but otherwise healthy. The cranial bones were natural. All the vessels of the membranes of the brain were full of blood. The surfaces of the convolutions were flattened and pressed together, but little subarachnoid fluid existing. The subarachnoid tissue covering the cerebellum contained fluid blood. The general structure of the cerebral hemispheres was natural in every way, but in the left hemisphere, rising some distance above the level of the roof of the left ventricle, and extending anteriorly and posteriorly beyond it, was a large cavity outside the corpus striatum, which appeared as it were dissected from the nervous mass. This cavity contained several ounces of fluid and clotted blood, as did both lateral ventricles. The septum of the latter was broken down, the structures forming

their boundaries being all more or less softened. The parietes of this large cavity were very shreddy, and contained small patches of ecchymosis; but the neighbouring parts were not particularly altered by softening or discoloration. At the bottom of the cavity, under the outer portion of the corpus striatum, was seen the rounded outline of the upper surface of an aneurism, dark in colour and very firm; and at the base of the brain, on separating the middle from the anterior lobe on the left side, the lower part of the aneurism was seen almost of a white colour. The sac was divided into two lobules, as if by the pressure of one of the branches of the artery. The entire aneurism was equal in size to a small walnut; and the surfaces of the cerebral convolutions around it at the base were indented, but not softened by it. The arteries at the base of the brain were not atheromatous.

[A brief account of the case will be found in the Pathological Society's Transactions, vol. vii, p. 127.]

CASE XIII. *Aneurism of the Aorta.* J. B., aged 35, a cabman, had been attending in the out-patient room for a fortnight previous to his admission, on account of cough and pain in the chest. For two or three days, hæmoptysis had accompanied the cough. His tongue was coated, pulse sharp, urine high coloured, and bowels costive; and he complained of severe pain in the left side. Salines with antimony were given, and he was bled to twelve ounces, but with only temporary relief. At night he complained much of acute pain in the epigastrium and over the dorsal region. On the left side of the chest there was dulness at the lower part of the lung, and above crepitant râles. The second day he brought up a considerable amount of frothy blood; and expired in the afternoon.

The body was examined twenty-one hours after death. Frothy serum mixed with blood was issuing from the mouth. In both pleural cavities were elongated bands of old adhesions. The posterior part of the left pleural sac was full of fluid blood, and a large ragged opening existed here, communicating with an aneurismal pouch of the descending aorta. This pouch was large enough to contain a small apple; its opening into the artery was of very large size; its walls were very irregular in thickness, and at one part so thin as to be nearly transparent. The trachea and bronchi were deeply stained with blood; and the tissue of the lung in the part corresponding to the aneurism was loaded with blood, so as to sink readily in water. The aneurism must, therefore, have had some communication with the interior of the lung; but this opening had been accidentally cut across in taking the lung out. The tissue of the lungs and heart appeared healthy. At the base of the right lung was a small mass of old cretaceous deposit. The other viscera were natural.

[To be continued.]

UNIVERSITY COLLEGE HOSPITAL.

STRANGULATED INGUINAL HERNIA: OPERATION: SECONDARY HÆMORRHAGE FROM THE SPERMATIC CORD.

Under the care of J. E. ERICHSEN, Esq.

[From Notes by T. B. JEWETT, Esq., House-Surgeon.]

JOHN R., aged 43, was admitted on January 17th. He had had hernia on both sides, but larger on the right, for twenty-five years, and had worn a double truss during the whole period. The hernia on the left side came down on the morning before his admission, in consequence of a violent effort.

On admission, the penis was found to be entirely embedded in the scrotum, which was as large as a child's head. On the right side there was a hydrocele, but the canal could be felt to be patent. On the left side there was a large hernial swelling, in which no impulse could be felt on coughing, and which yielded a dull sound to percussion over its whole extent. The taxis had been applied before his admission, and was repeated, in the warm bath, after admission, but unsuccessfully. The operation was therefore proceeded with at once. Mr. Erichsen first tapped the hydrocele, and evacuated a pint and a half of yellowish serum; he then proceeded to the hernial tumour. The neck of the sac was exposed, and an attempt was made, but without success, to reduce the tumour without opening the sac, the neck of which was much thickened. It contained a knuckle of congested small intestine, and some thickened omentum adherent to the epididymis and cord. The omentum was tied and removed. He was ordered a few doses of calomel and opium; but the pills seemed to occasion vomiting, so a dose of laudanum was given. The bowels acted next day.