CALCUTTA EYE INFIRMARY.

ILLUSTRATIONS OF SOME OF THE CURABLE FORMS OF AMAUROSIS.

By WILLIAM MARTIN, Esq., F.R.C.S., late Superintendent of the Infirmary: and Professor of Ophthalmic Surgery in the Calcutta Medical College.

II. AMAUROSIS FROM CACHEXIA.

IN the preceding division were put a certain number of cases exemplifying the disease as arising from a state of general debility of the system. We now come to examples of a condition of the retina, producing a nearly similar train of symptoms, but arising from a somewhat different cause; that is, a cachexia, caused not by mere excessive loss of the nutritious fluids of the body, but by some poisonous action exerted by the prevalence of a particular disease or train of diseases.

CASE I. Jaddoo, aged 30, was admitted as an in-patient, September 27th, 1856, with amaurosis. The past year he had suffered from syphilis, for which he had been salivated. Two years previously he had enlarged spleen. He had become anasarcaous, and had suffered from rheumatism. He saw objects as through a dense fog. There was slight strabismus. He was ordered tonic mixture, and counterirritation.

October 6th. He was improving in both eyes. He afterwards suffered from cough, dyspepsia, etc., and was much benefited by the use of sarsaparilla. He also suffered from dysentery.

November 4th. He was in better health, and the sight had improved, he said, six parts out of sixteen.

November 26th (two months after admission). He had continued to improve, and was discharged relieved.

REMARKS. This is a peculiarly interesting case, as there were a number of predisposing causes—syphilis, mercurial action, rheumatism, spleen-disease, anasarca, acute ophthalmia, any one of which might be enough to induce a cachectic state of the body; but which, although acting upon the constitution in conjunction, all or several of them, only produce a temporary or functional amaurosis. The treatment was varied according to the many changes of disease which were exhibited, and much occurred to depress the vital powers; yet after about two months stay in Hospital, he was reported as so much better, that, according to his own account, he was half cured, and the improvement was progressive.

CASE II. Toonoo, aged 55, was admitted as an in-patient into the Infirmary, on September 30th, 1856, with amaurosis. He had suffered three years previously from syphilis, for which he was salivated. A year before admission, he had attacks of ague and spleen disease. He was ordered quinine and soda.

October 4th. He said his sight had slightly improved.

October 11th. He continued to improve. The tonics were continued, and counterirritation was applied.

November 5th. He left Hospital relieved. He said he was half cured; and the improvement was progressive.

REMARKS. Here was also a combination of causes for a special cachectic state of body, and which only induced a functional amaurosis.

CASE III. Horee, aged 16, was admitted as an in-patient on October 22nd, 1856, with amaurosis. Three years previously he was subject to fever and spleen disease. Two years before admission he had catarrhal ophthalmia; and this was followed by nyctalopia. He was ordered local depletion, and counterirritation, followed by tonics.

November 10th. He said his day sight was better; but he could not see better after sunset. He continued to improve somewhat.

December 6th. He was discharged relieved.

REMARKS. The causes here were spleen-disease and attacks of acute ophthalmia. These induced a state of functional amaurosis, which was at any rate partly curable; probably it only required time for a perfect cure.

CASE IV. Beedee Bodun, aged 39, was admitted as an out-patient of the Infirmary, May 22nd, 1856, with amaurosis, complicated with jaundice. He was in a cachectic state. Ten years before he suffered from syphilis, and was affected by the action of mercury. He also, when admitted, suffered from rheumatism, and was an opium eater. He ceased to attend.

REMARKS. I find I have nothing to note as to the progress of the case; but still, from the similarity of its symptoms to those which proved to be only cases of functional disease, I think we may have a strong presumption that this was one of the same kind.

CASE V. Luckeenarian, aged 25, was admitted as an out-patient, July 8th, 1856, with amaurosis. He had for the past thirteen years suffered from fever, nyctalopia, muscæ volitantes, etc. The disease was complicated with hemiopia, the sight being in the following peculiar condition, showing a state of chronic irritation of portions of the nervous system of the brain and eye. With both eyes he could only see objects which lay on his right side. With the right eye only, he could see features, etc., indistinctly, but equally well on all sides. With the left eye only, he saw on all sides equally, but could not distinguish features. He could only see objects moving about between his eye and the sun.

REMARKS. Here was a peculiar cachectic state of constitution, brought on originally by the fever, which had previously caused various disordered conditions of the internal tunics. There was a state of the system different from mere anæmia or general debility. Unfortunately, I knew nothing of the progress of the case, as the man did not continue his attendance.

CASE VI. Ramdoon, aged 35, was admitted as an out-patient, August 28th, 1856, with amaurosis. Four years previously he had syphilis, for which mercurial fumigations were used, and he became salivated. Some time afterwards he had secondary syphilis, which was not severe. He said it disappeared without treatment in fourteen days. Five months after that, the right eye became inflamed, and soon afterwards the left. The action of the pupils, both associated and distinct, was normal. He could not see distant objects; near objects better, but not clearly. He saw best with a small light. He was ordered tonics, and stimulating drops to the eye.

September 3rd (sixth day). His sight was a little clearer.

September 10th. He saw both near and distant objects better.

September 13th. He was improving daily. He now saw near objects with perfect distinctness.

September 16th. He ceased to attend. The inference is that he would in time have been cured.

REMARKS. The syphilitic taint of the constitution, as well as the great improvement of vision from the treatment employed, are well marked in this case.

CASE VII. Seebchurn, aged 30, was admitted an out-patient on September 1st, 1856, with amaurosis. He had been much subject to fever. Some years previously he suffered from gonorrhœa and buboes. He had also had piles, dyspepsia, and occasional diarrhœa. The impaired state of vision had come on gradually within the last twenty-four days. He was ordered simply tonic mixture.

September 17th. He had more strength.

September 23rd. His sight had improved within the last few days. He was in progress to be cured, but left Hospital.

REMARKS. The bad state of the body here seems to have arisen principally from disordered function of nutrition, of which the dyspepsia, hæmorrhoids, etc., were the most prominent symptoms.

CASE VIII. Kalleedoss Dey, aged 40, was admitted as an out-patient on December 26th, 1856, with amaurosis, immediately following cephalalgia. He had suffered from rheumatism, and had been twice mercurialised. He was ordered a tonic mixture, containing sarsaparilla, with iodine and iodide of potassium; and five grains of blue pill at night.

January 9th, 1857. He still suffered from pains in the head. Counterirritation was applied, and decoction of bark ordered.

January 19th (twenty-fourth day after admission). He said he was stronger. His sight remained the same. The medicines were continued. He ceased to attend.

REMARKS. The cachexia here seems to have been due to the mercurial action, having been produced twice when the powers were enfeebled by rheumatic disease; from the similarity of its symptoms to those of other cases of functional amaurosis, I have not the slightest doubt that it may be put in the same class.

GENERAL REMARKS. In the group of cases, the outlines of which are here given, the indications of treatment were, gradually to restore the tone of the system; to bring back the functions, particularly of the liver, stomach, and skin, to a healthy state; and to cause the elimination, if possible, of the poisonous influence which had pervaded the system. These indications were not always readily fulfilled; much perseverance was required, and it was found necessary often to change the medicines used: the patient, moreover, often became dissatisfied with the slow progress made, and discontinued attendance. The improvement, however, which resulted in the greater number of cases, although not so nearly amounting to an absolute

cure, as in the preceding group, was sufficiently satisfactory. The medicines which were found of most use were gentian or cheyretta, quinine, preparations of iodine, and occasionally mercurials.

[To be continued.]

24, George Street, Hanover Square, W.

Original Communications.

REMARKS ON THE THEORY OF ELIMINATION IN THE TREATMENT OF DISEASE.

By C. HANDFIELD JONES, M.B., F.R.S., Physician to St. Mary's Hospital.

[Read before the Harveian Society.]

It has been well said by one of the masters of our profession, that the development of a principle is like setting up a brilliant light in the midst of a large hall, which enables the eye to take in the whole in all its fair proportions; while the investigation of details is like exploring the same hall in all parts by the aid of a taper. The latter process is most necessary; without it there can be no safe ascent to higher ground, no sound induction; but yet it ought always to be conducted, as far as possible, under the guidance of some *informing* (shaping) idea, which may connect and colligate our facts, and enable us at last to arrive at a law or principle. I am much inclined to think, that of late years we have been over-busy in accumulating facts without sufficient reference to principles, or, to use Dr. Whewell's term, colligating ideas. Blood-globules have been counted, tissues have been scrutinised both healthy and morbid, secretions analysed, symptoms described, and râles defined minutely. Far be it from me to decry this painful labour; it has produced fruit, and will, I doubt not, yet more. But have we not, amid all these particulars, thought too little of the general truths which they contain, and which we ought to strive to educe? I believe strongly that each age has its mission, its more particular work to do; and the mission of our age may be the accumulation of facts, that of a succeeding age their interpretation. But still I would we had more workers after the example of an eminent member of this Society, lately passed away from among us, whose observations and experiments always went straight to a point. I need not say I allude to Dr. Marshall Hall.

The topic which I wish to bring before the Society to-night is the consideration of a principle of wide application in therapeutics; viz., how far, in various diseases I shall refer to, we ought to proceed on the view of endeavouring to eliminate or get rid of some *materies morbi* which we suppose to exist. In many disorders we have more or less certainty of the imbibition of a poisonous material; and it would seem a natural inference, that to get rid of this material should be the chief aim of our efforts. It appears to me that this conception, in a more or less distinct form, is extremely prevalent, and that there is some risk of its being carried too far, and prompting an injurious practice. Thus, Dr. Todd bases his treatment of scarlatinal drowsy on the elimination of the poison. Dr. G. Johnson recommends castor-oil in cholera to procure the excretion of the poison by the bowels, and regards Bright's disease as the result of the renal cells striving to remove morbid materials from the blood, and becoming damaged in so doing. Mr. Simon speaks of cancer as an "eliminative effort"; and Mr. Paget appears to take nearly the same view.

These are high authorities; and I fear I shall be considered guilty of heresy in presuming to differ from them. How far I may be justified, it will be for my audience to decide.

Let me ask attention, first, to one or two introductory remarks. When we meet with a case of poisoning from the ingestion of deadly drugs, such as opium, arsenic, corrosive sublimate, or with a case of snake-bite, we do not think much of trying to expel the poison from the blood, and so relieving the sufferer. We know this would be a vain attempt, and so we set ourselves to counterwork the virus, to stay its deadly action by remedies which have an opposite effect. The snake-bitten man imbibes brandy, arsenic, ammonia, etc., plentifully; and if the poison be not too potent, his system is rallied from the fatal prostration, and he recovers. The animal to which prussic acid or woorara has been administered is apparently lifeless, and if left to itself will soon be really so; but we use artificial respiration for a time, and at length the poisonous influence passes

by, and the respiratory muscles resume their play. So, also, in a case of opium poisoning, we oppose the tendency to fatal coma by artificial excitement. Here, then, are exemplars for us; we see clearly in these cases what course we must follow. We must aid Nature to get over the difficulty; we must not think to take her work out of her hands; but we must try to sustain her powers where they fail. In one instance, indeed, we can do more, and eliminate the poison by means of drugs. I refer, of course, to the detachment of lead from the tissues of the body, and its excretion in the urine, by the agency of iodide of potassium, as first rendered probable by M. Melsens, and recently confirmed by Drs. Parkes and Sieveking. This, however, is almost, if not quite, a solitary instance.

Another remark I will offer is this; that it seems of more importance to have a sound, vigorous vitality of the tissues, than that the blood should be of absolute purity. If the life of the solids is feeble and languishing, they can do little, even with the best blood. On the other hand, if they are vigorous, they will nourish themselves and function well, although their nutrient fluid be of inferior quality. Not long ago I had a patient who was markedly jaundiced, but who went about his business, ate and drank as usual, and had no sensation of illness. Another patient has long suffered from Bright's disease, with anasarca, and his urine is still highly albuminous; but for the last six months he has enjoyed exceedingly good health, has been free from dropsy, and looks robust and vigorous. In both these cases (and they are no very uncommon ones) the tissues nourished themselves well out of blood which was far from healthy. Again, with regard to miasmata: a weakly man is exposed to the infection of fever; he imbibes it, and after a time droops and sickens. A companion of stronger frame exposed to the same escapes altogether. What has made the difference in these two? Both must have received the poison into their blood as a matter of physical necessity; but the nervous system of the weaker has succumbed, while that of the stronger has resisted the poison. Or an army is retreating over a line of country pursued by a victorious foe. Both are exposed to the same influences; but the one, dismayed and depressed, is visited heavily by disease, while the other, exulting and confident, is comparatively healthy. Again, if two boats' crews go up the same African river, and are alike exposed to the miasma, while the one is protected by a daily modicum of quinine wine, which is denied to the other, it is notorious that the latter will be to a man smitten with fever, and that the former will almost, if not entirely, escape. The quinine, like the mental stimulus, increases the resisting power of the nervous system: so that the poison received into the blood is inoperative. Once more, when pus is injected into the veins of an animal, the effect depends much on the vigour of its system. A feeble animal—as a rabbit—dies surely; a stronger resists and survives.

I hope that what I have said will not be taken to mean that I consider it of no importance whether the blood be contaminated or not. I am as fully satisfied of the mischief which may ensue on blood-poisoning as the greatest humoralist can be. My only aim is to hold the balance even, and to set forth duly the resisting power of healthy tissues.

Passing on to the proper subject of this communication, let us first take for our consideration the *exanthemata*—variola, scarlatina, measles, erysipelas. What shall we regard as the true principle of correct treatment in these, so far as we find that we have any power? It is certain that the morbid phenomena depend on the reception of a poison into the system. Can we hope to neutralise or expel this? As to the possibility of neutralising the virus, medical science, we must confess, is wholly silent. As to expelling it, the opinion seems held by many that the exanthem is an eliminative effort, and therefore it would seem that, even if we prefer another channel to that selected by nature, we ought to pursue the same indication. But is it certain that the exanthem, or any other of the morbid phenomena that occur, are eliminative efforts? May we not just as well regard them as reactions of the tissues against the poison, as so many foci of irritation? Doubtless we must not suppress the cutaneous eruption, because it is better to have the skin inflamed than the bronchial or intestinal mucous surface. But neither do we get any good by increasing the eruption as much as possible. Take the case of *small-pox*; the cutaneous is not the only channel by which exudation of the infecting matter takes place. Dr. Copland states "the miasm of small-pox is given out from all the mucous, cutaneous, and excreting surfaces—the exhalations, the secretions, the excretions, the matters in the vesicles and pustules, and the scabs, all contain this poisonous material." Indeed, there is reason