mately adherent to the periosteum around; it was, moreover, sunk deep in the bone, which presented a cup-like cavity. Perforation of the bone was not actually made out; nevertheless, as there could have been but a very thin plate of bone left, Mr. Athol Johnson thought it advisable, after emptying the cyst, not to remove the small portion adhering to the centre of the cup-like depression. The child did well. The parts healed, with a depressed scar adherent to the bone beneath.

When seen a few days ago, the wound was firmly healed, but the depression had not been filled up. The child was in perfect health.

II. SYME'S AMPUTATION.

We had the opportunity of seeing lately three stumps of Syme's amputation; and, as there is at present some difference of opinion with respect to the results of this operation, we have thought it advisable to describe the state of the parts. We accordingly append a short description of each case.

CASE I was that of a girl, named Sarah Fennell, a patient of Mr. A. Johnson, at the Children's Hospital, in whom the operation was performed in June 1854, for disease of the tarsus, she being at that time 5 years old. There was nothing peculiar about the operation; no sloughing took place. Four years have now elapsed since the operation, and she suffers no inconvenience in the stump, being able to bear her weight with comfort to herself. She wears a sort of bucket-boot, made by a common shoemaker, and can walk or run about as well as any girl of her age. A cast has been taken.

CASE II is that of a boy, at the same hospital, named W. C., aged 5, in whom the operation was performed fourteen months since, for disease affecting the os calcis and the astragalus. An attempt had been made at another London Hospital to save the foot by removing the diseased bone, but unsuccessfully; and, the health beginning to suffer, Mr. Johnson performed Syme's operation. The part united well, without any sloughing; and the boy is able to run about on the stump, without any shoe at all. He wears, however, a similar boot to that of Sarah Fennell.

In both cases, the amount of shortening of the limb is very slight. A cast has been taken of this stump also.

CASE III. The third case was shown in the theatre of St. George's Hospital, a few days since, previous to the removal of the stump by amputation through the leg. The patient, a young and healthy man, had had the operation performed two years ago, in the country. His convalescence was very rapid, so much so, indeed, that he was only confined to bed for a fortnight; and the wound seems to have been soundly healed in a month. It was a long while, however, before he began to use it, as he was waiting till he could procure an artificial foot; and, when he did do so, he soon found that the stump was painful, and that the skin was cold to the touch. The pain was not so great as to prevent him getting about; but at night, when he had been using the stump for some hours, it used to become much worse, and to be accompanied by spasmodic movements of the limb. This tenderness to pressure was remarkable at one point of the stump just below the middle of its anterior extremity, which was supposed to depend on some affection of the anterior tibial nerve. The skin was blue and cold. As the patient represented that the annoyance which the stump occasioned to him was intolerable, Mr. Pollock acceded to his request to remove it.

On examining the stump, it was found to be quite soundly healed, and was in every respect a very healthy and good one, except that the nerves terminated in enlarged extremities, forming a species of small tumour. This was most distinct on the posterior tibial nerve, in which the swelling was larger and lobulated. It was formed principally, if not exclusively, of fibrous tissue, and adhered firmly to the back of the tibia. The swelling on the anterior tibial nerve was smaller, but was more tucked in under the section of the bone, so that it was probably more pressed on in walking. The arteries were pervious down to the end of the stump. It was noticed that the fibula was prolonged lower than the section of the tibia, as if its end had grown since its division. To this elongated portion the tendo Achillis was inserted.

REMARKS. The first two cases are good instances of the great advantage of this operation, when successful; viz., that it enables persons to dispense with the services of the instrument-maker, and to get on perfectly well with a boot made by their own shoemaker. How great this advantage is to the poor, no one will question. Who would hire a domestic servant with a wooden leg? while, in Mr. Johnson's first case, there is every probability that the little girl will be so occupied when she grows up. It is indeed said by some instrument-makers, that

they find it easier to adapt an artificial foot to well made stumps of the leg, than to those left by Syme's amputation; but what we are contending for now is, that it is very important, when possible, to enable patients to dispense with the aid of an instrument-maker entirely.

With regard to the third case, the stump has indeed proved useless, and has been obliged to be sacrificed. This, however, did not arise from any cause essential to the operation, but from an accidental affection of the stump, which might just as well follow on an amputation of the leg. It may, indeed, possibly be the case, that, in cutting the posterior flap, operators are apt to leave the nerve so long as to come on to the lower end of the bone, and thus to be in the way of pressure when the stump is used. This was not clearly proved in the case before us, since the posterior tibial nerve, which would be most likely to be left long, was not apparently so much in the way of pressure as the anterior; but it is worth remembering, that such a cause may be the occasion of future tenderness in the stump, as nothing would be easier than to take off any portion of either of these nerves which might seem advisable during the operation.

CALCUTTA EYE INFIRMARY.

VARIOLOUS CORNEITIS.

By WILLIAM MARTIN, Esq., F.R.C.S., late Professor of Ophthalmic Surgery, Calcutta Medical College.

SMALL-POX has long been recognised as a prolific cause, predisposing or exciting, of various diseases of the eye. These more commonly affect the external parts; but not rarely we find iritis and inflammations of the internal tunics generally, owing to a recent attack of small-pox. Of affections of the external parts from this cause, the most troublesome are inflammations of the conjunctiva, sclerotic, or cornea; ulcers of the cornea, inflammation of the lacrymal passages, etc., etc. I propose to relate a few instances of corneitis, arising at a not remote period from an attack of small-pox.

CASE I. Noor Ali, aged 5, was admitted as an in-patient May 9th, 1856, having secondary ulcer of the cornea, with a corneal abscess. This commenced nine days after the separation of the crusts, in an attack of variola not occurring from inoculation. Much inflammation was present. The treatment consisted in the application of leeches and a blister, and the administration of a purgative.

May 13th. He was ordered to take a grain of disulphate of quinine three times a day, and to have cold water applied to the eye.

May 16th. The quinine was repeated four times a day.

May 20th. There was an extensive sloughing ulcer. The medicines were repeated; and three grains of mercury with chalk were prescribed, to be taken twice a day.

May 23rd. The ulcer was inclined to heal.

June 2nd. The ulcer was nearly healed. The matter of the abscess had become absorbed. The cornea was densely opaque. The powder was omitted; the quinine and cold water application were continued.

June 12th. He was doing well. The medicine was continued, and a blister was applied.

June 20th. He left the Hospital, having only corneal opacities left.

REMARKS. In this case, the commencement of the attack of ophthalmia took place at a period more remote from the invasion of variola than is usual. There could be no reasonable doubt about the cause. It is to be observed, that the disease was of as severe and intractable a nature as a primary variolous ulcer could have been, although all external manifestations of the constitutional disease had disappeared for many days. This is perhaps to be accounted for by the fact of the constitution itself having had time to become tainted by a special cachexia. The curability of such a dangerous local affection, when coexisting with this cachectic state of the system, is interesting and instructive.

CASE II. Dwarkanauth, aged 30, was admitted as an in-patient May 27th, 1856, with variolous corneitis, commencing twenty days after the appearance of the eruption in variola supervening upon inoculation. The treatment consisted in leeching, blisters, calomel, with opium and quinine. He was eventually cured.

REMARKS. Here also there was a sufficient interval between the invasion of the constitutional disorder and the local affection to make it probable that the variola acted as a predis-

posing rather than an exciting cause. I have generally found that the cases of variola which were induced by inoculation have been milder than others; and this may account for the facility with which this patient was cured. I conclude that this may be attributed to the absence of any marked constitutional debility.

CASE III. Goolab, aged 25, became an out-patient May 17th, 1856, with variolous corneitis. The disease commenced twelve days after the separation of the crusts. The same remarks are here applicable with regard to the interval of time between the variolous outbreak and the invasion of the corneitis.

CASE IV. Ramjaun Ali, aged 22, became an out-patient June 6th, 1846, on account of corneitis and its results—dense opacities of the cornea. The inflammation had commenced fourteen days after the invasion of the fever. The opacities were in part removed.

REMARKS. In this case, the interval of time was not as great, but the local affection seems to have been of a mild character; for I was only called upon to treat the consequences of the corneitis, and the case was in progress when the patient left off attending. These results were not indelible.

CASE V. Onrajee, aged 9, became an out-patient on June 9th, 1856, having chronic abscess of the cornea. Three months previously, an attack of variola came on, not from inoculation. The friends said that, at this time there was a small-pox pustule on the surface of the eye, and that the crust from this became separated at the same time as from the pustules on other parts of the body. Inflammation of the eye came on ten days after this separation. The treatment consisted of local depletion, purgatives, hydrargyrum cum creta, etc.

June 23rd. The mouth had become slightly affected; but the patient was improving, and the matter of the abscess was becoming absorbed.

July 23rd. The abscess was healed. The opacities of the cornea in its vicinity were thinning. The patient was in process of being cured.

REMARKS. The corneitis does not seem here to have been part of the external manifestation of the variola, but to have occurred subsequently. The pustules, said to have been in coexistence with the small-pox in the body, most probably only affected the eyelids.

CASE VI. Banepur, aged 10, was admitted as an out-patient on July 14th, 1856. Variolous ophthalmia was present in both eyes. In the right, in which the attack was of a primary nature, the cornea had sloughed; in the left, the disease had appeared secondarily—the effect being extensive ulceration and opacities of the cornea. The attack of small-pox had occurred six months previously.

REMARKS. This case is particularly instructive; as, some time after the invasion of the primary eruption, which was attended by so much local disease that one cornea was destroyed by sloughing, another distinct disease, involving also specially the cornea in the other eye, was set up. Nothing seems more probable than that the primary affection was a variolous pustule, having its seat in the conjunctiva and cornea, running a rapid course, and producing mortification of those parts in the same way that variolous pustules act in other parts of the body; and that after a certain interval of time, when the local affection had subsided, the constitutional taint predisposed to the secondary inflammation which seems in such cases to have a tendency to attack some of the vital parts of the organ, which had escaped the more dangerous primary disease. This secondary affection being altogether milder than the other; in this instance, there is every reason for thinking that the disease was perfectly curable.

GENERAL REMARKS. The cases above noticed afford good instances of the particular forms of the corneitis, which owes its origin to variola. In almost all, the disease showed itself in the most severe form, the one which rapidly runs into destructive ulceration of the structure, instead of into the granulo-vascular degeneration. This proneness to a destructive form of inflammation would be easily accounted for if we could look upon the disease as simply a variolous pustule formed upon the membrane, as in other parts of the surface of the body, but the history of the case will seldom permit us to look upon the case as being one of this description. Indeed, it has been doubted whether the specific variolous pustule is ever formed on the surface of the body, by observers who have had so much experience in small-pox as Dr. Gregory and Mr. Marson, who say that the phlyctenula or pustule that is found to arise in the consecutive stages of variola, are identical with those which arise in common catarrhal or scrofulo-catarrhal conjunctivitis (see Wharton Jones's second edition, page 163).

However, I have seen several cases in which a pustule or ulcer was formed upon the conjunctiva or cornea, ultimately affecting both structures at the same time with variolous pustules on other parts of the face and body. In such cases the pustule has been much more rapidly destructive in its effects than where it has occurred in the succeeding stages of the small-pox, and has in fact produced the same changes, leading to mortification of the subjacent parts, as the variolous pustule does upon the cutis. There seems to be no reason why the specific pustule should not occur as well upon the conjunctiva, which possesses a combination of characters of the skin and mucous membrane, as upon the skin of contiguous and continuous structures. The last of the cases given above seems to confirm this view, for one eye had been lost by a destructive affection of the cornea, which had occurred contemporaneously with the pustules on the body; the other eye being only secondarily affected with an inflammation which was much milder in character than the disease which had caused mortification of the cornea in the first eye. Nothing seems more probable than that in this case we had an exemplification of the two kinds of variolous affections of the cornea; in the one the specific pustule leading to destruction of the part implicated; in the other, the milder form of sympathetic secondary corneitis having its origin in a predisposition, caused by the disease having taken effect upon the constitution. At the same time, the fact of only one case in six reported having the appearance of being an instance of specific variolous pustule of the eye, indicates that such a case must be rather rare.

With regard to the period after the accession of an attack of variola, at which the secondary affection, which ushers in corneitis and ulcer of the cornea, comes on, it seems to be subject to considerable variations. Some authors say that it is about the eleventh or twelfth day from the first appearance of the eruption, and when the secondary fever has commenced, (see W. Jones, second edition, page 162); but in the six cases above noted the time varied from the tenth to the twenty-second day after the appearance of the eruption, and the average was eighteen days, so that when this affection occurs secondarily, in most cases probably it is on account of not so much the secondary fever, as of the dyscrasia produced by the disease.

24, George Street, Hanover Square.

Original Communications.

ON THE TREATMENT OF CONTRACTED JOINTS.

By HOLMES COOTE, F.R.C.S., Assistant-Surgeon to St. Bartholomew's Hospital, and to the Royal Orthopaedic Hospital.

[Read before the Royal Medical and Chirurgical Society, May 11th, 1858.]

I BEG to call the attention of the Society to the Treatment of Contracted Joints; and, in so doing, I shall limit my remarks to the knee, although the principle of treatment must be the same in every articulation of the body. But, in the first place, I must assume the following propositions as granted.

1. Diseases of joints, whether acute or chronic, may be almost invariably referred to that state of constitution called struma in the young, and to rheumatism or gout in the adult.

2. The progress of the disease is always painful and tedious, but no morbid changes occur in any of the tissues by their nature irreparable. The synovial membrane may be thickened, but that is the first stage leading to its absorption; the articular cartilage may be removed, but the extremities of the bones become covered with other tissues; ligaments, softened and elongated, will in course of time become firm, and will even recontract, if put under proper and favourable conditions.

3. The severer changes with which we meet in practice are mainly due to want of proper maintenance of repose of the joint during the course of the disease. Where do we notice instances of inflammation of the cancellous texture of the bone leading to abscess? At the lower extremity of the femur, at the upper and the lower extremities of the tibia, but not in the bones of the upper extremity. These accidents happen, then, just to those bones which support the weight of the body.