several times during the day. There is just a slight limp in her walk.

On referring to the case-book, it is stated that, at the operation, Mr. Fergusson removed a slice half an inch thick from the femur, and two portions from the tibia, together about an inch and a quarter in thickness.

ST. GEORGE'S HOSPITAL.

SUPPURATIVE INFLAMMATION OF LIVER FOLLOWING A BLOW.

By G. GODDARD ROGERS, M.D., Medical Registrar.

J. B., aged 15, a plumber on board ship, was admitted under Dr. Page, on July 13th last, with considerable distension of the abdomen. His face was pale, but his aspect was natural, and without any appearance of distress. He had never suffered from colic or lead palsy. Three weeks before admission, one of his shipmates struck him violently on the pit of the stomach three times, and ever since he had suffered from constant pain in the epigastric region and frequent vomiting. Clots of blood sometimes appeared in the vomit. On the 11th, he brought up two potfuls of vomit, but had no return of sickness previous to admission. The bowels had been exceedingly constive, nothing but a few scybalous lumps having passed, and then only two or three times since the blow was given. The abdomen seemed to be distended with flatus, and continued so for three or four days. Repeated doses of castor oil failed to open the bowels, but they responded to an injection (the simple "enema oleosum" of the Hospital *Pharmacopœia*); and after free action had been established, the pain at the epigastrium diminished. The urine was clear and free from albumen. On the 24th, there was more pain and increased fulness in the right hypochondrium. He also complained much of nausea, and lost his appetite. A great deal of this uneasiness no doubt depended upon his having surreptitiously obtained some gooseberries. The stools passed on the 28th contained much mucus and gooseberry husks.

August 3rd. Watery stools passed after the injection. He felt altogether better.

August 5th. The stools were natural in colour, and free from mucus.

August 7th. After this date, there was not much change in his appearance for nearly three weeks. The abdomen became much reduced in size. He took sherry wine, and a little grey powder, rhubarb and Dover's powder every night, and nitre draught with iodide of potassium three times a day. The bowels were generally obstinately confined, and he had to take castor oil, or sulphate of magnesia, in addition to using the injection.

August 27th. Rigors came on, with severe pain in the right hypochondrium, which was not relieved by the application of leeches. Day by day the abdomen increased in size, and it looked as if an abscess were pointing externally over the hepatic region. The integument was red and very tense, and the whole belly was hot. He complained of great nausea, and vomiting came on on September 8th; at the same time, a quantity of yeasty-looking and bloody fluid was passed by the bowels.

September 22nd. The sickness became more distressing. Hydrocyanic acid, creasote, and other remedies were given. Three doses of castor oil and an enema were administered before the bowels could be made to act. The sickness continued throughout the night. In the morning, the abdomen was softer. Over the lower margin of the right lobe of the liver and over the epigastric prominence there was excessive tenderness. In the umbilical and left iliac regions percussion gave a clear sound.

September 27th. At 6 p.m. he was attacked with violent hicough, and pain in the throat and chest. He was unable to speak up to the 7th of October.

On the 29th September, the urine had to be drawn off. The remarkably loud hicough continuing day and night, galvanism was applied to the pit of the stomach on the 2nd October, and repeated on the following day. Towards midnight the hicough ceased. By signs he gave us to understand that he had no sensation in his tongue; and on the 5th Oct. he professed not to feel anything touching his face, whilst the scalp remained as sensitive as ever. On the 5th Oct. the bowels were very freely unloaded, as much as two chamberpotfuls coming away. A great portion of the matter thus passed was puriform and yeasty. After this, the size of the abdomen was very materially diminished; his voice, as before stated, returned on the 7th Oct., and he rapidly improved in his general

appearance. On the 18th Oct., the abdomen was found more distended again, and he was ordered to keep his bed, which he had left for some days. A full dose of castor-oil was at once given, and much greenish puriform matter was brought away. On the 20th and 21st Oct., the purgative treatment was again adopted, the stools being light-coloured, but more healthy-looking. From this time all abdominal swelling and uneasiness departed, and he left the hospital perfectly cured on the 3rd November.

REMARKS. Abscess of the liver, caused by a blow, is of far less frequent occurrence than at first sight would seem likely. For this statement, however, we have the authority not only of Andral in his *Clinique Médicale*, but also of Dr. Budd in his elaborate work on the *Liver*. The same degree of force is far more likely to rupture the viscus, closely packed as it is, than to set up suppurative inflammation; and in the case before us, a considerable time (nearly eight weeks) elapsed before the formation of matter was fully established, showing that the liver does not readily take on this form of inflammation from any external violence. It is very probable that much of the extreme abdominal hardness was owing to sympathetic tension of the recti muscles, concerning which Dr. Budd has some valuable remarks. The loss of voice was one of the most remarkable features of the case, and I have heard no satisfactory explanation given of it, and am quite in the dark myself as to its true cause. I believe that a great part of the lad's declaration by signs of his loss of sensation is not to be depended upon. His case excited an unusual degree of interest, and he was, I fancy, anxious to achieve a spurious notoriety by exhibiting the most anomalous symptoms.

Original Communications.

OBSERVATIONS ON CERTAIN FORMS OF DISEASE OF THE LUNGS.

By EDWARD LATHAM ORMEROD, M.D. Cantab., F.R.C.P., Physician to the Sussex County Hospital.

II.—PNEUMONIA. VARIABLE AMOUNT OF SYMPTOMS IN DIFFERENT CASES; CHRONIC PNEUMONIA (?); PNEUMONIA OBSCURED BY THE SUPPRESSION OR EXAGGERATION OF PARTICULAR SYMPTOMS; UTILITY OF FREE PURGING; PAIN IN THE ABDOMEN AS A SYMPTOM OF PNEUMONIA; EPIGASTRIC PULSATION; DELIRIUM; COMA; PNEUMONIA COMPLICATING OTHER DISEASES; PNEUMONIA OF FEVER, OF RHEUMATISM; ACCESSORY PNEUMONIA; CONNEXION BETWEEN PNEUMONIA AND JAUNDICE; CRITICAL PNEUMONIA.*

THERE are few diseases which manifest their presence by more distinct symptoms and physical signs than pneumonia, as a general rule. There are few, on the other hand, where, on occasion, these are more strangely altered or more entirely suppressed. Sometimes the assistance of the physical signs is quite superfluous, so clearly is the nature of the disease expressed in other ways. The pain, for instance; the dusky face; the high nervous excitement; the short frequent cough and rusty expectoration,—may leave auscultation only to tell us just whereabouts the disease is, and how much of the lungs it occupies. But sometimes these symptoms are almost entirely wanting. There may be no pain, no cough, nor expectoration—nothing to attract attention to the chest at all. The signs of imperfect aeration of the blood may be masked by anæmia, or be explicable on some erroneous hypothesis which other coexistent symptoms suggest: or pain or nervous excitement may so far predominate as wholly to alter the features of the case, and to lead us to seek their respective causes in the head or the abdomen, rather than in the chest.

Again—to put all the difficulties and obscurities in the foreground—the physical signs too, in their turn, may fail us. The condition of the patient, for instance, may prevent our examining his chest; or (and this is not so very uncommon a case), from the disease occupying the centre of the lung, rather than the surface, the physical signs may be wholly in-

* The paper on Pneumonia, which constitutes a portion of this series, was read in an abridged form at the meeting of the South-Eastern Branch of the British Medical Association, at Brighton, in this year. I trust, in publishing it now *in extenso*, and in connection with the rest of the series, that I am doing no more than fulfilling the wishes of the members so kindly expressed on that occasion.