great relief to the breathing. The sulphate of copper was continued at longer intervals for thirty-six hours, together with beef tea, port wine, etc.

July 16th. He was much improved.

July 17th. The improvement continued. The sulphate of copper was given once every four or six hours: sickness followed each dose, with more or less detachment of membrane. His swallowing being now more free, beef-tea and port wine were given largely; and this day he was prevailed on to use the following gargle:

R. Acidi hydrochlorici mxxv ; mellis rosæ ℥ss ; aquæ ℥v . M.

From this time he gradually improved, and would have been soon convalescent, but for considerable sloughing of the external fore part of the throat, that had taken place from the vesicating application used at the onset, so that he was not off my hands, quite well, until August 2nd.

His mother, a young widow, who was in constant attendance upon him, had a smart attack of sore-throat, evidently of the same character, but of a slighter degree; it was removed by the use of hydrochloric acid gargle, with external counterirritation on the throat, and saline aperients.

REMARKS. In the foregoing case, although little had been made public at that time of diphtheria, and no case of the disease had previously come under my notice, I had no doubt that it was a more formidable disease of the throat than I had met with for some time, if I except cases of croup. From the intractable state of the little patient, there was not a good chance of using remedies; and although, from having seen many milder cases since, in which the tincture of sesquichloride of iron has appeared of service, yet I feel certain that the sulphate of copper saved this child's life, as I have frequently found in severe and hopeless croup cases, and which, I think, is worth a trial where other remedies fail. There was in this case no evidence of contagion; scarlatina was not prevalent in the neighbourhood; nor was there affection of any internal organs.

Gulstonian Lectures

ON

FEVER AND INFLAMMATION.

DELIVERED BEFORE THE
ROYAL COLLEGE OF PHYSICIANS, LONDON,
1859.

By WILLIAM ADDISON, M.D., F.R.S., Fellow of the College.

LECTURE II (continued).

XI.—INFLAMMATION AS A THERAPEUTICAL OR DEPURATIVE REACTION, IN CASES OF FEVER.

In some fevers, or in some cases of fever, the natural depurating organs are sufficient; or, by proper medical treatment, they may be roused to a sufficiency for the elimination and discharge of the morbid matter made over to the plasma from the diseased corpuscles. If this can be accomplished, there will be no call or necessity for any preternatural depurative reaction between the plasma and the common tissue. In such case, therefore, the person has, and must go through, the fever; that is to say, the blood corpuscles must pass through the phases of their disorder; but he is saved, by judicious medical treatment, from a local inflammation, because distemperature of the plasma, consequent upon disorder of the corpuscles, is met and relieved by the natural working of the depurating organs. These organs act upon the plasma; and inflammation is an action between the plasma and the vessels. By one or the other, or by both ways, the fluid of the blood may be relieved of hurtful matter: and, as the severity and duration of symptoms of fever are a measure of the severity and duration of disorder in the corpuscles, so, we apprehend, the severity and duration of the consecutive inflammations are a measure of ease or difficulty with which the morbid matter separates, or sloughs off, from the rest of the fluid, and is made over to the common tissue for discharge. But, in cases of fever, we apprehend that neither the natural organs nor inflammation can effect this depurative purpose, so long as disorder is limited to the corpuscles.

The *materies morbi* of the corpuscles—of whatsoever nature this may be—must leave them, and be discharged into the plasma, before any depurative means can come into play for the final expulsion of it from the fluid of the blood. This

appears to be the *rationale* of our inability, by any means which have hitherto been tried, to cut short the progress of a fever.

The argument respecting the therapeutical properties of inflammation in distemperatures of the fluid of the blood was partially discussed in the first lecture; and we now proceed with what further we have to say on this subject.

CASE. At 8 o'clock in the morning of Dec. 28th, a physician, who was assisting at the *post mortem* inspection of the body of a lady who had died of puerperal peritonitis, unfortunately pricked his finger. Twelve hours afterwards, he felt some pain at the part; and he had it touched with nitrate of silver. During the night, shiverings came on, and he felt extremely restless. On the morning of the 29th—the next day—the finger was swollen, and red lines extended up the arm. In the evening of that day, the symptoms were not abated, and there was great prostration. On the 30th, the hand and arm were greatly swollen, the glands in the axilla were affected, and the pain was very great. On the 31st, the pulse beat from 90 to 100 in the minute; and the breathing was heavy and irregular, with torpor and drowsiness. In the evening, all the symptoms were increasing; and now an erysipelatous blush from the axilla extended over the side of the chest. During the night, the breathing became difficult, and the drowsiness passed gradually into deep stupor. Death took place at six o'clock in the morning of January 1st, not quite four days from the infliction of the wound.

In this case, the phenomena, in all important respects, are similar to those observed in traumatic erysipelatous fever, and in puerperal fever. The circumstance of the disease arising in the manner related—namely, from inoculation of a poison from the body of another person who had had puerperal fever—establishes the relation between it and the contagious fevers, and shows that the fatal termination in so short a period is to be attributed, not to inflammation, but to disease of the blood. If this be assented to, the case is taken out of the category of inflammatory diseases, and is placed in that of blood-diseases.

But if this and analogous cases—if erysipelatous fever, puerperal fever, gout, small-pox, and the other exanthematous fevers—be considered as blood-diseases, a great step indeed will have been taken, in the direction we are arguing, towards removing inflammatory reactions altogether from the pathological list; and a wide avenue is opened for a reconsideration of their true import.

Moreover, much doubt is thrown upon the value of the labours of the pathological anatomist, who may regard effects left by internal inflammation in fatal cases of blood-poisoning as showing anything whatever of the nature or seat of the disease. For the question arises, whether inflammation, and the suppurations which may appear in contagious fevers, are not appropriately placed in the same class with inflammation and suppuration in small-pox, sloughing carbuncle, and necrosis of bone; all of which are indisputably therapeutical reactions, the only difference being that, in the one class of cases, the actions arise for therapeutical purposes in the solid parts—the common tissue; in the other, for therapeutical purposes having reference to the fluid of the blood.

If we impartially review phenomena of inflammation as a matter of natural history, and begin with the simplest cases—scalds, burns, sloughs, and fractures (injuries to the common tissue), and boils, eruptions, gout, and small-pox (from injury to the qualities of blood)—we can scarcely fail of perceiving, in both classes, that the forms and amount of the action depend upon, or are governed by, the amount or extent of injury sustained. And, if hindrances protract the process of repair, so also analogous difficulties protract depurative forms of inflammation. If keeping peas in a sore protracts granulation and discharge, so also perseverance in unwholesome articles of food will protract ulceration. In mechanical injuries, the cause of the injury (the heel of the horse, or the cart-wheel), the part injured (the torn flesh, or the broken bone), the extent and nature of the hurt (contusion, laceration, and comminution)—all these, and also the subsequent reaction (the process of repair), are objects either of sight or of touch, or of both. On the other hand, in injuries to the blood, all these things are, and to a great extent must remain, matters of reasoning and deduction. We have said, that very little can be demonstrated of the vital and depurating processes constantly going forward in the blood of the living person. In the engrafted small-pox, it is true, the poisonous matter introduced into the blood, and the consequent inflammation and suppuration in the skin, are seen; but the essential part injured—the elements of the blood—the extent of their injury, and the depurative reactions

going on amongst them, and which connect the introduction of the virus with the fever, and the subsequent inflammation and suppuration, are not seen. Likewise, in the fatal case just related, there was evidence of the introduction of poisonous matter into the blood through the wound of the finger; but the rapidity with which it spread throughout, and destroyed the normal quality of the blood, was only to be proved by the rapid course of the symptoms and the fatal termination. In this case, we argue that the inflammation and swelling of the glands in the axilla were reactionary efforts to arrest the course and eliminate morbid matter, though they failed: and they failed upon the same ground that similar efforts fail in cases of severe mechanical injury; namely, because the injury inflicted on the blood was so great that life succumbed before measures of relief could come effectually into play. In severe mechanical injuries, persons sometimes die before, or soon after, a process of repair is established; so likewise, in injuries to the blood, persons will die before any inflammation appears.

XII.—DISEASES OF THE CORPUSCLES OF BLOOD.

With respect to disease or abnormal metamorphosis of the corpuscles of the blood, and its association with symptoms of fever, numerous observers have described their darker colour, and the easy transudation of their colouring matter in typhus. Denis, for example, describes the blood in typhus fever as deficient in fibrine. He says that air had no effect in reddening it; and, on analysis, it was found to contain ammonia. Dr. Armstrong observed the blood in typhus from the temporal artery as dark as that from the veins.

The general appearance of blood in malignant fevers has been described by Huxham and Fordyce. At the first commencement of symptoms of fever, the blood was sometimes buffed; but the clot beneath was always of a loose texture, scarcely cohering, and very dark in colour. If the patient was bled two or three days after the onset of fever, the blood was found incoagulable, having the appearance as when spirits of hartshorn is added to blood as it runs from the vein, which darkens its colour, and prevents coagulation.

In the yellow fever of the West Indies, blood has been observed to be hotter than in health. As fever progresses, it becomes black and thin. Dr. Blair says that, in many instances, the corpuscles were found so much dissolved that, in several specimens of fever-blood, but few of them could be observed.

It has been found that a diminished amount of carbonic acid is discharged from the blood by respiration in cases of fever; and this fact, taken in conjunction with the dark colour of blood, is conclusive as to one, at least, of the functions of the corpuscles being disordered in fever.

Rokitansky, Simon, Perry, and many other accurate observers, all speak of the dark colour and changed state of the corpuscles of blood in malignant fever, of the incoagulability of the plasma, and of the staining of the tissues and of the serum by the colouring matter which transudes the corpuscles.

As regards the microscopical appearances of the corpuscles of blood in fever, we hesitate at present to lay any stress upon them; because in persons in health, they very speedily change their figure and outline in a very uncertain manner. Some of them become notched with projecting points, and otherwise changed in outline; others assume globular forms; they are influenced as respects these changes, it would appear, by the temperature of the glass upon which they are received for examination, and by the way in which they are covered by the thin glass. Some are seen paler, and some smaller than others; some adhere closely together in rolls, others float about separately without the least disposition to adhere to their neighbours. All these varieties may be seen in the same small quantity of blood which is alone available for microscopical inspection, especially if the exterior edges of the film of blood be brought into the focus of the microscope.

Notwithstanding these obstacles to the drawing any positive conclusion from the microscopical appearances of the blood-corpuscles in cases of fever, it is quite as likely to be by the microscope as by chemistry that future advances in the pathology of these bodies will be made. For how can any bulky chemical manipulation satisfactorily eliminate results from the plasma from results from the corpuscles? especially if, under the influence of reagents, as we shall show, they throw out matter from their interior into the fluid in which they swim, and yet preserve their integrity or individuality. "Hitherto, at all events", as Rokitansky has well observed, "chemistry cannot be said to have excelled, as respects the pathology of these bodies, the achievements of a circumspect anatomical survey,

notwithstanding the limited resources at the disposal of the latter."

EXPERIMENT I. Take a slip of glass, such as is used for mounting microscopical objects, receive on it a very small drop of blood, and place close to it with a pipette a drop of any fluid, chloroform, ale, weak sugar and water, etc., and the quantity of the extraneous fluid should not exceed the quantity of blood. Upon now dropping on the two fluids a thin piece of glass, their nearest edges will mingle, but in various proportions.

In these experiments we find the *outline* and the *interior* of the corpuscles of extremely various appearances in the same experiment, but nothing appears thrown out from them.

EXPERIMENT II. Proceed as in the last experiment, but let the fluid used be sherry wine. The corpuscles, along the line where the blood and wine mingle, will soon begin to throw out molecules around their circumference, many of which pass away into the fluid; others grow out into long tails, which remain attached to the corpuscles, and terminate in a knob. They also wave about in a very singular manner. After some time (half an hour), the tails or filaments become modulated, and then break away from the corpuscles, and when they have done so, they continue their singular movements in the fluid.

Some years ago, when making observations on the plasma or liquor sanguinis of blood, drawn by venesection in cases of fever and inflammation, we observed in the fluid a vast multitude of minute molecules. (*Medical Gazette*, vol. ii, 1841-42.) And the molecules seen in the experiment here related, are precisely the same in magnitude and appearance as those we saw in the fluid of blood drawn in the cases of fever. Now we believe that the blood-corpuscles do not immediately lose their vital or chemical properties when withdrawn from the body. Therefore, we regard the remarkable forms and actions they exhibit under the influence of sherry wine, as phenomena of a species of reaction, which, amongst multitudes of them, is various or unequal, hence the various appearances presented. In some, the resistance offered to the extraneous fluid is more easily overcome than in others.

We have frequently examined with the microscope blood taken from persons affected with scarlet fever, with reference to the appearances in the plasma, and have always noticed the following facts. The colourless or plasma corpuscles are much more numerous than in persons in health, and especially so if the blood be drawn from the skin as the fever is passing off and the epidermis beginning to exfoliate. They are of different sizes and present different appearances. In the open spaces between the rolls of the red corpuscles, irregular masses of granular matter and numerous free molecules are seen floating in the fluid.

In cases of diphtheria, we find that the plasma presents the same appearances as are seen in scarlet fever. Formerly we supposed the free molecules observed in the plasma of blood, drawn in cases of fever and inflammation, to be derived from the colourless corpuscles; but now that we have seen them thrown out by the red corpuscles, there is actual proof that these corpuscles, in their reactions upon extraneous substances, do themselves throw off matter into the fluid in which they swim. The appearances, then, which we have seen in the fluid of the blood in cases of fever, and the behaviour of the corpuscles in the experiment we have related, appear to corroborate our conclusions respecting the excretions of the corpuscles passing into the plasma, and the association of these excretions with phenomena of fever.

Before proceeding with our argument, it will be convenient now to refer to some collateral incidents which require our notice.

First, it may be objected to our proposition respecting the antecedent of fever, namely, disorder or disease of the blood-corpuscles, that in venosity of blood, where a poisonous element of the corpuscles fails of being discharged by respiration, the symptoms are those of brain disturbance, and not of fever. In *morbus cæruleus* there is no fever necessarily.

To this objection it may be replied, that the presence of a venous quality in the corpuscles does not imply disorder or disease in them, in the same way that it is concluded to arise from miasms in the air. The substance—carbon or other matter—which gives to the corpuscles their venous character, is a substance natural to them, an essential ingredient of their composition in certain parts of their course in the circulation. Therefore, it is to be expected, it would occasion no unusual reaction on the part of the corpuscles themselves, though the brain suffers; whereas, a poison from the air may be presumed something quite heterologous to the corpuscles, and doubtless they react with more energy against a foreign injurious

matter, than against anything which is a part of their normal composition.

In cases of blood-poisoning through the stomach, such as drunkenness from alcohol, narcotism from opium, and salivation from mercury, the locality and character of the symptoms point out which organ it is that suffers first or most from a particular poison diffused through the plasma. And if in these examples fever be absent, the argument is, that the parenchymatous organ is affected before the corpuscles of the blood, fever appearing when they partake of the disorder.

In cases of blood-poisoning through the lungs, on the contrary, the symptoms begin with fever, and the argument is, that the corpuscles of blood are here affected first, before the plasma or any of the local parenchymata; fever denoting an abnormal metamorphosis of the corpuscles.

A venous state of the corpuscles of blood is not an abnormal metamorphosis. The condition comprehended in the term venous is one natural to them. Misplaced venous corpuscles disorder the brain; but fever does not appear, because the venous state is natural to the corpuscles. The absence of symptoms of fever, when venous blood circulates arterially, is therefore no valid objection to the proposition we are arguing.

Again, we have hitherto purposely abstained from any reference to the brain or nervous system as causes of fever and inflammation, not because we underrate their powerful influence over blood and the secretions, but because so important a part of our subject requires a special consideration, which we can here but briefly indicate.

It has been argued in the former lecture, that the elements of the parenchymatous organs and the corpuscles of blood have the common properties of other living cellular bodies; and among these are properties of affinity, which differ in relation to different substances. And there can be no doubt that the corpuscles of the blood, as respects the various substances they encounter in their course during circulation, have much more intimate relations with (or a greater affinity for) some than others. For example, in the lung, a special reciprocal action takes place between the corpuscles which are within the vessels and the air which is outside them. And, as if to facilitate this action, the coats of the blood-vessels in the cells of the lung are reduced to an extreme degree of thinness.

Now, it may be argued of any other special organ, where the coats of the capillary vessels are reduced to a still greater degree of thinness than in the lung, that they are so for a similar purpose. Thus, in the brain, an organ very largely supplied with blood, the coats of the extreme vessels are so thin that we fail to trace them in all their various ramifications: indeed, so entirely are they altered that the elements of the organ have but a slight coherency. And, unless the contrary can be shown, we may infer this disappearance of the ordinary properties of blood-vessels to be for the purpose of removing all obstacle to the closest possible contact between the corpuscles of the blood and the parenchyma of the brain. This inference is corroborated by the fact that the brain is the organ specially affected by an abnormal circulation of venous corpuscles. That is to say, the brain is the organ which first detects the presence of an abnormal venous quality in the corpuscles.

If, then, we may regard the thinning away of the coats of the capillaries of the brain as facilitating a contact-action between the corpuscles of blood and the substance of the brain; and if, moreover, we are able to appeal to the known effect of venous corpuscles upon the functions of the brain denoted by drowsiness, stupor, delirium, and coma, as proof of a special action between the brain and the corpuscles, then we may claim the constant occurrence of similar cerebral disturbances in fever as corroborative of the view which bases phenomena of fever upon abnormal metamorphosis of the corpuscles of blood; the brain—to use a chemical phrase—being the test of the condition of the corpuscles.

All the functions of a living body may be comprehended as a series of actions and reactions; and, if it be admitted that the corpuscles of blood do perform a special function in the brain, there must necessarily be reaction from the brain upon the corpuscles.

CASE. A young married woman, aged 19, was delivered of her first child. The labour was natural, and she went on favourably until the fourth day from her confinement, when her husband stayed out late at night, and returned home drunk from a fair, very much knocked about. By this she was thrown into a state of great nervous excitement. Very shortly after, she was seized with a numbness of the legs and shivering, and then with pain in the head and wandering of the mind. The secretion of milk was interrupted; the skin became hot and dry; no

sleep could be procured; and the pulse beat 120 in the minute. The wandering of the mind passed into furious delirium, and all the symptoms of fever and excitement continued for several days. It was only by judicious medical treatment and careful nursing that the disorder was subdued on the seventeenth day.

In this case, we argue that the fever arose, not immediately from the nervous shock, but from disorder of the blood-corpuscles; and, as our researches have led us to interpose distemperatures of the fluid of the blood between errors in diet and inflammation, and disorder of the corpuscles of blood between ærial miasms and fever, so analogously of mental emotions and nervous shocks, when they occasion or aggravate fever or inflammation, we conclude they do so by disordering, or adding to the disorder, of one or other, or both parts of the blood. Errors in diet do not produce inflammation, unless the fluid of the blood be distempered: also a miasmatic air does not produce fever, except the blood-corpuscles be affected; so, too, great convulsions and loss of consciousness (in epilepsy, etc.) pass away without fever, if the blood be not affected. The close sympathy between states of the corpuscles of blood and the brain, then, supports our proposition that symptoms of fever and the generation of poisons in the blood are to be based upon disorder of the corpuscles.

[To be continued.]

Abstract of a Lecture ON THE CLASSIFICATION AND GEOGRAPHICAL DISTRIBUTION OF THE MAMMALIA.

DELIVERED IN THE SENATE HOUSE OF THE UNIVERSITY
OF CAMBRIDGE, MAY 10TH, 1859.

By RICHARD OWEN, D.C.L., F.R.S., etc.

THE learned Professor expressed his sense of the honour which had been done to him by the appointment to the Sir Robert Reade's Lectureship,* and the pleasure it would afford him to give to the members of the University a brief account of some of the prominent features of one of the divisions of the animal kingdom. He had selected the subject of the Classification and Geographical Distribution of the Mammalia, because it afforded an opportunity of taking a general survey of one of the most interesting domains of nature in the short period allotted to a single lecture.

He detailed briefly the principles and peculiarities of the classifications which had been respectively formed by Aristotle, Ray, Linnaeus, and Cuvier. The first of those naturalists had evinced a very extensive and correct knowledge of the mammalian group, by including the cetaceans in it, and by the manner in which he had divided it into the three great classes of BIPODA, TETRAPODA, and APODA. From the time of the publication of his remarkable treatise, *‘Αἱ περί τὰ Ζῷα Ἱστορίαι*, a period of more than two thousand years elapsed without any advance in the subject; and nothing indicated more forcibly the decadence or dormant condition of the human intellect, during a great part of that period, than the dearth of new facts and generalisations in natural history. To Ray, a member of the University of Cambridge, was due the merit of again opening up this interesting field of knowledge, of adding many new facts, and especially of pointing out clearly the distinctions of class, order, genus, and species; but in his generalisations he was, in some points, inferior to his great predecessor, whose classification he in the main adopted. Cuvier had given a great impulse to the science; and, by dint of vast research and unusual capacity for generalisation, had worked a far more complete and satisfactory classification than had hitherto been attained. His classification had been generally followed in the

* The endowment for this lectureship was given to the University in 1524, by Sir Robert Reade, Lord Chief Justice of the Common Pleas in the reign of King Henry VIII. A statute approved by Her Majesty, by order in Council, April 6th, 1858, directs that the lecturer shall deliver one lecture in term time in every year. The lecturer is to be appointed by the Vice-Chancellor during the Lent Term in every year, and to deliver his lecture in the Senate House, on a day to be fixed by the Vice-Chancellor. The present Vice-Chancellor accordingly appointed Professor Owen, being desirous that the office should be held for the first time under the new statute by one who would shed a lustre upon it, and who at the same time is not a member of the University.