MIDDLESEX HOSPITAL.

FRACTURE OF THE RIBS WITH EXTENSIVE EMPHYSEMA :
WITH CLINICAL REMARKS.

By A. SHAW, Esq.

G. B., aged 39, a carman, was admitted on September 10th, 1859, into Middlesex Hospital, under the care of Mr. Shaw, for fracture of several of the ribs on the right side. For many years he has been subject to chronic bronchitis, and up to the present time has been a patient of a Dispensary for Diseases of the Chest. On this day, while unloading his waggon of beer-barrels at a public house, he fell into the cellar, and struck his side with great force against the edge of a barrel. He was brought to the Hospital within an hour after the accident. Three or four of his ribs, in the central and back part of the right side, were broken, their ends being loose and grating. Over the seat of the injury, for a space of about eight inches square, there was emphysema, marked by the swelling and peculiar crackling feel caused by air contained in the cellular membrane. No blood was perceived in the sputa. The dyspnoea was very great. As this was aggravated by the application of the rib-roller, it was removed; and a broad piece of adhesive strapping was drawn firmly over the injured part. The patient being unable to lie down, a bed chair with pillows was provided: he had a purge of calomel and jalap: also a linctus for the cough: and diet of gruel.

September 11th. His difficulty of breathing has increased: the act of expectoration causes great distress: what he brings up is thick mucus. Pulse 82. The emphysema can now be felt all round the chest, over the abdomen, and down the arms. He was ordered a saline mixture, with ipecacuanha wine and chloric ether: and broth for diet.

September 12th. The emphysema has extended to the neck and head, and is now in all the upper half of the body. The face is greatly puffed out, all the lines and wrinkles of the features being obliterated, and the skin tense and glossy; both eyes are quite closed, the eyelids being distended, like segments of round balls, beyond the level of the orbits. Even under the scalp, near the vertex, the crackling of the air can be felt. The scrotum and skin of the penis are very largely distended. The emphysema does not extend to the lower extremities; it terminates abruptly on both sides at the line of Poupart's ligament. The frequency of the cough and painful efforts to expectorate, greatly harass him; the matter brought up is still tenacious mucus. In the intervals of the fits of dyspnoea, the respiration, though laboured, is comparatively tranquil; not being so difficult as to lead to the idea that much air is confined in the cavity of the pleura. As he is unable to lie down, and his body frequently falls forwards, his rest is greatly disturbed; his pulse is weak, and he complains of exhaustion. Brandy, in small quantity, and ammonia, are to be administered when he is faint.

September 14th. He is much the same; the brandy is to be discontinued, as he complains of increased pain in the side.

September 15th. A new source of distress has been added in his having had an attack of diarrhoea; it subsided after a few doses of chalk and aromatic mixture. The emphysema in the face is diminished; he can partly open one eye.

September 16th. The cough is much less troublesome. By substituting numerous pillows for the bed-chair, he has rested more comfortably, and got some sleep. Fish diet was ordered.

September 18th. He goes on favourably. The swelling of his face is much less; he can now partially open both eyes.

September 23rd. His breathing is easy. Since last report, the rib-roller has been applied, and has added to his comfort. He was ordered olei balenæ ʒi, three times a day, after meals.*

September 29th. The emphysema is nearly gone. He is dressed, and has been out of bed for two hours.

October 11th. His cough has almost ceased, and he has gained flesh. He was discharged to-day.

Mr. Shaw has favoured us with the following

CLINICAL OBSERVATIONS, which were addressed to his class. The case presents an example of ribs fractured by *direct* force. That distinction is important, as, in such kind of fracture, the jagged ends of the ribs are driven inwards, and are in danger of penetrating not only the pleura costalis, but, as in our patient, the lung itself.

The signs of the lung being wounded are, bloody sputa and

emphysema. In this case, there was an absence of expectoration of blood. Owing to the elasticity and immediate collapsing of the lung at the wounded part, vessels of considerable size may be opened, and yet the hæmorrhage may not be great.

In emphysema, the air escapes from the wound of the lung during the act of expiration: it is owing to the same compression by which the air contained in the lung generally is expelled through the natural passages, the bronchi and trachea, that the air contained in the portion of lung near the wound is expelled into the cavity of the pleura, through the newly formed outlet, viz., the wound. By degrees, the air which thus issues from the wound collects, in greater or less volume, within the cavity; but it does not long remain there: it is gradually expelled, and that principally by the efforts of coughing. The pressure of the compages of the chest, exerted upon the confined body of air, causes it to escape out of the pleural cavity through the interstices of the fractured ends of the ribs and the lacerated tissues around. In that manner the air gets access to the cellular membrane of the exterior of the thorax; and as the pumping goes on, the emphysema extends. The whole trunk, arms, neck, and face, in our patient, were blown out by the extravasated air. But it was observed that the lower extremities were not affected. The explanation of the latter fact I conceive to be, that, as the patient could breathe only in the sitting posture, and his body was commonly inclined forwards, there were produced at the flexure in the lower part of the abdomen, by that position, a constriction and pressure, which had the effect of preventing the air from descending into the cellular membrane of the lower extremities.

In this case, notwithstanding its absence in the legs, the emphysema was certainly more extensive than is commonly met with. Can you think of any peculiarity in the patient's condition that may account for that? In my opinion, the dyspnoea from which he suffered, in consequence of his chronic bronchitis, had an important effect. The quantity of sputa collected daily in the porringer, you observed, was considerable; it was also expectorated with much effort. We may conclude, therefore, that when he met with the accident, the bronchial tubes were obstructed more or less by tough tenacious mucus. Now, in proportion as the respired air, which fills the lungs in ordinary breathing, is expelled, during expiration or coughing, with difficulty, so much greater will be the tendency of the air contained in the portion of lung near the wound to be emitted in large quantity through that abnormal passage into the cavity of the pleura. Hence the painful and laborious expectoration, and the great and rapid extension of the emphysema, both strikingly observed in this patient, may be justly connected together as cause and effect. I would also suggest, as a reason for the absence of bloody sputa, that just as the air contained in the lung near the seat of the wound could not be expired along the bronchi and trachea, so the blood extravasated into the minute bronchial tubes at the wound could not be expelled through the same passages.

The question may be asked: was the distension of the cellular membrane over such a large extent of the body, and which, especially in the face, had such an alarming appearance, a real source of danger; or was it rather to be considered in a contrary view? That question will be best answered if you reflect on what would have been the consequence if all that air, instead of having been distributed over the exterior of the thorax, had been pent up in the closed cavity of the pleura. How would the patient's breathing have been affected? The first effort would have been a collapse of the wounded lung; it would have been flattened against the mediastinum, and the breathing through it greatly impeded, or stopped. Secondly, if air had continued to be pumped through the wound in the lung, the cavity of the pleura would have become, in the course of a short time, so much overdistended that the heart and mediastinum would have yielded to the pressure—they would have been thrust to the opposite side—they would have encroached on the space pertaining to the sound lung; accordingly, that lung, on which the act of respiration would then have entirely depended, would have been compressed, and incapable of performing its office.

I have witnessed, in a case of fractured ribs, where the emphysema was slight, the dreadful dyspnoea produced in the manner I have described; and also the extraordinary, instantaneous relief given by puncturing one of the intercostal spaces. As soon as the cavity of the pleura was perforated, the air rushed from the orifice with a force that blew out a lighted candle. Accordingly, when we bear in mind to what an excessive degree our patient suffered from difficulty of breathing, owing to the effects of his long-standing bronchitis, it may fairly be

* The oil of the black-whale of the Southern Seas. It has been long used in the Middlesex Hospital as equal in virtue to the cod-oil; than which it is cheaper, and less objected to in taste.

conjectured that if, in addition to that, his breathing had been oppressed by compression of the lungs from air accumulating in the cavity of the pleura, he could not have survived. We may, therefore, look on the escape of air into the cellular membrane, external to the chest, as having been a source of safety—like the steam let off by the engineer, from an overcharged boiler.

In the course of watching the case, we had to keep our minds directed to three distinct sources of difficult breathing: first, that which might depend on the patient's former disease of the chest, of the nature of which we had no exact information; secondly, that which might be the ordinary effect of several ribs having been broken at the same time; thirdly, that which might arise at any moment from the air that escaped at the wound becoming confined, and increasing in quantity within the cavity of the pleura, so as to compress and stop the breathing in one or both lungs.

You will have observed that we applied the rib-roller at first, and immediately afterwards removed it; but again, when the patient improved, returned to its use. It is not unfrequently found, in cases where more ribs than one have been broken without any lesion of the lung, that the chest-bandage, however applied, with or without pasteboard or the like to protect the side, cannot be borne, and we are obliged to abandon it. This is more likely to occur when there has been previous disease of the chest, old bronchitis, or emphysema, in elderly persons, who are the people most liable to have their ribs broken. But, in this case, I was afraid that, if the thorax had been bound firmly, or with tightness sufficient to do any good, two bad results would have ensued: first, it was to be apprehended that the pressure directed on the seat of the fracture by the roller would have stopped the exit of air from the cavity of the pleura, or prevented its free entrance into the cellular membrane; and secondly, supposing air had accumulated in the cavity, and caused compression of the wounded lung, the bandage would have had the effect of interrupting the play of the sound side of the thorax, and, by impeding the respiration in the only lung left to perform the office, would have added greatly to the dangers of the patient, suffering already as he did from chronic bronchitis.

Original Communications.

ON THE SUCTION-METHOD OF REARING INFANTS.

By F. J. BROWN, M.D., Chatham.

My object in this paper is to point out the value of milk as the food of infants, and the importance of administering it by the suck-bottle; also, to condemn the practice of pap-feeding.

I wish to show not only that milk is the natural food of infants, but also that suction is the natural mode of taking this kind of food during the infantile period of life. In speaking of suction, I do not mean exclusively the sucking of the mother's breasts; I also include the sucking of a bottle with prepared mouthpiece. I use the word suction to show that the natural mode of feeding in infants is by sucking, in contradistinction to the eating or swallowing of spoon-food.

Pap of every kind I hold in abhorrence. I declare war against it. I tell the mothers and nurses that feeding by pap is an excellent way to kill their children; that, although the majority of infants live and thrive on spoon-food, yet that many perish; and that spoon-food is the cause of the flatulence, eructations, vomitings, and gripes, that are considered to be natural to infants. I believe that these unpleasant conditions are mostly due to the spoon-food; that they are far less frequent in children that are suckled either at the breast or by means of the bottle, *provided that the acts of suckling be not repeated at intervals shorter than three hours.*

Children thrive admirably on breast-milk, or on cow's milk diluted with one-sixth part of water, and slightly sweetened with white sugar. The amount of sugar should be very small; and the milk should be given lukewarm, with the child lying in the semi-recumbent posture, in its nurse's arms. The child may take its fill (from four to eight ounces) each suckling period: thus it will take from one to two quarts, or even more, of milk and water daily. The child may be taught to do without sucking for four or five hours in the night.

The child should not have anything given it in the way of food until the teeth begin to appear. Nature sends the teeth to indicate that mastication is about to replace suction.

About this period, it is right to give a small quantity of pap; namely, white home-made bread (free from alum), scalded or boiled slightly, and sweetened by a small quantity of white sugar. If such bread cannot be obtained, give tops and bottoms, or rusks, prepared for use in the usual manner. Gradually increase the feeding, and give beef-tea and gravy, and, finally, portions of any wholesome food that may be upon the table.

When the teeth appear late, the feeding must also be adopted late.

Milk must be the food of toothless infants. Still, after nine months have passed from the birth of the child, although there may be no teeth, a small amount of food may be given; but I should advise that it be beef-tea, rather than bread, in these cases. The cordials and baby-mixtures, so commonly used for infants, would almost disappear from our homes, if children were not sought to be reared as young birds. Birds throw food down the throats of their gaping progeny, and nurses imitate them by shovelling pap into the mouths of their little ones until the food lies in the mouth unswallowed, the child scarcely caring to swallow it. Mark the difference in the results: the young bird thrives in direct proportion to the amount of solid food swallowed, whilst the infant thrives in an indirect ratio.

A few words now respecting the sleeping of infants. Infants ought not to sleep in the same bed with their parents: they should sleep in a cot by the side of the bed—one of equal height with the bed. This would save the child from the danger of suffocation, and would render the mother's bed more comfortable for herself. It is believed that many infants perish, not by being overlaid, but by being asphyxiated by carbonic acid; the child sinking down in the bed under the clothes, and so being exposed to the carbonic acid.

The few remarks that I have made contain nothing that is not already known and taught by all medical gentlemen acquainted with sanitary science; but the public are utterly shocked at the heretical nature of the doctrine, and cry out upon its promoters as if they had a design upon infantile life. I have seen so many fine children reared by the suction of milk, and so many children damaged by spoon-food, that I cannot forbear raising my voice a second time* in favour of the one method, and adversely to the other. I believe that thousands of illegitimate children, that are put out to nurse, and that now perish, might be reared by the milk-suction method. Certainly many of these poor innocents are diseased, and die of syphilis; but for this affection in infants there is a remedy that acts most satisfactorily; it is the unguentum hydrargyri fortius of the *Pharmacopœia*, applied over the abdomen. No medicine is required internally. I have seen many children cured in this way of syphilis, and also of strumous glandular disease of the abdomen.

If it were not unsound policy to interfere with the operations of trade—to disorder the balance between supply and demand—I should recommend that adults in the enjoyment of good health should forego the use of milk, so that babes might obtain an abundant supply of their natural food. There are so many weakly mothers, and so many unmarried women that bear children, that a large supply of cow's milk is needed. I expect that all the cows' milk consumed in this country in tea and coffee would be required for infants, if the milk-suction method of rearing children were exclusively adopted. For my part, I should be glad to see the use of milk confined to infants, to the aged, and to the sick of all ages.

To sum up:—

1. The natural food of infants is milk.
2. The natural mode of feeding, in infants, is by suction.
3. Milk must be the food of toothless infants.
4. The teeth are Nature's indices of the masticatory process.

AN ANALYSIS OF TWO THOUSAND CONSECUTIVE CASES IN MIDWIFERY OCCURRING IN PRIVATE PRACTICE.

By G. RIGDEN, Esq., Surgeon to the Canterbury Dispensary.

[Read before the East Kent and Canterbury Medical Society.]

I HAVE been induced to communicate the following analysis of cases in midwifery to the Society, not only as giving the result of an individual practitioner's experience during a period of many years, but as affording in some degree a contrast between the statistics of private practice and those derived from obstetric institutions, where the average of bad cases, from the

* See my paper on the First Food of Infancy, in the "Journal of Public Health and Sanitary Review" for April 1856.