operated upon with the other, and endeavour forcibly to extend the foot. By this means the plantar fascia is tightened. You then introduce a sharp pointed knife on the outside of the fascia, pass it horizontally beneath it, and divide from below upwards. It should be observed that this operation, when the necessity for it exists, should be performed at the same time with the division of the anterior and posterior tibial tendons.

In cases where the increase of the transverse arch is extreme, the fourth and fifth metatarsal bones pressed inwards towards the first and second, before you proceed to divide the tendo Achillis and flex the foot, it will be necessary to unfold the arch, so to speak, and to raise the fourth and fifth metatarsal bones, which will be found partially dislocated at their tarsal extremities; or the patient, after flexion of the foot, will have his whole weight thrown upon the outer edge of the fifth metatarsal bone, and be in a worse condition than he was before the operation. To effect this object, I have used a "double-screw pad" (Fig. 14), having three pads; two small ones, applied

Fig. 14.

under the fourth and fifth metatarsal bones; and one larger, to be applied on the upper surface of the first and second. By means of the screw, a steady continued pressure and counter-pressure may be kept up; in applying which, however, great caution must be exercised to avoid *undue* pressure, as well as pain. With these precautions, the worst cases may be effectually treated.

When the foot has been thus unfolded, to effect which will be found by far the most difficult part of the whole treatment, you proceed, as in the former case, with division of the tendo Achillis, etc.

After the foot is fully flexed, apply a boot and iron, similar to the one represented in Lecture II, which must be worn constantly until the foot has gained strength. In infants, a tendency to rotation inwards of the whole leg is at times present; so that, when the child begins to walk, the foot, although previously in position, is directed inwards and towards the opposite extremity, so retaining the deformed appearance. This tendency ceases of itself at the end of four or five years, if not interfered with. It may, however, be prevented, in the meantime, by the application of irons attached to the boots on the outer side, and extending to the hips, with free joints throughout, and a strap at the upper extremity, passing from one iron to the other, behind the pelvis, by tightening which, the inversion of the foot may be prevented.

The removal of the os cuboides has been recommended and had recourse to, for the treatment of adult varus. It is not only perfectly unnecessary, but a direct and inexcusable mutilation.

UNIVERSITY OF ST. ANDREWS. The next examination for the degree of Doctor of Medicine, will commence on Monday, the 24th of September. Fellows and members of the Royal Colleges of Surgeons of England, Edinburgh, and Dublin, of the Faculty of Physicians and Surgeons of Glasgow, and licentiates of the London Apothecaries' Company, are eligible for examination. Every candidate is required to communicate by letter with Dr. Day, the professor of medicine, fourteen days before the period of examination, and to present himself to the Secretary for registration, on or before Saturday, the 22nd of September.

Jettsonian Lectures

ON

THE OPHTHALMOSCOPE AND ITS APPLICATION.

DELIVERED BEFORE THE MEDICAL SOCIETY OF LONDON.

By HAYNES WALTON, Esq., F.R.C.S., Surgeon to St. Mary's, and to the Central London Ophthalmic Hospitals, etc.

LECTURE II (concluded from page 598).

IN passing to speak of the morbid changes of the *Retina* and *Choroid*, I must adhere to what I said in my first lecture as to conciseness and mere outline, and give but a very brief notice. My time will not admit of more; besides, ophthalmoscopic detail cannot be taught in this way; to attempt it would be to ensure tediousness, and not profit.

We now get a general idea of the vitality of the eye, as we judge of a person's general health and vigour by a look at his face. We now speak of anemia, or of a reddened fundus of the eye, as we have been wont to say the countenance is pale or plethoric, and, as a rule, the indication is equally valuable for generalisation. It is a matter on which we very frequently want to be informed, and a glance will suffice our purpose.

Retina. Perhaps the commonest morbid change in the retina is *Hyperæmia*. The colour of the fundus of the eye is changed, being more red, and sometimes of a dark red. A portion of the optic disc is obscured by vessels, or the whole is covered thinly, so that the disc is yet made out, or thick enough almost to obscure it, and then its position may be more readily traced by the vessels as they pass outwards, but which soon lose their distinctness.

This is a state that always exists in a greater or lesser degree with the train of symptoms called by Dr. Mackenzie "asthenopia", and supposed by him to be a defect of adjustment only. His reasoning is clear and powerful; his conclusion is inevitable, and, but for the ophthalmoscope, every one must have been satisfied with it. It is worth referring to as an example of the fallacy that we may fall into by attempting to solve such subjects by reasoning alone. I may recall the disease to the minds of my hearers by telling them that "asthenopia" is the "choroiditis" of Tyrrell. It is the affection of those who for the most part get their living by minute sight. They are obliged to discontinue with their work, because the vision as it were gets fatigued, and the objects confused or indistinct, and vanish from view altogether. A slight cessation, a wink of the eyes, a look at a distant object, or, in the worst cases, a pause for a minute or two, enables them to continue. It is strange that the indistinctness comes on when the vision is required to be sustained on minute things only, and not when large and distant objects are looked at. But it is not peculiar to the overworked eye. I have seen it in children learning to read, in the idle man of pleasure, and in the vigorous mechanic. I begin to think that it is almost a defective of organisation, rendered worse or apparent under much or peculiar use of the eye, and the general inutility of treatment, short of cessation of that employment which renders it annoying, tends to this conviction. I could fain dwell longer on it.

Among other ophthalmic revelations of retinal changes, are those of *dots*, or *patches of white*, usually seen in the aged, and associated with feeble vision. I do not know that their nature has been made out.

Edema of the retina, rendering the membrane pale and of a yellowish tinge, less shining than in health, and apparently raised around the optic disc, is fully recognised.

Pigment Deposit on the retina, in every form of spot, and in extent from a few dots to almost complete covering, with but small interspaces, is the most remarkable change in this tunic, and not less is the absence of any coexisting external symptoms. I have seen an amount of deposit that seems incompatible with any vision on first examination of a patient for slight feebleness of vision, which is, of course, a physiological puzzle. Many times I have doubted the visual power till I have carefully tested it.

Choroid. The choroid, like the retina, becomes *hyperæmic*;

but the most striking change is *Atrophy*, chiefly manifested by the removal of its pigment; a common consequence of chronic inflammation. The alteration generally begins near the optic disc. At first a yellowish change is seen, and then the orange-coloured choroidal vessels appear, and these become more distinct, ultimately disappear, and leave a white spot, which is often mistaken by the inexperienced for the optic disc, especially if the disc be injected. With the loss of pigment, there is a deposit of grey or deep black spots in other parts, as if the colouring matter was removed from one portion and taken to another. Associated with this, extravasated blood, in small quantities, is not very unfrequent, so that the whole fundus of the eye has a mottled and variegated appearance, and withal a person may be able to read. These alterations rarely exist without some disease of the retina.

Early in my professional career, indeed in my studentship, I suspected that in so-called *Syphilitic Iritis* there was something more than the name at first led me to infer, and this chiefly on account of the subsequent damage, the very defective eye that often remained after the iris had recovered. I was particularly attracted to this in a young woman, during my dressership. The eye seemed perfect when the inflammation was passed, yet she could hardly count my fingers. Well do I remember this being attributed by the gentleman, whose patient she was, to deficiency of active treatment in the first instance, a remark which explained nothing. The evidence of implication of the eyeball by several local and general symptoms were unheeded by me, for I looked on them, as I was taught, merely as the consequences of an inflamed iris, that being the part which the disease primarily attacked, and on which it spent its violence. It was with inexpressible delight, some years after, that I read Dr. Jacob's most valuable monograph on inflammation of the eyeball, and became convinced by it that iritis, so called, was not disease of the iris alone, but general inflammation of the eyeball; and, indeed, who ever saw the iris discoloured and paralysed by inflammation from syphilis, or rheumatism, or struma, without vision being materially impaired? I have ever found that Dr. Jacob's writings stand the test of time, and are less eclipsed by the advancement of knowledge than those of any modern ophthalmic author. Ophthalmoscopic examination does not render them obsolete, but adds confirmation.

It is no longer a secret why syphilitic eyes become spoiled. With but faint implication of the iris, and little symptom of redness and pain, the retina may be spoiled, as we see by the deposit of lymph, and choroidal damage is subjoined by structural alterations. I once gave a wrong prognosis in a case of rheumatic inflammation of the eyeball, that annoyed me for a long time, and created a great desire to discover the cause of an eyeball being so insidiously lost. With but very slight external evidence of implication of the iris, I gave assurance of complete recovery; but, to the great disappointment of all, blindness ensued. This, together with ophthalmoscopic experience, has taught me always to give a guarded opinion, and I may here remark, with regard to prognosis in general, that most of us are too apt, in critical cases, at a great risk to our reputation and comfort, and without any adequate advantage to ourselves, to be too prophetic, and to promise too much. It is true that patients almost force us into it, and when we commit ourselves, it is our desire for a favourable issue that betrays us.

A more extended knowledge of syphilitic disease of the eye, has convinced me that the posterior part of the eye, certainly the retina and the choroid, and perhaps also the vitreous humour (which is indicated by haziness and flocculi), are always involved. I never saw an example without impaired vision, and all observers know that vision may become misty prior to any palpable iritis. Before I ever looked through an ophthalmoscope, I made the vision the chief ground of diagnosis in the incipient stage of the disease. Many a case of reddened eyeball presents itself, and looks so like iritis, that the most experienced observer cannot say at once that there is or is not this affection; but the test that never fails me is trying the seeing power, and if I find no deficiency here, I am sure that there is only surface inflammation.

A surgeon in very large practice, in London, and well known to all of us, sent very lately for me, to see a patient whom he had attended with rheumatism, and now, as he supposed, afflicted with rheumatic iritis. The eyeball was much injected, and whether superficially or not, I could not tell. The iris was clear. I tried the sight, and when I found that the almost microscopic inscription on my watch could be read, I was satisfied that there was no iritis—no "eyeball-itis". The calomel

and opium pill, which was just commenced, was left off, the usual diet was allowed, and an astringent lotion given. In a few days the eyes were well.

More than all this, the ophthalmoscope teaches us that the retina may be the sole seat of syphilitic action. My attention was first drawn to this in a medical pupil saturated with secondary syphilis, and who lost the sight of one eye without any external symptoms. Internal inspection, however, showed that there was inflammatory exudation over the fundus of the eye, obscuring the optic disc and retinal vessels. When I was last consulted, fortunately for him, his sight was almost clear.

Among the appearances at the fundus of the eye, after the acute stage of syphilitic eyeballitis has passed away, are dots and patches of deposit of different forms and sizes, more or less darkened with pigment, over the retina. The optic disc is often changed, sometimes contracted and ill defined, and greyish; sometimes, apparently, more spread out, and presenting many abnormal aspects. The choroid becomes patchy with white, yellow, and brown; its pigment is lost in spots, allowing the vessels to be seen, and exposing the sclerotic coat.

I must beg the gentlemen who honour me with their presence not to suppose that the ophthalmoscope has increased the difficulties of ophthalmic practice. I fear that this idea is not uncommon, on the contrary, it renders it easier, after a man has taken the requisite trouble to learn certain preliminaries. What time used not to be spent in investigating cases of impaired vision, to discover to which of the subdivisions of amaurosis it belonged? How numerous the questions, how tedious and unsatisfactory the examination? Now we at once have a peep, and see things as they are, and when we are puzzled, the patient is not made to suffer for it. I believe, too, that the ophthalmoscope will reduce ophthalmic literature; the treatises will devote less space to the diseases of vision, because less is needed. There will be more fact, which is always put in a small space, and less of wordy theory.

There is just one topic, respecting therapeutics, that I venture to introduce. Do not abandon a patient, and say that he is incurable, merely because you see this or that change in the eye; it is one of the dangers of a speciality. Let nothing short of disorganisation prevent you from seeking indications of treatment, and catch at the shadow of a chance. We know so little of the natural reparative power, of function and of vitality, that we ought not to dare, except under remarkable and well recognised conditions, to give a fatal prognosis.

Do not look on a diseased eye merely as such; remember that this organ is but a part of a whole, and that recovery, or partial recovery, often ensues contrary to the expectation of the most experienced. Which of us has not seen this exemplified in disease of the lungs, and kidneys, and liver, by specialists in these departments?

I have brought here tonight patients whose diseases will illustrate most of the morbid affections that I have described.

Allow me to thank you, Mr. President and gentlemen, for the attention I have received from you while giving these two meagre lectures. Soon, I shall give to the profession, in the new edition of my work *On the Diseases of the Eye*, a full and complete chapter on all that practically relates to the ophthalmoscope, and to which I beg to refer you for much that I could not even allude to here.

THE NEW CENSUS ACTS. The new Acts for taking the Census in England and Ireland next April have been issued. In England the Secretary of State is to superintend the taking of the Census. Every registrar's sub-district is to be formed into enumerators' divisions. Enumerators are to be appointed, and householders' schedules to be left at all dwelling-houses, to be filled up as to all persons abiding therein on the night of Sunday, the 7th of April, as to the name, sex, age, rank, profession, or occupation, condition and relation to head of the family, and "birthplace of every living person", and also whether there were any blind or deaf and dumb. There is nothing as to religious profession in the English Act, but there is in the one relating to Ireland. The schedules in England are to be collected from the houses on Monday, if possible, the 8th of April, and to be corrected if found to be erroneous. The masters of gaols, workhouses, hospitals, etc., are appointed enumerators for the occasion, and all overseers and other parochial officers are bound to act as enumerators. Returns are to be obtained of the houseless poor during the night of the 7th of April, and of all persons travelling or on shipboard.