Letters to the Editor

We are writing to express our concerns regarding the recently published article by Kumaran et al. (2005), entitled "Structure and mechanism of ADP-ribose-1"-monophosphatase (Appr-1"-pase), a ubiquitous cellular processing enzyme." In this article, the authors describe the crystal structure of the yeast protein YMX7_YEAST (product of ORF YMR087w) alone and in the presence of ADP-ribose (Appr). The authors claim that this protein is the yeast Appr-1"-p phosphatase, and propose a mechanism for the phosphatase.

We note that there is no evidence in the article by Kumaran et al., or elsewhere in the literature, demonstrating that YMX7_YEAST (YMR087w protein) has the claimed Appr-1"-p phosphatase activity. On the contrary, we showed previously that GST-YBR022w co-purifies with Appr-1"-p processing activity (Martzen et al. 1999), and recently showed that YBR022w protein (now called Poa1, phosphatase of Appr1"p) has Appr-1"-p phosphatase activity (Shull et al. 2005). Poa1, like YMR087w protein, is a member of the large A1pp family, as defined in the SMART database (<http://smart.embl-heidelberg.de/>) but, to our knowledge, is currently the only member of this family of proteins with demonstrated phosphatase activity. Other members, such as YMR087w protein, may in fact be phosphatases that can act on Appr-1"-p, as implied by the article by Kumaran et al.; indeed, our data suggests there is another significant source of this activity in crude extracts. However, the possibility that YMR087w protein has Appr-1"-p phosphatase activity is distinct from the declarative statements in the title and text of the Kumaran et al. paper. An alternative possibility is that the shared property of members of this family is their binding of ADP-ribose, as we have determined for Poa1 (K_i , 17 μ M), and as demonstrated by co-crystallization of YMX7_YEAST (YMR087w) and ADP-ribose.

[Kumaran, D., Eswaramoorthy, S., Studier, F.W., and Swaminathan, S. 2005. Structure and mechanism of ADP-ribose-1"-monophosphatase (Appr-1"-pase), a ubiquitous cellular processing enzyme. *Protein Sci.* **14**: 719–726.]

Martzen, M.R., McCraith, S.M., Spinelli, S.L., Torres, F.M., Fields, S., Grayhack, E.J., and Phizicky, E.M. 1999. A biochemical genomics approach for identifying genes by the activity of their products. *Science* **286**: 1153–1155.

Shull, N.P., Spinelli, S.L., and Phizicky, E.M. 2005. A highly specific phosphatase that acts on ADP-ribose 1"-phosphate, a metabolite of tRNA splicing in *Saccharomyces cerevisiae*. *Nucleic Acids Res.* **33**: 650–660.]

Neil P. Shull, Sherry L. Spinelli, and Eric M. Phizicky
University of Rochester School of Medicine
Rochester, New York, USA

The yeast proteins specified by YMR087w and YBR022w, members of the A1pp family, were selected by us for structure determination in a structural genomics approach (www.nysgrc.org) because no structural model was available for this family of proteins. The structure of the protein specified by YMR087w, expressed from the cloned gene, was completed in January 2003. Since the family was annotated as Appr-1"-p processing proteins, we also determined crystal structures with bound ADP and ADP-ribose (PDB 1TY8 and 1TXZ). These structures, plus modeling of Appr-1"-p in the active site, are consistent with the detailed and specific catalytic mechanism for Appr-1"-pase we presented, based on the alignment of functional groups in the structure. This structural interpretation seemed to us a sufficient basis to conclude that the protein has this activity (although we do refer to it as a putative function in the abstract).

D. Kumaran, S. Eswaramoorthy, F.W. Studier, and
S. Swaminathan
Brookhaven National Laboratory
Upton, New York, USA