followed by temporary inflammatory swelling, which, however, soon subsides.

The average interval I have allowed to elapse between each operation, has varied from five days to a fortnight, and the quantity of blood abstracted from one to three drachms.

Smearing the surface with vaseline, and protecting from the outer air, especially in cold weather, is the only additional treatment needed, besides such internal remedies, saline aperients, alteratives, etc., as the general condition of the patient may indicate. I may add, I have seen no complication, or untoward symptom, result from this method. The use of alcohol must be emphatically denied.

I repeat, in all instances, lasting and satisfactory results have been procured, and are attributable to the local depletion, which allows recovery of tone of vessels, the contractibility of which has not been permanently destroyed, and also to the invisible scarring which, by after-contraction, reduces the vascularity of the skin. This after-scarring is hardly discernible.

I regard this method as capable of preventing cellular, and fibro-cellular, hypertrophy, which results from long continued passive vascular congestion of the nose.

CASE I.—Elizabeth W., aged 42, single. Duration, five years. She had frequently taken spirits for alleged "weak heart." Debility, dyspepsia, and dysmenorrhœa. Total, 15,750 punctures, distributed over fifteen sittings, and a period of seven months. Commenced January 1883. Result permanent and satisfactory.

CASE II.—A surgeon, suffering from acne rosacea and vascular hypertrophy. Duration, eighteen years; commencing at the age of 15. He had suffered from atonic dyspepsia. In twelve separate sittings, extending over a period of seven months, 11,150 punctures; commenced January 1882. Result permanent and satisfactory.

CASE III.—Patrick P., aged 50. Plethoric; health good. Duration, six years. Very abstemious. Total, 52,200 punctures, included in twenty-two separate operations, over a period of five months. Commenced December 1882. Result to present, permanent and satisfactory.

A POINT IN THE TREATMENT OF PURULENT OPHTHALMIA.

Read in the Section of Ophthalmology at the Fifty-Second Annual Meeting of the British Medical Association.

By EDGAR A. BROWNE, M.R.C.S. Eng.,
Surgeon to the Liverpool Eye and Ear Infirmary.

In purulent ophthalmia, the indications of treatment are (1) to wash away the infective material as thoroughly and as early as possible; (2) to render the conjunctival epithelium and secretions as nearly as may be aseptic.

Too much stress cannot be laid on the necessity for securing the most perfect fulfilment of the first of these conditions. On this head, all practical surgeons are agreed. The intensity of an attack is greatly modified by the period at which the first thorough cleansing takes place. We frequently have the opportunity in gonorrhœal infection of forming a rough estimate of the virulence of the discharge by inspection of the penis, and it must have occurred to most of us to have noticed that cases in which the urethral inflammation has been at its height have, when cleansed immediately on discovery, been less violent in their course, and less disastrous in their results, than milder cases which were at first neglected. But the configuration of the upper lid renders thorough cleansing extremely difficult, and perfect antiseptics is, of course, impossible. The main difficulty lies in reaching the upper sulcus. If, in a case of moderate severity, without much swelling, we evert the upper lid, and carefully press the conjunctival *cul-de-sac* into view, by means of a probe, we often find, even immediately after a dressing, that this important portion of the mucous tract has neither been effectually cleansed by the washing, nor affected by the astringent. In other words, the strongest focus of infection receives the weakest treatment. To obviate these disadvantages, I have employed a lid-irrigator, which enables us to throw a stream of pure or medicated water into the sulcus for any length of time we please.

The instrument resembles an ordinary lid-elevator, but is rather deeper in the claw, and constructed of hollow plated tubing, instead of wire. It is furnished above with a handle, and behind a small curved nozzle is connected by means of an India-rubber pipe, six or seven feet long, with a reservoir capable of containing about a quart of solution. The limb that passes beneath the upper eyelid is pierced

with six fine holes, through which the fluid issues as a douche. A stop-cock, or spring-clip, is provided to regulate the flow.

In use, the reservoir is filled, and hung up. The surgeon or attendant stands behind the patient, who holds his head slightly inclined over a basin. The elevator is gently insinuated under the upper lid, and the stream turned on.

The points requiring attention are these: 1, it is necessary to ascertain that the orifices are patent; they are liable to become clogged; four in full play are sufficient; 2, the reservoir must be well elevated to give a good fall, and it must be nearly full to give a good pressure. I generally have a cord run over the curtain-pole, or a picture-nail, so as to get the greatest elevation the room will allow. The jets should be about a foot high when the instrument is held free from eyelid. If the play be not free, we do not attain our object. 3. The application must be prolonged; a quarter of an hour for a first dressing is the minimum I should allow myself. In severe cases, a longer period is advisable. The points aimed at are: continuous flow and prolonged soakage. Now, as to the fluid to be employed. An antiseptic that shall be effective, and yet not too strong; a tissue-irritant is required. I have tried a great number of substances at one time and another, but latterly my inclinations have set in rather strongly towards a preparation of trichlorphenol. I was first attracted by an account, in the *London Medical Record* for April 5th, 1883, of trichlorphenol as an antiseptic, by Dr. Dianin of St. Petersburg. He recommended a solution of trichlor-phenolate of calcium as an application in erysipelas. But the calcium solution is not fitted for eye-work; it deposits a precipitate; it is not easy to make or preserve at an uniform strength; and some specimens have unaccountably changed colour. With the assistance of Messrs. Clay and Abraham, I have managed to attain a satisfactory solution with a magnesia base. It can be made of a uniform strength of 5 per cent.; it is stable and free from particles.

Trichlorphenol—that is, carbolic acid with three atoms of hydrogen replaced by chlorine—is said to be twenty-five times more powerful as an antiseptic than carbolic acid. I have no means of judging of the truth of this statement. Practically, I find a two per cent. solution amply strong enough for the first applications in gonorrhœal ophthalmia, and one per cent. for average use. It gives rise to a good deal of smarting, which ceases directly the flow is stopped. Some skins are easily irritated by its flow over the cheek; the majority are not affected. The skin may be protected by vaseline if necessary. I use it twice a day, for about a quarter of an hour. In one very severe case I have used it three times a day. As the case progresses, a one per cent. solution is sufficient; in some, a half per cent. In order to test the antiseptic value of the solution, I asked Dr. Alexander to allow a certain number of cases of gonorrhœa to be treated in the Lock Hospital of the Liverpool Workhouse. The method of irrigation was employed by means of an elastic catheter passed into the urethra, and an elevated reservoir. Twelve cases were treated, with an average of nine days' treatment; the longest case occupying eighteen days, the shortest three, one four days, and three of six each. The cures were complete. Some credit is doubtless due to the method of irrigation.

This is, I think, a sufficient test to warrant me in recommending this solution for further trial. No remedy, no method of treatment, can have its real value appraised till it has passed through many hands.

As regards the value of the method of prolonged irrigation, I have no doubt; and I feel justified in recommending the little instrument shown as a convenient and efficient means of employing it.

THERAPEUTIC MEMORANDA.

CUCAINE IN DENTISTRY.

CUCAINE, in my hands, 24 per cent. solution, has been quite valueless in preventing the pain attending tooth-extraction. Applied to the gum several times, at intervals of some minutes, it has partially saved the patient the pain of adjusting the forceps; but the actual extraction, involving, as it does, the fracture of the alveolus and rupture of the vessels and nerve at the apical foramen, has been accompanied with the usual pain attending it. Applied to sensitive dentine, it is not at all certain in its action; and to an exposed, ulcerated, or inflamed pulp, not nearly so effectual as our old remedies, carbolic acid and creasote. The oil of cloves, with which the cucaine appears to have been mixed, will account for the success attending its application in the case reported in last week's JOURNAL.

MORTON SMALE, Dental Surgeon,
89, Seymour Street, W. Westminster Hospital.