ment. It is conveyed by contagion, and may also arise *de novo*, from, I believe, a mixture of animal and vegetable matter in decomposition under certain circumstances; but there is little, if any, evidence to show that it is caused by sewage simply, or by animal or vegetable decomposition alone.

EXCISION OF THE RECTUM.

By D. LOWSON, M.D., M.R.C.S.,
Surgeon to the Hull Dispensary.

MUCH has been done, of late years, in the way of improving and perfecting the operation of excision of the rectum; and yet many surgeons are doubtful of its advantages, and prefer either inguinal or Callisen's colotomy. French surgeons, in particular, have had untoward results; and, with loud voice proclaiming their disapproval of the radical method, have advocated various modes of alleviation and compromise. The Germans, on the other hand, foreseeing the advantages likely to accrue, have quietly set themselves to render it easier of performance, safer from risk, and more satisfactory as regards the conservation of the action of the sphincter. In England, the name of Harrison Cripps is well known in this connection, and his researches into the nature of the stenoses and obstructions in this region have led to a more scientific discrimination of the cases fitted for operation.

"Whether excision or colotomy," is a question which must be mainly settled in each individual case; but I venture to formulate the following rule: that when the finger, pushed through the stricture, reaches the upper limit of the ring, or mass, *in front*, excision, and not colotomy, should be resorted to. When the disease extends higher, each case must be decided on its own merits; but of this there is no doubt, that colotomy is not, even in the best hands, a safer operation than extirpation. Indeed, statistics (according to Bryant) are more favourable for the latter operation. The advantages of colotomy are ease and rapidity of performance, as well as an absence of disease in the tissues cut through. *Per contra*, there is the danger of setting up peritonitis, cellulitis, etc. There is also the inconvenience of having the opening into the bowel in an unnatural situation; and, when all has been done, the disease itself is left. Extirpation is more tedious and difficult; but it has the great merit of being an attempt, at least, to remove the disease radically; the anus is retained in its normal position; and considerable, if not at all times perfect, sphincter power remains.

There are several ways of performing this operation, the later modifications aiming at the preservation of the sphincters. Where the anus and skin are much involved, it is common to make a circular or elliptical incision, so as to include all the disease, to separate the mass from its connections in the pelvic outlet, divide the bowel above the affected part, and stitch it to the integument. In annular strictures, an ano-coccygeal incision may suffice, though it is not a good one in most cases. Hüter of Greifswald recommends a horse-shoe incision (Fig. 1), the curve of the shoe encircling the front of the anus, and the ends terminating in the ischio-rectal fossæ. The flap thus formed is thrown backwards with half an inch of rectum, to be replaced after removal of the disease, and the two pieces of bowel are stitched to each other. In the case to be related, I adopted an incision as in Fig. 11. Here the curve extends round behind the anus, and the

Fig. 1.

Fig. 2.

limbs, passing forward and well outward, terminate considerably to one side, and a little in front, of the bowel, and, from the centre of the convexity, a straight cut is carried backwards over the dorsum of the coccyx and sacrum.

Cancer or cancerous stricture is the disease for which excision is most frequently required. In cicatricial stenosis, the bougie or incision, followed by dilatation, is the practice usually followed, or the linear rectotomy of Verneuil; but, "in all cases," Bryant remarks, "there comes a time when treatment by dilatation ceases to be beneficial." He then proceeds to recommend colotomy. Why not excision? In the following case it had, unfortunately, been delayed rather too long.

A. M., aged 37, had for nineteen years suffered from fistulæ *in ano*, and, though twice operated upon, they did not close. Nine years ago, Mr. Brewer of Huddersfield found him suffering from stricture of the rectum. Since then, he has had little peace or comfort, suffering much from irritation and inflammation in and around the bowel; narrowing

and flattening of the extruded fæces, if constipated; or acrid liquid discharges, mixed with small scybala. During the last twelve months, when under Dr. Hanson of Huddersfield, the symptoms, though alleviated as much as medical treatment could, latterly became aggravated; and, when he came to be examined, he was worn and a good deal wasted with pain and anxiety about his state. Obstruction had become worse, and defæcation more difficult and exhausting. There were two fistulæ discharging matter near the anus—one opening through an old abscess, and the other directly into the rectum. The finger could with great difficulty be pushed through the stricture, which commenced half an inch above the end of the bowel, and the finger-tip just reached its upper limit in front, where it ended rather abruptly. Behind, it extended higher; the passage was crooked, but the lining felt healthy, and there was no approach to nodulation, while a tough mass of cicatricial tissue filled the whole pelvic outlet.

The patient's weakness was the only indication against the operation, and it did not seem to be extreme; and, as he was aware his life could not be much prolonged, he was anxious to go through whatever was deemed fit. Extirpation was decided upon, and carried out with the assistance of Dr. Hanson, Dr. Coutts of Waterfoot, and Mr. Macnab of Hull. An incision was made as in Fig. 11, and the flap dissected forward with half an inch of rectum. A straight cut was next carried back over the dorsum of the coccyx, which was separated by a bone-pliers. The mass was then separated from the front of the sacrum, peeled off the obturator internus muscle, and, with a catheter in the bladder, to indicate the position of the urethra, the separation was completed in front by the *écraseur*. The dissection was done in the main with the finger, with occasional aid from Richardson's saw-scissors. The mass, being now loosened all round, dropped down, bringing into view a healthy but muscular rectum, crossing the anterior surface of which was a distinct white line, marking the reflection of the peritoneum. A surprising amount of healthy rectum lay between this and the diseased mass—about an inch. The atmospheric pressure had carried up the recto-vesical pouch, in the same way as Sims' speculum acts in vaginal examinations. The *écraseur* divided the bowel without hæmorrhage; and, in fact, very little blood was lost at any stage, two small perineal branches only being secured. The stitching together of the two pieces of bowel proved a troublesome operation, the more so from the section of the bowel being rather oblique. Drainage-tubes were then introduced, and the external incision closed.

The patient revived wonderfully after a very tedious operation, and in the evening seemed quite lively; but the pulse was small and rapid, beating at 120. Next day, he kept up well. The pulse, however, did not moderate, and the thermometer stood at 100.5°. This was the highest temperature reached. On the morning of the second day, he was very weak and collapsed, and hiccough had proved troublesome all night; the temperature of the extremities was low, and a cold damp moistened the skin. The wound was healthy-looking, but inactive, and the discharge rather copious, and of a serous nature. In the evening, he remained in much the same state, the hiccough being still very harassing. The third day found him much worse, and about noon he was evidently sinking. He died seventy-two hours after the operation.

The cause of death in this case was evidently shock, reaction having been weak from the first. Had the patient made up his mind sooner to have "something done," there is considerable reason to believe the result would have been different, and it is questionable if colotomy would have been more successful.

The incision employed in this operation has, I think, certain advantages over Hüter's. It is more suitable for the removal of the coccyx, which is a considerable gain in a case like the above, where a large mass has to be taken away; and I am not aware that the absence of the bone in question is in any way detrimental. Professor Esmarch of Kiel makes an excellent suggestion in this connection. He recommends that the lower piece of bowel should be removed altogether, and that the upper portion should be stitched to the margin of the anus. This is an improvement in more ways than one. By drawing the bowel through the central opening of the flap (anus), the process of stitching is transferred from the inner to the tegumentary surface, rendering it easy, and uncomplicated with numerous long threads. But of greater importance than this is the advantage in healing. When the two pieces of bowel are stitched above the anus, we have a circular wound, through which the bowel-contents have subsequently to pass; and, unless it heal quickly by first intention, which it rarely does, the connection is broken the first time the bowels are moved, and the fecal matter finds its way through the wound, causing a fistula, which is a long time in closing. I had an experience of this sort, where an annular stricture was removed through an ano-coccygeal incision. The bowels could not be kept constipated for more than five days, and

the result was as related. It is (now) six years since that operation took place. The patient is now well, having good sphincter power, except in diarrhoea; but the healing was tedious and troublesome. Any one who has had experience in excision of the rectum will, I am sure, appreciate Professor Esmarch's suggestion.

A NEW FORM OF VAGINAL PESSARY.¹

By J. G. SINCLAIR COGHILL, M.D., F.R.C.S., VENTNOR.

A CASE of retroversion of the uterus, presenting conditions of unusual difficulty, first suggested the idea of the instrument here delineated.

It is on the plan of the familiar trestle used by builders for supporting stages or platforms, and is made of soft caoutchouc enclosing a steel spring-wire frame. The pessary consists of a transverse bar superiorly, and lateral fenestrated supports or wings. These latter are inclined outwards from the points of their attachment to the transverse bar, and also follow a curve downwards and forwards, corresponding

as nearly as may be to that of the vagina. The wings are also everted anteriorly, not only to accommodate the distension of the bladder, but to prevent the lateral rotation of the instrument. The slight concavity of the upper surface of the bar seems to give more fixed support in cases where there is great uterine mobility, or greater than usual

weight of uterus, but it may readily be bent so as to become convex if need be. The bar is intended to be placed in the vaginal cul-de-sac (*forix*), either behind or before the cervix uteri, according to the nature of the displacement, while the wings rest on each side of the pelvic floor.

This pessary is very easily introduced, on account of its comparatively small size and the compressibility of the wings. The free curved extremities of the latter are approximated closely between the forefinger and thumb of the right hand, and the bar is passed through the vulva with its long axis antero-posteriorly; then a quarter of a turn, with gentle sustained pressure, brings the wings with their curve forwards within the vagina, and, the hold being relaxed, they separate, and their

inferior curves at once adapt themselves to and rest upon each side of the floor of the pelvis. It is only then necessary to pass the forefinger up between the wings that adjust the bar behind or in front of the cervix, as desired. I have found that a pessary of this shape will give more efficient support than a Hodge's, or Albert Smith's pessary quite two-fifths longer in the vertical axis. It is necessary to bear this in mind in the selection of the necessary size. In very many cases this question of mere size is a most important consideration.

The exact *modus operandi*, even of the familiar Hodge type of pessary, is still a moot point among gynecologists; but the action of this trestle pessary is quite simple, and readily enough understood. Like every other form of vaginal pessary, it can only influence the position of the uterus indirectly, through its ligamentous connections. The uterus has, of necessity, a considerable range of physiological displacement; and indeed, so long as its utero-sacral attachments or broad ligaments do not permit a material displacement downwards, an amount of anteversion or of retroversion, quite pathological strictly speaking, may be borne without giving rise to any distressing symptoms. The transverse bar of the trestle pessary, when placed either in the posterior or in the anterior vaginal fornix, gives a point of support to the cervix, at the point of its junction to the body of the uterus, and at the same time raises it with its broad ligaments, restoring the normal tension of the latter, the loss of which is so important a factor in determining the lesion. The restored equilibrium is maintained by the lower segments of the wings resting on the sides of the pelvic floor, above the tuberosities of the ischia, where there are no sensitive or mobile viscera. The bladder and rectum are thus enabled to distend and contract in the discharge of their functions, without impediment on their part, or disturbance of the pessary.

As I have already said, this instrument was designed to meet the exigencies of a case of retroversio uteri presenting local and reflex symptoms of extreme urgency; among others, persistent and painful menorrhagia with intense nausea and gastrodynia. The patient was an unmarried lady. The displacement was of many years' standing, and complete, the fundus resting in Douglas's pouch, and easily recognised through the upper posterior vaginal wall, while the os uteri was tilted upwards and forwards, behind the symphysis pubis. The whole uterus was curved backwards, but not completely flexed. The situation was further complicated by enormous congestive hypertrophy of the dislocated organ, the sound measuring nearly six inches of cavity, and by the presence of a fibroid in the left posterior wall of the fundus, by which, probably, during a period of laborious nursing, the malposition of the organ was originally determined. After reposition, which was a work of extreme difficulty, principally on account of adhesions, every variety of pessary of the Hodge type was tried by eminent gynecologists, both of Edinburgh and London, as well as by myself; but the relief afforded by these aids was only temporary, and never complete even when the instrument was kept *in situ*, on account of the weight and extreme mobility of the uterus, while the frequent necessity for readjustment was a source of the greatest trouble and distress. The difficulty with the Hodge's pessary was the impossibility of a sufficient fulcrum for the lower bar, so as to support the weight above. This lower bar was continually slipping forwards across the vulva, while the fundus uteri was allowed to fall again into Douglas's pouch, over or by the side of the upper bar. It became evident, after much consideration, that a fixed support on the pelvic floor was the only means of meeting the indications of the case; and to obtain this I had the pessary I have described made for me by Messrs. Maw and Co. The effect of its introduction was immediate and complete; and, as far as nearly four months' experience can enable one to judge, promises to be permanent. Indeed, the results far exceeded my most sanguine expectations. The menstrual and other functions resumed their normal character, all the distressing reflex disappeared, and the patient, who had to be carried up and down stairs for the most part, and could hardly walk or drive without extreme suffering, was at once enabled to move about freely, and is rapidly recovering her lost health.

I have since had the opportunity of employing this pessary in several cases, both of retroversion and of anteversion, and also in a case of prolapsus with great protrusion of the bladder and posterior wall of the vagina of very great difficulty, and with results so satisfactory, that I am sure it is an instrument that will be found useful in a large number of cases in which the weight of the uterus, extreme relaxation of its ligamentous connections, deficiency of perineum, or irritation of the bladder or other surfaces, necessitate a more fixed support than can be afforded by other forms of pessary acting more or less on the lever principle, with a movable fulcrum. Indeed, I venture to think that it may be used with advantage, even in a much wider range of cases than those I have indicated.

¹ This pessary is made in three sizes, of which the drawings are exactly to the scale of the largest.