THIRTEEN CASES OF HYSTERECTOMY, WITH REMARKS ON CARBOLIC ACID SPRAY IN ABDOMINAL SURGERY.

By THOMAS KEITH, M.D. Edin.,
Surgeon for Ovarian Diseases, Royal Infirmary, Edinburgh.

In the BRITISH MEDICAL JOURNAL of December 8th, 1883, notes are given of all my cases of supravaginal hysterectomy, twenty-five in number.

During the past year, other thirteen cases have been operated on, bringing the number up to thirty-eight hysterectomies, with three deaths. Besides these, both ovaries were removed twelve times to check the growth of bleeding fibroids. These were successful operations, more or less benefit having in every case already resulted. In only one has menstruation continued. In that case, a fragment of ovarian structure was left closely adherent to the tumour, and over that part there is still pain at intervals. The tumour is, however, much smaller, and the patient's general condition is better than before. In every case one menstrual period came on, generally beginning on the third day. In upwards of sixty operations of double ovariectomy that I have performed for disease, in only a single instance, to the best of my knowledge, has menstruation recurred regularly, and in that case, also, part of one adherent ovary was left behind.

Thirteen Cases of Hysterectomy.

		Date of Operatn.	Age.	Weight.	Description of Tumours, Adhesions, etc.	Result.
26	Infirmary	Dec. 1883	38	7 lb.	Ovaries could not be removed; broad ligaments opened up	Recovered
27	Infirmary	Feb. 1884	43	10½	Bleeding fibroid; adhesions; much enucleation	"
28	Dr. Woolston	Feb. "	39	1	Bleeding fibroid; ovaries could not be removed	"
29	Infirmary	Feb. "	43	4	Subperitoneal; very extensive enucleation; ovaries could not be removed	Died
30	Infirmary	May "	58	13	Solid fibroid; entire growth since menstruation ceased; whole uterus removed	Recovered
31	Dr. Parker	July "	29	19½	Broad ligament opened up; much enucleation; adhesion to colon	"
32	Dr. Somerville	July "	34	12	Omental and intestinal adhesions; much enucleation; broad ligament opened up	"
33	Infirmary	Aug. "	46	9½	Solid tumour	"
34	Infirmary	Oct. "	44	9	Solid bleeding fibroid	"
35	Dr. Rodger	Oct. "	29	17	Omental adhesions; solid fibroid	"
36	Dr. Miller	Oct. "	28	13	Fibrous-cystic tumour; entirely subperitoneal	"
37	Infirmary	Nov. "	34	16	Firm and extensive adhesions to wall, omentum, stomach, and diaphragm	"
38	Infirmary	Nov. "	31	2½	Bleeding fibroid; ovaries could not be removed	"

In the cases of the small tumours—that is, tumours under nine pounds—the operation was begun with the intention of doing no more than removing the ovaries. In nine of the whole number, I failed to find one or both, or, having found them, was unable to remove them, or felt that there was so much risk in the attempt, that it seemed safer to go on and remove the uterine tumour altogether. In those cases where the removal of the ovaries was successfully accomplished, an incision just large enough to admit two fingers was, with some patience, all that was necessary. When that failed the wound was enlarged, and even after introducing the hand, and sometimes turning out the tumour, the satisfactory removal of the ovaries could not be accomplished.

In case 28, on applying a ligature to one ovary which was close on the tumour, the silk thread went right through the tissues, cutting the ovary clean off, and leaving a hole in the side of the uterus, the hæmorrhage from which could only be arrested by the removal of the tumour. Indeed, I know of no operation that may be liable to so many accidents as the removal of an uterine fibroid, or the interfering with it in any way. In this case, the parts were strangely soft and friable, tearing on the least handling, so that, in fixing the cervix, I was afraid to screw up as tightly as usual, for the wires threatened to cut through. The consequence was that free bleeding soon came from the stump, and, when I was sent for five or six hours after

operation, there was a pile of blood-clot between the thighs; all the dressing had to be removed, and a few turns of the screw stopped the oozing.

We have thus to do with the results of fifty operations for the cure of fibrous tumour. The list contains every case of uterine fibroid that I have ever in any way interfered with by abdominal section. So far as I am aware, the results of these supravaginal hysterectomies are the best that have yet been obtained; and, as these are my first cases of this operation, it is only reasonable to suppose that the mortality will become lower by a longer experience. It seems to me that the reason of this comparative success lies in the fact that no operation was done unless there appeared to be some strong necessity for the doing of it. The tumours were generally large, the patients had suffered, and were more or less broken down, by pain and hæmorrhages. No one was operated on in good health or in good condition. No case of pediculated tumour was ever meddled with, simply because I do not consider that operations are necessary for such cases. Almost all were done on account of repeated hæmorrhages and ruined health. The time chosen for operation in the feeble ones—and most of them were feeble—was a day or two before menstruation was expected. The tumour might perhaps be then at its largest, but the patient had regained more or less force from the losses of the previous period.

The fatal case in this series is a typical one of the way in which, with one or two exceptions, my fatal cases after abdominal operations have died for some years past. Death seems to begin from the hour of the operation, or rather during it, before even the patients are placed in bed. There is a cold surface, a rapid feeble pulse—150 to 180, and this never comes down till they die. This patient was a soft, pallid, unhealthy woman from Shetland, never strong, and always ill fed. She had many scars of old strumous sores in her neck, hands, legs, and arms, whence pieces of dead bone were removed in her youth. When she first came to the Infirmary, she was so feeble, from long drenching hæmorrhages, that the menstrual periods were a succession of fainting attacks, and she often looked as if she would die. After some time, she was sent to the country, and was well fed and cared for. She improved, but the pulse remained feeble, sometimes scarcely perceptible. She was 43 years of age, and at first was advised, if possible, to fight on without interference till the menopause. She was unwilling to return home after having come so far. She was long about the Infirmary, and had seen many come, and go away well. So it was agreed to remove the ovaries. There was no encouragement to remove the tumour, which came down into the pelvis without any cervix. It was scarcely movable, and both broad ligaments were opened up. As she went under the ether, she begged me to take away everything, if at all possible, for she would rather not live as she had been for so long. Unfortunately, neither ovary could be found. The wound was enlarged, to admit the hand; the left was then felt very low down, and adherent. It could not be got at so as to apply a ligature round it. The tumour itself was enveloped by the broad ligaments, and was entirely subperitoneal. The bladder covered its anterior surface, and it could only be partly pushed into the wound by strong pressure from the vagina. After much trouble, the tumour was enucleated out, and wires placed as low as possible. The base could not be properly fixed outside, and, while trying to arrange this, a very large uterine artery below the wire was injured, and could only be secured by retaining on it a pair of locking forceps. She was cold and pulseless when put to bed. By evening the pulse was 160, but scarcely to be felt. This continued all next day, the temperature not rising above 99°. With stimulation, she lived into the next day.

I have no one way in dealing with the attachments of uterine tumour. At present, each case must be a law unto itself, and of this part of the operation there is much to be learned. A few of the simpler cases may be treated entirely extraperitoneally. Generally, the broad ligaments must be left inside; and sometimes the whole attachment, when there is much enucleation, must be so treated. Sometimes the treatment may be entirely intraperitoneal by means of Kœberlé's *serre-neud*, or it may be half intra- and half extra-peritoneal. These cases require much care in the after-dressing, though the convalescence is much shorter than when the whole is left outside. I am hopeful that the cautery will yet be the best and safest of all the methods of dealing with some of these tumours. The more I use it in ovariectomy, the more I like it. It is simply perfect, and its employment seems to me to be "a higher exercise of our art" than the ligature, which, apart from the chances of hæmorrhage, embraces ten times the amount of tissue that is really necessary. That a more perfect way will soon be found I have little doubt. This will do as much for uterine tumours as Baker Brown's intraperitoneal method has done for ovariectomy ever since 1864.

In two of these operations, and both were amongst the early cases

done many years ago, there was a mistake of diagnosis. The tumours were supposed to be ovarian, and were not recognised to be uterine till the abdomen was opened. Both patients recovered. In all the rest, a careful and correct diagnosis was made.

I often ask myself the question, Does a mortality of 8 per cent. justify an operation for a disease that, as a rule, has only a limited active life, that torments simply, and that only for a time, though of itself it rarely kills? The mortality of an ordinary uterine fibroid, if left alone, is nothing approaching a death-rate of 8 per cent. Most of the cases on which I have operated were known to me for years before; only the extreme cases were done; in nearly all, the lives were useless, and the risk of operation was clearly understood. Considering the nature of the cases, it seems to me that these operations were, perhaps, justifiable; and, if these were barely justifiable, what can be said of those ghastly lists of hysterectomy where the mortality is one death in every two, one death in every three, or even one death in four or five. Dr. Bigelow, of Washington, has lately collected all the cases placed on record up to March 1884. At best, this must be an imperfect list, and can only show the least bad side of the operation. Of 359 operations done by 16 of the most successful operators, there were only 227 recoveries and 132 deaths, or a greater mortality than one of every three operated on.

As in ovariectomy, I have long ago given up the use of carbolic spray during these operations. In only 6 of these 38 was it employed; one of these died. There was blood in her urine the next day; the day after, albumen, and then came an attack of acute mania, from which she died. Of 32 done without spray, there were two fatal cases, and both died from exhaustion. In truth, there is nothing in all my work that has so thoroughly broken down with me, as the carrying out of the so-called "perfect Listerism" in the surgery of the abdomen, by means of the carbolic spray. I expected much, but have got nothing, after years of vexation and disappointment; and I am now very much where I was before I ever heard of it. My results in ovariectomy were 1 in 26 just before I began the spray. They are almost the same now, since it has been given up. It is true that there was a long series of success, and, at the time, I thought that this was due to the use of the spray, and not to the steady use of the cautery, as I now think that it was. There were 80 consecutive cases of recovery—that means 2 deaths in 82, or 1 in 41. But of the whole number of 120 spray-cases, there were 7 deaths, or 1 in 17. In the Infirmary, the results were disastrous, 1 out of every 7 died. Fortunately, there were only 21 operations in all done under the spray; and my impression is—I may be wrong—that these would have lived had the spray not been used. Since the spray was abandoned, there have been done in the infirmary 88 ovariectomies up to this date, December 8th, 1884. These include cases done by the assistant as well as by myself. The number of deaths is 2, or 1 in every 44 operated on; neither of these died of septicæmia. In one—a very difficult case of large, very solid, tumour, opening up the broad ligament—the patient died a week afterwards of uræmic poisoning, the kidneys having secreted nothing for two days. The right ureter was found to be included in one of the many ligatures. This is the only disaster that has happened to me during the operation in fully 550 times that I have opened the abdomen for the removal of uterine or ovarian tumours. The other fatal case was that of a feeble woman coming out of a severe operation, with a pulse at 170; and it never fell.

In the BRITISH MEDICAL JOURNAL of the 28th October last, Mr. Thornton made the general statement that his results in ovariectomy establish the valuable protection afforded by the spray; and that, practically, septicæmia is removed from the causes of mortality in abdominal surgery; and he quoted certain experiments made by Professor Chiene on the atmosphere of his own wards in the Infirmary here, as also supporting the principle of the protection afforded by the spray. Mr. Thornton's operations were performed in the Samaritan Hospital, which is practically a private house, with every comfort. He had done 100 cases of ovariectomy with 3 deaths; that is, he had done 101 cases with 4 deaths. The results I have already given are even more favourable without the spray. Not that I think one or two per cent., on one side or another, is of much consequence in proving anything; for, when the percentage is so low, we get within the range of accidental circumstances. But, if septicæmia be banished in all abdominal surgery, what are the results, and what the cause of death, in the numerous operations for the removal of uterine fibroids that Mr. Thornton has performed? On this subject he gives no information; but Dr. Bigelow's tables tell what the mortality was. A death-rate of one in every three of fibroids of all kinds spoiled his protection-spray theory, and generally did not suit his purpose, though the readers of the JOURNAL were, all the same, left under the impression that the spray, etc., had banished septicæmia from all abdominal

operations. It is said somewhere, as an excuse for bad results, that the antiseptic principle cannot be properly carried out in hysterectomy. Then what is the use of it? I have also read somewhere, as another excuse for fatal results, that, in cases of ovariectomy where the cyst has been previously tapped, the spray does not answer; and that the tapping—done, perhaps, long before—is to be blamed for the death. It is difficult to see why; for, in my hospital-cases, nearly one-half were previously tapped, once or oftener, before operation. Hysterectomy is a much more dangerous operation than ovariectomy; and an ovarian tumour, often tapped, generally presents greater local difficulties, and its removal is accompanied by a greater risk. Be this as it may, the fact is that we must look for greater success in hysterectomy in the developing of the technical methods of operating, just as Mr. Baker Brown did twenty years ago, when, by a simple change in his way of treating the pedicle of an ovarian tumour, he at once lowered the mortality of ovariectomy by two-thirds; only in this case the London surgeons would have none of it, but worked away on the old lines, losing one patient out of every three or four, while this great improvement really seemed to be wilfully neglected; and now, when the real discoverer of a perfect intraperitoneal method is long lying in his grave, they begin to wonder what Mr. Baker Brown really had to do with the advancement of abdominal surgery.

Those who teach that the carbolic spray in abdominal surgery is anything else than an useless ceremony can make of these results what they may. One thing is, however, certain; both ways cannot be the right way. If a "single germ," getting into the abdomen during an operation, play the mischief it is said to do, then few cases done in the old way ought to recover at all; whereas, in the heart of the very surgical hospital, and almost next door to where Professor Chiene tells us that the atmosphere is laden with death-carrying germs, less than one only out of every forty have died after ovariectomy. What is one to make of all these things? If Mr. Thornton gets a fatal result in every third case where he removes a uterine fibroid, with his complete and perfect Listerism, spray, and all the rest of it, am I to go back to these ways, when I get one out of sixteen without them? By no means. The antiseptic principle, which I believe in as much as anyone, can be carried out by simpler means than these; and, for myself, I have almost gone back to the boiled water and soda of twenty years ago. It is, unfortunately, a sad fact that, ever since surgery began, the great evil was done by the surgeon himself. It was the willing and tender, though unclean hand, that carried most of the poison into the wounds. It is to this that Lister has put a stop. With a proper antiseptic, a surgeon is now made to be clean in spite of himself, is compelled to have safe sponges, safe ligatures, clean instruments, and, above all, clean fingers. If one be careful enough—and few are careful enough—one may do all this, as Mr. Lawson Tait does, with boiled water and soda. Some such precautions are essential; beyond these, with ordinary care, we need not disturb ourselves much as to what is in the air. Yet it was a pleasant doctrine to believe in, to put the whole blame of a bad result that should not have been a bad result, upon some indefinite unknown something in the air—something beyond ourselves. It was no fault of ours; it could not have been helped. Everything was done that could have been done. I fear we are all apt to blame place, persons, things, accidents, circumstances—anything you like under the sun—rather than ourselves.

CLINICAL REMARKS ON A CASE OF PAROVARIAN DROPSY.

By THOMAS OLIVER, M.D., M.R.C.P. Lond.,

Physician to the Infirmary, Newcastle-upon-Tyne, and Lecturer on Practical Physiology in the University of Durham College of Medicine.

THE possibility that a cyst occupying the position of an ovarian cystoma may be other than ovarian, is not, I think, sufficiently borne in mind by most of us. A short time ago, I was present at an operation for the removal of a cyst which was considered ovarian. The abdomen was enormously distended, the distension giving to the abdomen a peculiar square appearance. The patient, who had reached the period of middle life, had noticed the tumour seventeen years previously, after the birth of her last child. The abdominal wall was so very thin that the operator, with one or two scratches of the scalpel, went immediately through skin, an extremely attenuated layer of muscular fibre, peritoneum, and what afterwards appeared to be adherent cyst-wall. Forthwith there escaped perfectly clear thin fluid, clear as water. At first it was thought to be ascitic, but it was, in reality, the contents of the cyst; and, in all, fifty pints of fluid escaped. It