I heartily endorse. The worst feature of the present system is, that it produces men, so imperfectly educated in a vast number of subjects, and at so high a pressure, that the majority seem incapable of pursuing a steady system of self-culture, after they have obtained their degrees or diplomas; and, without this, hope there is none of their becoming efficient practitioners. Without self-culture, and the thoughtful investigation of the clinical facts which come under the observation of the young physician, advance in the true knowledge of his profession is hopeless; and medical reform, be it ever so carefully planned, will, when obtained, be but a barren victory.

ON THE SOURCES AND THE EXCRETION OF CARBONIC ACID IN THE ECONOMY.

Read at the Section of Anatomy and Physiology at the Annual Meeting of the British Medical Association, in Belfast.

By J. J. CHARLES, M.A., M.D., D.Sc., F.R.U.I.,
Professor of Anatomy and Physiology, Queen's College, Cork.

Not many years ago, it was a general belief among physiologists that the generation of carbonic acid occurred chiefly in the lungs, owing to the direct oxidation of part of the carbon contained in the blood; and, accordingly, the lungs were regarded as the chief, if not the sole, centre for the production of the animal heat. Respiration, in fact, was described as a combustion-process occurring in the lungs, a process with which the rest of the body was not in any way directly concerned, except, perhaps, in supplying to the blood some of its effete carbonaceous material to be burnt up in the course of respiratory oxidation. Chemical and physical investigations, however, have caused this view to be discarded. It has been found, for example, that the venous blood going to the lungs contains a large quantity of carbonic acid, and is absolutely warmer than the blood leaving the lungs, as has been proved in the case of the blood in the right and left sides of the heart respectively; the blood, moreover, which leaves the liver and other glands when in a state of activity, has been shown not only to possess a higher temperature, but also to be richer in carbonic acid than the blood entering these organs.

In short, the temperature of the blood, according to Claude Bernard's and Korner's observations, is highest in the liver, that of the blood in the hepatic vein being, on an average, 0.7° Cent. to 0.9° Cent. higher than that in the portal vein; and the next highest temperatures have been noted in the blood flowing respectively from the brain, the salivary glands, and the muscles, particularly when these parts are in an active condition. It is true that Heidenhain (*Physiologie*, von Dr. L. Hermann, Band v, theil 1, s. 274) has arrived at results differing somewhat from those of Bernard, for he finds that the blood of the liver has a temperature not exceeding that of the blood of the spleen, or of other abdominal viscera; yet there is no doubt about the temperature of the blood from the liver being high as compared with that of the blood from the lungs, muscles, etc.

What, no doubt, has greatly tended to direct so much attention to the importance of the tissues in respiration, has been the result of the investigations of the properties of protoplasm in the simplest forms of animal life. The primitive amœba has been shown to absorb oxygen, and give out carbonic acid; and the natural inference is, that all tissues containing protoplasm act after a somewhat similar fashion. The closest attention has, therefore, been directed of late to the minute chemical changes occurring at the tissues themselves; and, accordingly, we find Liebig, Matteucci, Valentin, and Hermann pursuing their investigations directly upon living muscles that had been removed from the animal body, and enclosed in tubes inverted over mercury; Ludwig and Schmidt analysing the blood before and after it had passed through the living and active muscles of the animal so as to detect its alterations; and Hammarsten investigating the gases of the lymph with the same object in view.

In the case of secreting glands, likewise, observations have been made upon their recent secretions, as well as upon the changes effected in the blood during its passage through them. The saliva, bile, urine, and the like normal fluids result directly from the action of the histological elements of the secreting tissues, and are, therefore, more immediately influenced thereby than can be the lymph or blood flowing through the same tissues; but by investigations on the secretions, as well as on the lymph and blood of the tissues, we obtain not only the quantity of the gases in the tissues, but likewise their tensions.

The importance, indeed, of these investigations into the gases of the secretions is very great; for, by their means, much light is thrown

upon the seats of the oxidation processes in the economy. By a very simple experiment an example of this oxidation, with the concurrent disappearance of oxygen, and the possible formation of carbonic acid, may be satisfactorily demonstrated. If we examine with a spectro-scope, in a dark room, the red coloured light that passes between two of the fingers held together in front of a good lamp, the spectrum of oxyhæmoglobin will show itself. If, however, we apply a ligature to the fingers close to the palm, in order to prevent the entrance of fresh arterial blood, and examine as before, instead of the two bands of oxyhæmoglobin, there will be seen, in the course of a few minutes, only the single band of reduced hæmoglobin.

I. SOURCES OF CARBONIC ACID.—With these preliminary remarks, we may now discuss the possible sources of carbonic acid in the organism. Some of the better known of these we shall merely glance at in passing; others, however, which are less familiar, as well as of a more hypothetical character, we shall consider at greater length.

1. The chief source of carbonic acid in the organism, beyond all question, is *simple oxidation*, in which a more or less direct combination occurs between the absorbed oxygen and the carbon of the food and tissues. The fatty matter of the food which is not stored up or directly consumed by the tissues requiring it, is ultimately transformed into carbonic acid and water, just as the like destination awaits the stored up fat of adipose tissue when it comes to be oxidised in the system. The carbo-hydrates (starches and sugars) taken as food, give origin likewise, when oxidised, to carbonic acid and water, although it is difficult to say whether these terminal products are direct or indirect. Indeed, there may be intermediate products, as in the case of the sugars-aceton sometimes appearing in the urine in cases of diabetes and continued fever; or glycosates may first be formed in the blood, and afterwards undergo complete oxidation. That some of these carbo-hydrates are oxidised during muscular activity, thus effecting a large production of carbonic acid, is also a very reasonable hypothesis. The proteids, further, may give rise to carbonic acid as the result of oxidation; but it is more probable that a direct complete metamorphosis of this kind takes place only to a very slight extent. As to the two views held regarding the part played by the albumen of the food; whether a portion of it enters at once into the constitution of the tissues while the rest circulates for a time in the blood, forming there unstable and readily destructible combinations; or whether the whole of it becomes tissue-albumen before undergoing subsequent metabolism; it would seem, without entering on a critical discussion of their respective merits, that the latter is the more likely; for the great increase in the amount of urea excreted after food, a fact which appears to be in favour of the former hypothesis, may be readily explained on the very probable assumption that the leucin and tyrosin, formed at the expense of the albumen in the alimentary canal, are absorbed, and thus form the immediate source, together, possibly, with kreatin, of the bulk of the urea then discharged.

Mention may also be made of Bidder's and Schmidt's opinion, that the albumen of organs is regularly destroyed at the rate of about one per cent. of the total albumen of the body in the twenty-four hours; and that, in cases of death from starvation, when the glands and muscles have lost 50 per cent. of their albumen, some of it has been oxidised at the tissues, and some of it liquefied and transferred to other parts of the body, as the brain, etc., for their nourishment.

Reference may likewise, with advantage, be here made to the respiratory exchanges which occur when certain tissues, such as muscle, brain, and bone, are placed in an atmosphere of oxygen. We find, then, that 100 grammes of muscle absorb 50 c.c. of oxygen, and exhale 56 c.c. of carbonic acid; 100 grammes of brain absorb 45 c.c. of oxygen, and exhale 42 c.c. of carbonic acid; and 100 grammes of bone absorb 17 c.c. of oxygen, and exhale 8 c.c. of carbonic acid. The consumption of oxygen and the evolution of carbonic acid, it will thus be seen, are highest in the case of muscle.

As to the influences regulating this destruction of the organic materials of the body, Hoppe-Seyler states that they appear to be intimately connected with the activity of the histological elements of the organs; but whether they are of the nature of fermentation or otherwise, it would, for the present, be impossible to say.

2. The second source of carbonic acid in the system is the *decomposition of albumen by hydration or dehydration*. In a paper (*Journal of Anatomy and Physiology*, 1882, p. 298) "On the Gases of the Bile," which I published in 1882, I put forward an hypothesis as to the probable mode in which albumen might, by its decomposition at the liver, produce carbonic acid. I there assumed that a molecule of albumen might be supposed to combine with 50 molecules of water, and yield 8 molecules of urea, 7 of glycogen, 5 of carbonic acid, 7 of oxygen, and 1 of sulphuric acid. The oxygen would not, of course,

remain free, but might enter into combination with reduced hæmoglobin or other substances. In favour of this assumption, it should be remembered that, of the 18 atoms of nitrogen in a molecule of albumen ($C_{12}H_{112}N_{18}O_{52}S$), 4 probably belong to the urea group; and also that carbonic acid has actually been obtained in the laboratory from albumen.

Schützenberger had, a few years previously, advanced an hypothesis somewhat similar to mine, although I was not aware of it at the time of the publication of my paper. He assumes that 100 grammes of albumen generate first 35.5 grammes of urea, and the residue, by subsequently combining with 12.3 grammes of water, gives origin to 27.4 grammes of carbonic acid, and 51.39 grammes of fat. He has further shown that albumen may be decomposed into carbonic acid, ammonia, oxalic and acetic acids, and that the ratio of the carbonic acid and ammonia to each other is the same as if urea had been made the basis of experiment. He also believes that the carbonic acid and ammonia may afterwards combine in the body under the influence of electrical currents, and thus form urea, dehydration having taken place at the same time. If this view prove correct, some of the carbonic acid evolved in the liver would not leave that organ as such, either by the blood or the bile, and would not, therefore, appear at the inner or the outer surface of the body in the gaseous form.

We can readily believe in the albumen of the tissues, particularly the muscular and the nervous, undergoing these chemical changes during its metabolism, and so account for the carbonic acid present in all the tissues, and for the urea found in some of them; but it is, moreover, probable that a part, at least, of the primary decomposition products of the albumen may be yielded up by the tissues to the blood, and thus conveyed directly to the liver, where the decompositions above described may ensue. That chemical changes, such as oxidations and decompositions, occur to a large extent at the liver, is indicated by the high temperature of the blood of the hepatic vein, as well as by the great quantity of carbonic acid in the bile. In fact, there can be no doubt that the destruction of albuminous material is most active at the liver, for this organ, as we know, contains about a fourth of the total blood of the body;—an amount equal, indeed, to that in the muscular system, although this system constitutes at least half the normal body weight; besides, there are to be found in the liver a great number of nitrogenous substances intermediate between albumen and urea, such as leucin, tyrosin, xanthin, hypoxanthin, uric acid, etc. According to most authorities, also, there is more urea to be met with in this organ than in any other in the body, many physiologists even believing that the bulk of the urea is formed here. The urea certainly cannot be derived from the blood by a mere process of filtration into the liver; for the nervous and muscular tissues, which do not directly produce urea, contain little or none of it.

Not only, however, may albumen be decomposed at the liver and produce carbonic acid, but there are good grounds for believing that the crystalline indifferent proteid—hæmoglobin, which is a higher albumen derivative, is broken up in the liver into a number of simpler bodies—an hypothesis advanced by Dr. Zuelzer of Berlin. (*Untersuchungen über die Semeiologie des Harns*, Berlin, 1884.) The following equation expresses the decomposition—2 hæmoglobin ($= C_{1200}H_{1920}N_{308}Fe_2S_6O_{563}$) + 85 H_2O + 123 $O = Fe_2O_3 =$

	C	H	N	O	S
6 taurocholic acid ..	156	270	6	42	6
24 glycocholic acid ..	624	1032	24	144	—
2 bilirubin	64	72	8	12	—
135 urea	135	540	270	135	—
4 cholesterin	108	176	—	4	—
113 carbonic acid	113	—	—	226	—
	1200	2090	308	563	6

[The fact that the use of albuminous food increases the amount of biliary acids formed at the liver, tells somewhat in favour of this hypothesis.]

Or, 13.332 grammes of hæmoglobin, with the necessary water and oxygen, yield 2.486 grammes of carbonic acid. A further decomposition of a smaller proportion of hæmoglobin with the formation of glycogen (taking glycogen as $C_{30}H_{50}O_{25}$; Schtscherbakoff) is also probable. Thus 2 hæmoglobin ($= C_{1200}H_{1920}N_{308}Fe_2S_6O_{563}$) + 501 O + 182 $H_2O = Fe_2O_3$ + 6 $H_2SO_4 =$

	C	H	N	O
2 bilirubin	64	72	8	12
150 urea	150	600	300	150
32 glycogen	960	1600	—	800
26 carbonic acid ..	26	—	—	52
	1200	2272	308	1014

Or 13.332 grammes of hæmoglobin yield 0.572 grammes of carbonic acid, and 12.96 grammes of glycogen.

Owing to the destruction of red blood-corpuscles which is constantly occurring in the spleen, a considerable quantity of hæmoglobin in solution is as constantly being carried to the liver; and if this be borne in mind, it will readily be perceived that the amount thus disposed of is far from inconsiderable. Moreover, red blood-corpuscles are also broken up at the liver; for glycogen, as Hoffman has pointed out, dissolves them. To give an idea of the rate at which old red corpuscles are destroyed, and hæmoglobin at the same time set free, I may mention that a calculation has been made which indicates that the duration of the life of a red corpuscle is at most about thirty days. Thus, taking the amount of iron in all the corpuscles of the body as one gramme, and the quantity excreted in twenty-four hours as 35 milligrammes, all the iron would be eliminated from the body in thirty days; for $\frac{1}{35} \times 24 = 30$ days.

Now whilst it must be admitted that some of the chemical changes I have described as taking place at the liver are to a great extent hypothetical; yet there can be little doubt that, as Heidenhain and others suppose, albumen in some form is decomposed in the cells of that organ, and biliary acids and pigments formed in the same cells as the glycogen. It is true that these two processes in the liver, resulting in the formation of glycogen and bile, though both are attended by a corresponding production of carbonic acid, differ markedly in that their periods of intensity do not synchronise; and, further, in that, whilst the biliary secretion continues during starvation, the formation of glycogen ceases. The secretion of bile, indeed, does not depend on the pressure of the blood; for the pressure under which it is secreted exceeds that of the blood in the liver, although the amount formed is influenced to a certain extent by its velocity and pressure, because with increased velocity and pressure of the blood, more albumen is brought in a given time to the liver to undergo metamorphosis.

II. SEATS OF EXCRETION OF CARBONIC ACID.—It has been computed that nine-tenths of the carbonic acid eliminated by the body passes off by the lungs, and the remaining tenth by the intestines, skin, and kidneys.

1. *The Lungs*.—About 900 grammes of carbonic acid, on an average, leave the body in the twenty-four hours by the lungs; that is, an amount of carbonic acid equivalent to 249 grammes (or 8 ounces) of solid carbon. This discharge of carbonic acid is chiefly due to diffusion, the tension of the carbonic acid in the venous blood being greater than that of the carbonic acid in the air of the air-cells. Of every five parts exhaled, one part is possibly derived from the blood-corpuscles, and the other four from the liquor sanguinis. In the corpuscles, there is probably some loose combination between the carbonic acid and the hæmoglobin; but in the liquor sanguinis, it is partly in solution and partly combined with sodic phosphate and carbonate.¹ Simple diffusion, however, is of itself scarcely sufficient to account for the total discharge, for we must also bear in mind that the gas is in a state of solution in its passage through the moist capillary walls and the epithelial lining of the air-cells. Further, the entrance of oxygen undoubtedly increases the tension of the gas in the blood, a fact which has been pointed out by Holmgren.

2. *The Skin*.—An average of nearly four grammes of carbonic acid is discharged by the skin in the twenty-four hours—that is, about 1-200th of the amount excreted by the lungs. The gas is exhaled from the sudoriferous glands, and partly also from the surface of the skin itself.

3. *The Secretions*.—I have already referred to the light which a full knowledge of the gaseous conditions of the secretions throws upon the respiratory interchanges at the tissues. Indeed, more is to be learnt from the study of the gases of the secretions even than from that of the gaseous exchanges, at the tissues, of either the blood or the lymph; for the secretions have been more directly in contact with the elements of the gland-tissues—being as it were part of the tissues themselves, and hence the importance of investigations in this direction.

The tension of carbonic acid in the cavities and liquids of the body, when these are surrounded on all sides by healthy tissues (as in the case of the bile in the gall-bladder and the urine in the urinary bladder) is greater than that of the carbonic acid in venous blood.² The tension of the gases in the tissues is determined indirectly from the tension of the gases in the lymph or in the secretions; but the

¹ Carbonic acid may likewise be united to paraglobulin in the red corpuscles. Setchenow believes that the colourless corpuscles also fix carbonic acid, their absorbing power for this gas being $\frac{1}{10}$ th that of serum.

² Tension of carbonic acid in arterial blood, 21.28 mm. of mercury; venous blood 41.0; bile, 50.0; acid urine, 68.0; hydrocele fluid, 46.5; fluid of peritoneal cavity, 58.5; lymph, 35.5.

latter mode is to be preferred, for in lymph the carbonic acid tension is less than in venous blood, partly owing to gaseous diffusion between the arterial blood and the lymph, where the vessels containing them are in close proximity—an objection, however, which does not apply in the case of the secretions.

The secretions, as a rule, are formed in the interior of cells, either directly from their own protoplasm, or by the agency of the latter from materials derived from the blood. The phenomena of secretion are accompanied by the disengagement of heat, readily demonstrable by means of the thermo-electric needle, both in the blood and in the secretion formed.

Of these secretions, we shall first consider—

a. The Bile.—In this secretion, collected directly as it flowed from the liver of the dog, I found, in a series of experiments I made about two years ago in Bonn, a total of 57 volumes per cent. of carbonic acid,³ 14 per cent. of which was evolved *in vacuo*, while the remaining 41.7 per cent. required the previous addition of phosphoric acid to disengage it; whereas oxygen and nitrogen there were only traces. That this proportion of carbonic acid is very large can readily be understood, when it is stated that there are only 34 volumes per cent. of the gas present in the arterial blood of the dog, and 46 volumes per cent. in the venous blood of the same animal. The tension of these gases in the bile of the dog, as taken from the gall-bladder (for I am not aware that the tension has been determined in the case of freshly secreted bile), is comparatively low—only equal to 50 mm. of mercury; but then it should be recollected that the tension of carbonic acid in venous blood is only equal to 41 mm.; the blood-pressure also in the portal vein is very low, perhaps not more than 10 mm. From the bile of the rabbit (the only herbivorous animal in which freshly secreted bile has up to the present been examined) I obtained a total of 109 volumes per cent. of carbonic acid, free and combined—a larger proportion than has yet been discovered in the fluids of any animal, but only slight traces of oxygen and nitrogen.

With regard to the possible source of the large amount of carbonic acid excreted at the liver, two views suggest themselves: (*a*) that the carbonic acid is produced in the different tissues and organs of the body, the liver included, and some of it then passes out by simple diffusion or solution from the portal and hepatic capillaries into the bile (as may occur at the pleura in lymph exudations), the hepatic cells not being specially concerned in the process; (*b*) that, in addition to this general formation of carbonic acid in the body, there is a special production of the gas at the liver, owing to the decomposition in the hepatic cells of such bodies as albumen and hæmoglobin. I have already given reasons in support of the latter hypothesis. Of the carbonic acid thus generated in the liver, part will enter the blood and a larger proportion the bile, particularly if this fluid is alkaline; just as the carbonic acid formed in the tissues passes off from them in the lymph and in the venous blood, but a greater proportion in the latter, because it exercises a stronger chemical attraction for carbonic acid.

The amount of combined carbonic acid in animal fluids or secretions depends, as a rule, on their reaction, whether alkaline, acid, or neutral. If the secretion be alkaline, more of the tissue carbonic acid will leave the body by that fluid than by one less alkaline; while, if neutral, or particularly if acid, scarcely any combined carbonic acid will be found in it, as compared with the total carbonic acid in the blood. A good example of this can be observed in the case of bile. If the bile of a dog is alkaline, as it is usually, 100 volumes may contain 57 volumes of carbonic acid; while, if acid, there may be in 100 volumes only 5 volumes of the same gas. Urine, again, which is acid, contains only 14 volumes per cent. of carbonic acid; while, in alkaline saliva, there may be present as much as 50 volumes per cent. But, in herbivorous animals, in which the bile is more alkaline than in carnivorous, there is a proportionately larger amount of carbonic acid, and this greater richness in the gas may be the case, as well, with the other fluids of these animals, though this has not yet been determined.

Accordingly, from the proportion of carbonic acid in the bile, it would be unsafe to estimate the total amount of carbonic acid generated in the liver; for, if the bile formed be less alkaline than the blood flowing away from the liver, the greater part of the carbonic acid will probably be conveyed away by its means to be excreted elsewhere. I am not aware that any experiments have been made as to the relative quantities of the carbonic acid in the blood of the portal and hepatic veins, an exact knowledge of which would be most useful in our present inquiry; and without it, I do not feel myself in a position to say more on this subject.

³ In this experiment, as in almost all referred to in this paper, the volume of the gas is given at a pressure of one metre, and a temperature of 0° C.

b. The Urine.—The presence of sodic phosphate and carbonate in the urine renders it capable of dissolving carbonic acid. Pflüger found that 100 vols. of urine yielded of free carbonic acid 17 volumes and of fixed carbonic acid set free by phosphoric acid, 0.2 volume. Some of this carbonic acid is probably formed by the cells of the kidney, because nitrogenous metabolism occurs here; but the larger proportion is possibly excreted in solution in the fluid discharged at the Malpighian capsules. An interchange of gases likewise goes on in the tubes of the kidney. The tension of the carbonic acid in the urine is higher than that in the bile, being equal to 68 mm. of mercury.

c. The Saliva.—The submaxillary saliva of a dog fed on flesh-meat yielded Pflüger 19 volumes per cent. of free carbonic acid, and 30 volumes per cent. of carbonic acid evolved on the addition of phosphoric acid—that is, about 50 volumes per cent. of the gas, free and combined; and when the animal was fed on a mixed diet, the total volume of carbonic acid was considerably greater. In its richness in carbonic acid we, therefore, notice that this secretion approaches bile closely, but if the small volume of the submaxillary saliva be taken into account, it will readily be perceived that, as compared with bile, only a small proportion of the carbonic acid is thus discharged.

d. The Milk.—This fluid, according to Pflüger, contains 7 per cent. of free and combined carbonic acid.

e. The Pancreatic Juice.—The amount of carbonic acid in this secretion has not yet been ascertained.

f. The Intestinal Secretions.—It is impossible to say what proportion of the gases in the alimentary canal is derived from the blood by diffusion, what from the air swallowed, and what from the fermentation and decomposition of the intestinal contents. As Planer's researches have shown, the nature of the diet affects the composition of the gases in the intestines. Carbonic acid is always present in large amount. A gaseous interchange between the intestinal contents and the blood very probably occurs, for in many animals (the air-swallowing fish, *cobitis fossilis*, for example) intestinal respiration appears to be important.

g. The Normal and Abnormal Exudations.—Reference may be made, in conclusion, to the carbonic acid in these liquids. They resemble dilute liquor sanguinis in composition, and may or may not be spontaneously coagulable; but they are always alkaline, and, therefore, capable of retaining much carbonic acid in solution. The gas is derived from the tissues by means of the lymph poured out into the serous or other cavity, or it leaves the blood in a state of solution, and accumulates according to the solvent or fixing-power of the medium; and its amount may be increased by decomposition or oxidation occurring in the exudation. Hydrocele fluid has yielded 48 volumes per cent. of carbonic acid, 24 per cent. being free, and 24 fixed; pleuritic effusion 44 volumes per cent. of carbonic acid, 15 per cent. being free, and 29 fixed; peritoneal fluid 10 volumes per cent., 7 per cent. being free, and 3 fixed; cedematous fluid of the extremities 24 volumes per cent., 17 per cent. being free, and 7 fixed. In many of the liquids to which I have referred above, the tension of the gas is greater than it is in the blood. Only traces of oxygen and nitrogen, it may be added, are to be met with in any of these exudations.

Dr. McVAIL (Glasgow) congratulated Professor Charles on the very careful and important experimental research, the results of which he had brought under the notice of the meeting. If the processes of estimating the gases in the secretions and excretions could be so simplified and systematised as to be possible of application in ordinary clinical work, an important service would be rendered to medicine. Up to the present, while physicians had made great advances in practical acquaintance with the greater mechanical actions of the body, such as respiration and the circulation, they could do little in the way of estimating the metabolic changes going on in the cells of the organs and tissues, although abnormal conditions there might be said to constitute the very fountain-head of disease. More particularly were Dr. Charles's remarks on the gases of the bile of great importance, and investigations pursued in the same direction could not fail to extend greatly the knowledge of the metabolic processes going on in the liver. He trusted Dr. Charles would continue the important work on which he had so successfully entered.

Dr. ANDERSON (Galway) said that it was of the highest importance that the gases as well as the solid constituents of a tissue or fluid should be examined. The fresh bile from the receptacle was undoubtedly a fluid good for analysis, and it must always be recollected that such a fluid had been taken from the gall-bladder. The bile might, and indeed must, undergo changes in its course from the capillaries to the gall-bladder. In passing along the intestinal tube, it was altered; whilst it remained in the gall-bladder, it became changed. The greater part of the oxygen of the blood was undoubt-

edly consumed in the tissues; but the blood in a tissue was itself undergoing change at all parts of its course. Oxygen must be consumed, and carbonic acid produced in the lungs, as well as in the liver or skin. Urine in the capillaries of the kidney was not the urine of bladder; the saliva of the acini was not the saliva of the duct; and the bile of the gall-bladder was not the bile of the hepatic capillary ducts. Chemistry alone could teach the accurate constitution of tissues, and to that science we must look for progress.

Dr. CHARLES said, in reference to Dr. McVail's remarks, that he was afraid the method of estimating the gases in liquids pursued at present was of too complicated a nature to be made use of in clinical investigations. As to the statement that the results of the analyses of the gases of secretions were not reliable, owing to decomposition occurring in the secretions after death, he held that this objection would not apply to his experiments, as they had all been made on bile flowing directly from the liver. Indeed, results of this kind were, in his opinion, as trustworthy as those derived from an investigation of the gases of the blood.

THE CURE OF WRITER'S CRAMP.

By A. DE WATTEVILLE, M.A., M.D., B.Sc.,

Physician in Charge of the Electro-Therapeutical Department, St. Mary's Hospital.

THE group of symptoms—neuralgic, paralytic, and spasmodic—the varying combinations of which, in individual cases, constitute the ailment known as writer's cramp or scrivener's palsy, has hitherto defied the most strenuous efforts of therapeutics. At least, the instances in which improvement has taken place are so few and far between, as to illustrate the saying that "exceptions confirm the rule."

I need not, in this paper, give a full account of the different theories propounded at various times in explanation of this singular neurosis. The fact of its clinical versatility, sometimes with a sensory, oftener with a motor predominance of symptoms (tonic, clonic, or parietic), certainly points to a strictly peripheral origin. Several observers have adopted this view, and Dr. Vivian Poore, who has devoted much attention to the etiology of writer's cramp, has advocated it with much ability, describing this ailment as an instance of what he calls "chronic muscular fatigue diseases." In this category, and intimately connected with the complaint now under consideration, naturally fall all those cases where groups of muscles brought into play in the performance of acquired co-ordinated actions, become the seat of nervous troubles. These "professional neuroses," as they have also been called by German writers, have received a considerable amount of attention during the last few years. Cases have been described among violinists, pianists, telegraphists, as well as among lawyers, tailors, machinists, and even ballet-dancers.

The pathognomonic symptom of the muscular disturbances in this class of cases is that the spasm or paralysis, as the case may be, becomes apparent only during the action which brings the particular co-ordination into play. With reference to the pain, when it does exist, it may be spontaneous; but if it be called forth or exacerbated by movements, it is only these co-ordinated actions which have the property of doing so.

Now with reference to the treatment of these "professional" neuroses, it may fairly be said that, not only has the *Pharmacopœia* been ransacked in the search of a suitable drug wherewith to combat the symptoms, but every known external means of treatment has in vain been tried. Successes in individual cases have occasionally been reported, but no method has proved regularly successful. Various shaped pens and supports for the hand and arm are but very poor substitutes for the healthy action of these parts in writing. Electrical and hydropathic applications have failed, even when combined with protracted rest and hygienic measures. Massage, if we are to believe sundry reports, has proved of service in certain cases; but, until five or six years ago, no one claimed to have discovered a method by which a large percentage of the sufferers from writer's cramp and other "professional" neuroses were speedily and permanently cured.

In 1881 Professor Charcot, on reliable reports published by some of the leading physicians and surgeons in Germany, called to Paris a gentleman, Mr. Julius Wolff by name, whose successful application of massage and gymnastics combined, had earned for him a great reputation in his native country. Two inveterate cases of writer's cramp

were placed under his care, with the result that, in both, the patients were able to resume their pens after two or three weeks' treatment (see *Progrès Médical*, January 21st, 1882). Since then the new method became known, by name at least, in this country; and we find it described, for instance, in the second edition of Dr. Ross's classical treatise on the *Diseases of the Nervous System*, vol. i, p. 606. (See also *The Year-Book of Treatment*, page 38.)

During the latter part of last year Mr. Wolff came over to this country for the purpose of demonstrating his method. The discouraging failures which had hitherto attended all my efforts for the relief of the comparatively numerous cases of writer's cramp and allied neuroses it had been my fate to meet, as contrasted with encouraging results obtained by the new system, made me anxious to obtain personal evidence as to its efficacy. I followed with keen interest the cases of two patients, treated under my eyes by Mr. Wolff, and of which I give a condensed account.

CASE I.—Mr. H. S., representative of a house of business, aged 38, was sent to me by Mr. Ernest Hart, with a view to deciding whether his case was a suitable one for the application of Wolff's method. The patient is tall and muscular, and though formerly rather delicate, has had no serious disease. There are some neurotic antecedents in his family. His present complaint began very gradually many years ago. He lost the power of writing quickly; his handwriting, which used to be good, became indifferent, from the uncertainty of his movements. Pains appeared in the forearm, upper arm, and shoulder. The thumb, index, and middle finger became the seat of cramps, accompanied with much pain. The spasm of the thumb made it slip off the pen. Then the hand began to turn. It took him fourteen minutes to write two lines in a very shaky manner. The patient had to give up his employment, and apparently improved by protracted rest; but not sufficiently to resume his writing. He was in this condition when I saw him; and as the case was one of typical writer's cramp, I immediately sent him to Mr. Wolff, under whose treatment he remained for four weeks. The operations of stretching, massage and local gymnastics, were repeated twice a day for from twenty-five to forty minutes each time. He has since called upon me to report himself cured.

Case I.

*I am suffering from
Writer's Cramp and
I am now cured
of the Cramp.*

CASE II.—Mr. F. B., merchant, a strong and healthy man, aged 35, without any antecedents worth noting, has, for over ten years, been unable to write properly. He began, without known cause, to experience a sense of fatigue in the hand and arm, but no pain. The thumb and little finger became more especially the seat of the feeling, which, however, did not preclude the possibility of writing. Actual weakness, however, set into these parts; later still, tremors and violent spasms made their appearance in the whole hand, in the thumb more especially. The power of guiding the hand was lost. On trying to write, the patient experienced such violent shaking, that his whole strength of will was unable to overcome it sufficiently to shape the letters. He could freely use his hand for every other purpose. He then gave up every attempt at writing. For the last three years he has been able to trace a few words. I saw this patient early in December, when he began Mr. Wolff's course of treatment. After five weeks, he was able to write for hours without pain, spasm, or tremor, and has now resumed an active share in all the duties of his profession. I may add that, as is usual in such cases, all the ordinary methods (electricity, hydrotherapy, etc.) had been tried previously, and failed.

I have appended facsimiles of the handwritings of these two patients before and after treatment. In addition to the tremulous, distorted character of the letters before treatment, the reader must bear in mind the time and efforts required, in both cases, to trace these few words, as well as the pain involved in the performance.