LECTURES

THE ANATOMY OF THE INTESTINAL CANAL AND PERITONEUM IN MAN.

Delivered at the Royal College of Surgeons of England.

By FREDERICK TREVES, F.R.C.S.,

Hunterian Professor at the Royal College of Surgeons; Surgeon to, and Lecturer on Anatomy, at the London Hospital.

LECTURE I.

MR. PRESIDENT AND GENTLEMEN,—The account of the intestinal canal and peritoneum in man, that I have the honour to bring before you, is derived from the systematic examination of one hundred fresh bodies. Through the kindness of my colleagues, Drs. Sutton and Turner, the pathologists to the London Hospital, I was enabled to open, before the performance of the usual necropsy, all the bodies of patients that had died of other than abdominal disease. The bodies, therefore, were quite fresh, and, in many instances, still warm. I had long been convinced that a study of this part of anatomy was rendered liable to many fallacies when conducted in dissecting-room subjects in whom decomposition had advanced, and in whom one could expect, for various reasons, some displacement of parts.

Certain questions as to vascular-supply I have followed out in injected preparations. I have also derived considerable information from the dissection of a number of fetuses of various ages.

Moreover, through the kindness and courtesy of Mr. Beddard, the Professor to the Zoological Society, I have been enabled to make a detailed examination of the viscera of many of the mammalia, the examination including the dissection of forty different species. This latter work has been of the greatest service in throwing light upon many obscure points in human anatomy.

The first question that may be considered is that which concerns the length of the intestinal canal in the human subject. I have made careful measurements in every instance, but the results, although voluminous enough, have been somewhat barren in interest. I find that the average length of the small intestine in the adult male (between the ages of 20 and 50) is 22 feet 6 inches, the extremes being 31 feet 10 inches in one case, and 15 feet 6 inches in another. The average length of the same part in the female is 23 feet 4 inches, the extremes being 29 feet 4 inches and 19 feet 10 inches respectively. The average length of the colon, in the same set of subjects, is 4 feet 8 inches in males, and 4 feet 6 inches in females, the measurement being taken from the root of the appendix or tip of the cæcum to the point where the meso-rectum ended. The extremes were, for both sexes, respectively 6 feet 6 inches and 3 feet 3 inches. I have convinced myself that the length of the bowel is independent, in the adult at least, of age, of height, and of weight; nor is the ratio between the measurements of the small and large intestine constant. A very long small intestine may be associated with a very short colon, or *vice versa*; or both segments may be unduly long or unduly short. Moreover, advancing age appears to have no influence upon the length of the intestine. The measurements of the bowel in subjects above the age of 50 years, are practically identical with those that have been just detailed.

I think, therefore, that it must be allowed that the differences in the length of the intestine—differences that, in the lesser bowel, may actually reach to no less than 15 feet—depend upon physiological, and not upon morphological, data. It is not unreasonable to assume that the nature of the food, the vigour of the digestive process, the activity of the abdominal nervous centres, will have more concern in determining the length of the bowel than will the height and age of the individual; and it may be that a time will come when physiologists will be able to express the value of certain alimentary functions in feet and inches. A study of the comparative length of the intestines in animals certainly seems to support a belief in the physiological bias.

The history of the growth of the bowel in infancy presents, however, some features of more definite interest. In the fetus at full term, the length of the intestine, and especially of the colon, is singularly constant. The average measurement of the small intestine

is 9 feet 5 inches; and of the large, 1 foot 10 inches. So regular are these measurements, that the greatest variation I have met with in the lesser bowel amounted only to 2 feet, while in the colon it was as little as 5 inches. During the first month after birth, it may be reckoned that the small intestine will grow about 2 feet; and a like rate of growth may usually be recorded at the end of the second month of extra-uterine life; but after that period, the development of the lesser bowel proceeds in a most irregular manner. Thus, in a child of one year, the small intestine measured 18 feet; while in another, aged two years, the length was only 13 feet 8 inches. Again, in one subject, aged six, the lesser bowel was no less than 21 feet in length; while in another child, eleven years of age, its length attained merely to 14 feet. It will be needless to observe that this growth of the small intestine has no concern with the general growth of the body; nor does it bear any relation to the weight of the child. Growth is intimately associated with activity of function; and the diet of young children among the poorer classes is often so unsuitable and erratic, that one must expect that the developing intestine is exposed to many fluctuations of fortune, and to many vascular vicissitudes. In one instance, the nature of the nourishment may lead to an apathetic, dull, and anæmic condition of the bowel; while in another case it may foster a perpetual intestinal tumult. It is impossible that such opposite conditions can have no effect upon growth; and, when it is remembered that all children start life with practically the same length of intestine, it is difficult to ascribe the extraordinary variations in growth that are conspicuous in the bowel during the early years of existence to other than physiological influences.

With regard to the growth of the colon, we may anticipate less conspicuous fluctuations. The functional activity of the large intestine is infinitely less than that of the small; indeed, the colon is little more than an internal receptacle, provided in order that faecal matters may be voided at intervals; and, in a certain physiological sense, the real anus is at the ileo-cæcal valve. The length of the colon in the fetus at full term is 1 foot 10 inches. When I came to examine my notes as to the length of the colon in infancy, I found that the measurement of this segment of the bowel at the age of one month was 1 foot 10 inches; at the age of two months, it was still the same; nay more, at three months, and even at four months, it was still the same—1 foot 10 inches. After this period, it grew steadily and regularly. In a subject a year old, it measured 2 feet 6 inches; at six years, it was about 3 feet; and at thirteen years of age, it was about 3½ feet in length. The absence of growth in the colon for at least the first four months of extra-uterine life is certainly remarkable. It is remarkable in this sense; in the fetus, the sigmoid flexure forms an enormous loop; it measures no less than 10 inches; while the rest of the large intestine measures a foot. Now, after birth, the sigmoid flexure rapidly ceases to be so exceedingly conspicuous; it soon begins to conform to the condition met with in the adult body; and, indeed, by the fourth month this segment of the bowel has relatively all the characters of the loop in the full grown individual. When measurements came to be examined, I found that the main part of the colon was growing at the expense of the sigmoid flexure; so that, in a subject that had reached the age of four months, this flexure measured only 6 inches, while the rest of the colon had acquired the length of 1 foot 4 inches. During this period of time, therefore, there had not been growth in length, but merely a readjustment of parts—a compensatory arrangement in subtraction and addition. It is peculiarly interesting to note that this rearrangement takes place without conspicuous disturbance of the investing serous membrane.

The Duodenum.—The length, course, and relations of the duodenum are singularly constant, and exceptions to the condition detailed in the familiar account of this portion of the bowel are exceedingly rare.

In the anatomical text-books, the duodenum is said to describe a single large curve of an almost circular shape, and is divided into three portions—an ascending, a descending, and a transverse. The first portion extends upwards, backwards, and to the right, to a point just beneath the neck of the gall-bladder. The next segment passes downwards as low as the second or third lumbar vertebra, while the third division crosses the column obliquely from right to left in front of the second lumbar vertebra. The end of the duodenum is thus clearly to the left of the spine. The first portion is invested by peritoneum, like the stomach; the second portion is covered by the serous membrane on its anterior surface only; while the third part receives a partial investment upon the same surface.

If the jejunum be considered to commence at a spot where the gut receives a complete investment of peritoneum and a distinct mesentery, then to the above description of the duodenum should be added

a fourth, or second ascending part. This fourth portion is practically constant. When the bowel has reached the left side of the column, it ascends vertically by the side of the spine. This vertical portion—which is covered entirely in front, and partly at the sides, by peritoneum—I have never found to be less than one inch in length. One anatomist (Dr. Bruce Young, *Journal of Anatomy and Physiology*, 1884, page 100) has recently described this terminal vertical portion, and insists that it is a constant feature in the normal duodenum. In one body, that of a male adult, I found this vertical portion so extensively developed, that the end of the duodenum was carried far up on to the under surface of the transverse meso-colon. The relatively small size of the duodenum, and its relation to the serous membrane, are points that are distinctive of the bowel in the higher mammalia. Speaking generally, it may be said that, in mammals, the duodenum forms a large loop, describes a regular and well-rounded curve, and is provided with an extensive meso-duodenum. In the primates, the meso-duodenum begins to disappear. In the ringtailed Lemur, there is a small but distinct meso-duodenum. In the common Marmoset, this fold is but slightly represented; while, in the fourteen different species of apes that I have examined, the peritoneum had practically the same relation to the gut that it has in man. The curve of the duodenum, however, varies greatly in monkeys. One extreme is represented by the simple angular horizontal bend met with in the Bonnet monkey and Barbary ape (Fig. 1 A), and another by the extensive vertical loop which I find that the duodenum describes in the Spider-monkey (Fig. 1 B). It is important to recognise that the end of the

duodenum is very firmly fixed in its place by the musculus suspensorius duodeni. This name has been given to a fibrous band, that contains, according to Treitz, some plain muscular fibre, and that descends to the vertical part of the duodenum from the left crus of the diaphragm, and the tissue about the coeliac axis.

The points about the duodenum to which I would direct especial attention, concern the fossa duodeno-jejunalis. This fossa has received little notice at the hands of anatomists, although it possesses great anatomical interest. It deserves attention, also, as being the seat of origin of retroperitoneal or mesenteric hernia, a form of rupture of which nearly fifty examples have been placed on record. If the transverse colon be thrown upwards, and the small intestine be drawn well to the right, the termination of the duodenum and commencement of the jejunum will be clearly displayed. The terminal part of the duodenum is covered in front by peritoneum. This covering is continuous above with the under layer of the transverse meso-colon, and below with the peritoneum covering the front of the lumbar spine, and passing down into the pelvis. On the left side, it is continuous with the serous membrane that invests the kidney and descending colon; while, on the right, it joins the mesentery of the small intestine.

When the fossa exists, a fold of serous membrane will be seen to pass from the parietal peritoneum, just to the left of the terminal part of the duodenum, and to be attached, in a vertical line, to the anterior surface of this portion of the bowel (Fig. 2). This fold, which is called the plica duodeno-jejunalis, forms above a free crescentic margin, which looks directly upwards; while, on all other sides, it is continuous with the peritoneum on the duodenum and posterior parietes. The fold is clear and translucent, is singularly free from fat and blood-vessels, is composed of two layers of the serous membrane, and forms a well defined pouch or pocket in the peritoneum.

The lateral attachment of the plica on the left side very commonly corresponds to the inferior mesenteric vein. It may come off between the vein and the duodenum, but never, so far as I have seen, from the peritoneum to the outer side of this vessel. To the bowel the plica is attached vertically, and nearly always along a line that separates the middle third from the left hand third of the gut. It may be attached nearer to what may appear to be the centre of the

bowel, but it is significant to note that, in any case, the line of attachment, if followed up on to the jejunum, will be found to be opposite to the line of attachment of the mesentery. The pocket, or fossa, formed by this fold, is of triangular outline, with its base above. The

opening of the fossa looks directly upwards. Its apex extends below the bend of the duodenum, a fact of significance in connection with the development of the fossa. The anterior wall of the pocket is formed entirely by the fold; the posterior wall is formed by the posterior parietes and part of the duodenum both covered by peritoneum. The capacity of the fossa varies greatly. In well marked specimens, it has a vertical depth of $1\frac{1}{2}$ inches; and will lodge the thumb up to the first joint. It will often take only the point of the little finger, and, in some instances, will readily receive two fingers as far as the first joints. The fossa normally lodges the duodeno-jejunal bend; and the production of a retroperitoneal hernia, by the protrusion of additional gut into this pouch of peritoneum, can be readily understood.

The plica and its pouch, although not constant, are met with with equal frequency in both males and females, and in individuals of all ages. Among the specimens selected as typical was one from a male

fetus at full term, and another from a woman aged 82. In the 100 bodies examined, I found 48 examples of this fossa duodeno-jejunalis. Among the variations met with in the fossa, the following may be noted.

The pouch may be very shallow, on the one hand (Fig. 3 A), or very long, narrow, and deep, on the other (Fig. 3 B). It may be placed at some distance from the duodenal bend (Fig. 3 C). It may be reduced to a very minute pouch of serous membrane limited to the duodenum alone (Fig. 3 D). Its opening may be in the form of a distinct ring, that in the specimen depicted would just admit a No. 12 catheter (Fig. 3 E). The pouch may be of remarkable breadth, and lodged in a conspicuous transverse fold running from the inferior mesenteric vein to the right hand border of the duodenum (Fig. 3 F). And lastly, the lower part of the pouch may be deficient, and the plica be represented by a simple transverse band running between the usual points of attachment (Fig. 3 G).

Before discussing the origin and mode of formation of the plica duodeno-jejunalis, it may be convenient to consider the relation that it bears to the inferior mesenteric vein. Although this vessel is closely associated with the plica, and although it is well known that many peritoneal folds are caused by the blood-vessels, yet I would venture to express a belief that this particular vein has nothing to do with the formation of this particular fold. It is well known that this vein is somewhat uncertain in its mode of ending, but the 100 specimens that I have examined show that the presence or absence of the plica is in no way influenced by the mode of ending or the disposition of the vein. In the human subject, the vein may end in one of three ways: 1, it may enter the superior mesenteric vein precisely at its junction with the splenic (Fig. 4 A); 2, it may enter the former vein at some distance from the splenic (Fig. 4 C); 3, it may enter the splenic vein at a right angle (Fig. 4 B). In 100 specimens, the vein was disposed as follows: condition 1 = 44 per cent.; condition 2 = 36 per cent.; and condition 3 = 18 per cent. In the remaining two cases, the vein took a remarkable course. It ran vertically upwards in front of the splenic vein, passed between the layers of the transverse meso-colon, parallel with and two inches from the bowel, and, having described an extensive curve, entered the superior mesenteric vein on its right side (Fig. 4 D). This is not unlike the course adopted

by the vessel in some animals, and the abnormal trunk was no doubt due to an enlargement of the communicating branch between the inferior and superior mesenteric veins, that normally runs in a curved manner along the transverse meso-colon. The ending of the inferior mesenteric vein in the splenic would appear to be a feature in the higher development of animals. In the mammalia below monkeys, the inferior mesenteric artery is a branch of the superior mesenteric, and the vein that accompanies the former vessel enters either the trunk of the superior mesenteric vein, or at the point where that vessel joins the splenic. In none of the mammalia below monkeys have I found the inferior mesenteric vein ending in any other way. In the last named class of animals, the inferior mesenteric artery comes off from the aorta, and the companion vein ends most usually in the angle of junction of the superior vein with the splenic. It may enter the trunk of the superior mesenteric vein some distance below the termination of that vessel. In only two instances, namely, in a Bonnet-monkey and in a Silvery Gibbon, did I find the inferior mesenteric vein discharging itself into the splenic.

In man, the inferior mesenteric vein takes a sharp curve before it ends, especially when it terminates in the superior vessel. It passes under

the lower border of the pancreas and behind the end of the duodenum. Before it is hidden by the pancreas, it often draws the serous membrane into a fold, near the free edge of which the vessel runs. This fold will be concave downwards, and will limit a fossa of varying dimensions that may pass up, as a kind of pocket, behind the lower edge of the pancreas. The fossa is sometimes large enough to conceal the thumb up to the root of the nail, and looks directly downwards. It often co-exists with the fossa duodeno-jejunalis, which it then serves to deepen; and is most usually met with when the inferior mesenteric vein ends either in the superior mesenteric, or at the point where that vessel joins the splenic (Fig. 5 B). Two quite unusual forms of this pouch are shown in Figs. 3 H and 3 I.

I believe this pouch to be of no practical importance, to have no interest either with regard to development or comparative anatomy, and to be in no way concerned in the production of internal hernia. I take it to be simply of the same nature as the folds that are frequently formed in the peritoneum by blood-vessels. Good examples of such folds are found about the cæcum, and a striking instance from the peritoneum near the descending colon is shown in Fig. 5 A. To

these vascular folds, and especially to the latter, I shall have occasion to allude later on.

The mode of formation of the fossa duodeno-jejunalis may now be considered; and, to render the account of its formation intelligible, some reference must be made to the general development of the intestinal tract. In its most primitive condition, the alimentary canal appears as a simple vertical tube, running down in the middle line, and connected to the spine by a simple vertical fold of peritoneum. (Fig. 6 A.) In a little while, the upper part of the tube enlarges, and becomes bulbous, and the rudimentary stomach is produced. The lower part of the tube retains its primitive connection with the spine, and retaining also very nearly its original vertical direction, becomes the descending colon and rectum, including a part also of the transverse colon. In fact, it persists as all that segment of the large intestine that is supplied by the inferior mesenteric artery. The intermediate part of the primary tube increases rapidly, and soon forms a large loop, which projects out of the still shallow abdominal cavity, and from which the intestine from the pylorus, to about the middle of the transverse colon, is ultimately developed. There is at first nothing to indicate the separation of the large intestine from the small; but soon a bulging takes place, at a point just behind the apex or middle of the loop; and this, in process of time, develops into the cæcum and its appendix. This large loop has a common mesentery, continued from the original vertical mesial fold of serous membrane that connected the primary intestinal tube with the spine. The loop has a narrow neck, formed above by the duodenum, and below by that part of the large intestine that subsequently becomes the right extremity of the transverse colon. Between these two segments of gut, at the neck of the loop, the superior mesenteric artery runs to supply the bowel. As it runs in the mesentery, it gives off branches from its right or upper side to supply the small intestine, while from its left border arise the vessels for the cæcum and ascending colon. (Fig. 6 B.) The intestine in the loop grows rapidly, and the mesentery increases in a corresponding ratio. That part of the membrane, however, that lies at the neck of the loop does not grow at the same rate. Thus it is, as Professor Flower has well pointed out, that the duodenum and the right end of the transverse colon never lose their primitive relation, in spite of the many subsequent changes that take place in the position and growth of the viscera. They continue to limit and bound the neck of the mesentery; and, as growth proceeds, this neck or root becomes proportionally narrower, while through it still runs the trunk of the superior mesenteric artery. In time, the great loop is withdrawn into the abdomen, and becomes twisted upon itself in a remarkable but definite manner. This twisting is due, in the main, to unequal growth in the two sections of the loop; for, while the small intestine has been

increasing in length with great rapidity, the colon has comparatively grown but little. Just before the twist takes place, the cæcum will probably be lying about the umbilicus, and will be placed, together with the rest of the large intestine, wholly to the left of the middle line.

When the change in position occurs, the cæcum mounts up towards the right hypochondrium; it passes in front of the loop of the duodenum, and ultimately descends to its final resting place in the right iliac fossa. As a result of the twisting, the small intestines are turned towards the left side; what was originally the right side of the mesentery becomes the left side; and the vessels to the small intestine are seen to come off from the left border of the superior mesenteric artery, instead of from the right. The mesentery has been rotated, in fact, half a circle. At the narrow neck of the great loop, the changes are less conspicuous. The right end of the transverse colon has passed in front of the duodenum; but they still form the boundaries of the narrow neck of the mesentery, and between them still passes the superior mesenteric artery. As development proceeds, the bowel grows somewhat out of proportion to the peritoneum about it, and this disproportionate growth is well marked in the two segments of bowel that form the neck of the great intestinal loop. As they grow, they become too large for these serous coverings; they to a great extent grow out of them, and thus it happens that the duodenum and transverse colon are brought nearer and nearer together, until at last their relations are very intimate, although their mutual positions have become modified. The development of the duodenum itself may now be considered. This part of the gut forms a loop of its own; a loop that starts from the pylorus, and ends where the gut becomes fixed by the musculus suspensorius. Its termination, therefore, is at the neck of the great general intestinal loop, and close to the trunk of the superior mesenteric artery. The duodenum, which is at first comparatively of large size, has its own meso-duodenum, which is attached vertically to the middle line, being a part of the original mesial mesentery. When the pylorus moves to the right, the loop of the duodenum moves with it, so that the left layer of the meso-duodenum becomes anterior, and the right layer posterior. When the twist takes place in the intestinal loop, the duodenum has no share in it. The general position of the loop remains unchanged. The end of the duodenum is dragged across the middle line from the right side to the left; the superior mesenteric artery passes over it instead of by its side, the dragging produced by the upward movement of the colon causes the terminal part of the duodenum to become vertical, and a sharp twist is formed in the gut where the duodenum and jejunum meet. The meso-duodenum and the mesentery are no longer continuous in the same plane. In time, the duodenum loses its mesenteric fold, partly because it grows out of proportion to the peritoneum, and partly because developing viscera in its neighbourhood draw the serous covering from it, and ultimately it actually acquires a large non-peritoneal surface.

The question now arises, as to the origin of the fossa duodeno-jejunalis; and that question can be best considered by leaving for a while the development of the parts in the human fetus, and picking up the thread among some of the lower mammalia. What has been said of the development of the intestines in man will apply equally in all general points to the rest of the mammalia. We are, however, especially concerned now with the duodenum. In nearly all mammals—with the exception of the monkeys—the duodenum retains its mesentery. This fold is often very extensive; it is not in the same plane with the mesentery of the small intestine; it is attached vertically along the middle line, and its posterior or right layer is continuous above with the corresponding layer of the gastro-hepatic omentum. It persists, in fact, as a part of the original primitive vertical serous fold. The duodenum ends, as in man, to the left

of the middle line, or rather to the left of the trunk of the superior mesenteric artery, and where it ends the bowel undergoes a very abrupt and pronounced twist. As a fairly typical example of the condition of parts, I have selected the duodenum of a hyena, and the drawing is from a dissection of an adult female (*Hyæna striata*), Fig. 7. It will here be noticed that from the terminal part of the duodenum a vertical fold comes off, which is attached precisely along the middle line, and which, indeed, arises from the same line as the descending meso-colon.

So far as I can ascertain, this fold has been barely noticed by anatomists, although it is almost constant where a meso-duodenum exists. The fold is singularly free from blood-vessels, and is attached to the duodenum along a line precisely opposite to the attachment of the meso-duodenum. It ends below in a free fold that is concave downwards. I have seen a like fold in a small human fetus, in whom the meso-duodenum had not yet been obliterated. The fold is evidently nothing more than the continuation back to the spine of the meso-duodenum. The terminal part of the gut has been drawn up between the folds of the serous membrane, partly by the effects of its own growth, partly by the traction of the rapidly developing small intestine. The free edge of the fold represents the inferior limit of the meso-duodenum before the terminal part of the gut has become vertical. Owing to the enormous development of the duodenal loop as compared with that of the lower colon, the meso-duodenum, that was originally above the meso-colon and in the same vertical line with it, has descended by the side of the latter until the two have at last become parallel. They are, indeed, fixed together at the spine. In many animals, owing, I imagine, to continued growth of the meso-colon, this duodenal fold has been carried with the latter from the spine; so that in such cases it appears to be inserted in a vertical line to the right layer of the descending meso-colon at a certain distance from the column. I venture to think that it is from this fold that the plica duodeno-jejunalis is produced.

In the first place, I may state that I have found no trace of this plica, nor of the fossa it produces, in any one of the animals I have dissected. In two of the carnivora, namely, in a Puma and in a Cape hunting dog, I found a fossa in the peritoneum, at a point where the meso-duodenum and the mesentery met. The fossa was placed transversely, was at some distance from the spine, was apparently due to the rotation of the bowel, and had no resemblance to the duodenal fossa in man. It would appear that the fossa follows upon the obliteration of the meso-duodenum. This obliteration depends, in part, upon the growth of the duodenum itself, and, in part, upon the rearrangement of the peritoneum during the progress of development. The proportion of peritoneum to intestine is much greater in the lower animals than it is in man. In most of the mammalia, there is an extensive mesentery and a meso-duodenum, the ascending colon is free, the transverse colon supplied by a liberal fold, and the descending colon attached by means of a large meso-colon. In man, on the other hand, the ascending colon has usually a large non-peritoneal surface. It has outgrown its serous covering. The descending colon, moreover, has been removed far to the left of the middle line, and very commonly has no meso-colon. Even when such exists, it will be no longer attached along the middle line in the place of the original fold. In most apes, the descending meso-colon, which is always present, is still attached practically along the middle line. In the Gibbon its attachment is transferred to the inner edge of the kidney, but in no ape that I have so far seen has the peritoneum been encroached upon to the extent that is so noticeable in man. In the human fetus, the meso-duodenum is probably obliterated by the downward growth of the cæcum. This part of the colon, as it develops, would drag upon the peritoneum, and would tend to obliterate all folds. More peritoneum is required by the cæcum and ascending colon, and it is obtained from that of the posterior parietes, and, in great measure, by the unfolding of the meso-duodenum. By such unfolding, the posterior layer of the meso-duodenum is dragged away from the loop, which now lies with its hinder surface bare; and what was once this posterior layer comes to line the adjacent part of the parietes below and to the outer side of the duodenum. The anterior layer of the meso-duodenum remains. It will be seen that, by such unfolding, the vertical fold from the duodenum (Fig. 7) will become obliterated; and the condition may thus be brought about that is met with in most apes, and in such human specimens as are marked by the absence of the plica duodeno-jejunalis. But the dragging of the peritoneum is not all in one direction. The left part of the colon is making almost equal claims upon the serous membrane, and is tending to drag it from the posterior parietes towards the left side. In this way, the vertical fold would be carried towards the left, and be moved in an upward direction; and would persist, if the demands upon the peritoneum

case, as the plica duodeno-jejunalis. The free edge of the plica will correspond to, although it will not be identical with, the free edge of the vertical fold. It is not necessary to assume that the duodenum moves with the shifting serous membrane. Repeated illustrations are given elsewhere, to show that peritoneum can be displaced without involving equivalent displacement of the viscus that it covers.

It is significant to note that both the plica and the vertical fold are attached to the same part of the duodenum, and along precisely the same line. They are both, moreover, conspicuous by their thinness, and by the absence of visible blood-vessels.

Several instances have been reported in which, as the result, probably, of intra-uterine peritonitis, the intestines have continued to occupy in the adult the position assumed in early foetal life. In such cases, the colon is found wholly to the left of the middle line, while the small intestines occupy the right side. The bowel has not been rotated in the usual manner, and the vasa intestini tenuis still arise from the right side of the superior mesenteric artery. The duodenum is quite to the right of the spinal column. Abnormalities of this description have been noted by Sir James Simpson (*Edinburgh Medical and Surgical Journal*, 1839, p. 26), Dr. Hilton Fagge (*Guy's Hospital Reports*, vol. xiv), Mr. Lockwood (*BRITISH MEDICAL JOURNAL*, vol. ii, 1882, p. 574), Professor Chiene (*Journal of Anatomy and Physiology*, 1868, p. 15), Dr. John Reid (*Edinburgh Medical and Surgical Journal*, 1836, p. 70), and Dr. Bruce Young (*Journal of Anatomy and Physiology*, 1884, p. 98). In the description of the parts given by the last named observer in a very able paper, this significant passage occurs: "Extending downwards from the inner side and apex of this curve of the duodenum, in the vertical antero-posterior plane, was a membranous layer, about two inches broad and one inch long, which lay two inches to the right of the middle line of the abdomen, and was attached below to the upper surface of the mesentery." Dr. Bruce Young thinks that this fold was a morbid adhesion; but one cannot fail to be struck with its resemblance to the vertical fold met with in animals, and which one would expect to find in this particular abnormality of the bowel.

It has been already said that the fossa duodeno-jejunalis has been credited as the cause of retroperitoneal hernia. If such a rupture form, the pocket becomes enlarged, and lodges more and more of the small intestine. The bowel finds its way behind the peritoneum; and in a complete case, such as that described by Sir Astley Cooper, the whole of the small intestine, with the exception of the duodenum, is hidden from view, and occupies a large sac that is formed in the middle of the abdomen, and surrounded by the large intestine.

Sir Astley Cooper described, however, another form of retroperitoneal hernia, to which he gave the name of the mesocolic. In this rupture, the small intestine was contained within a sac that formed a tumour to the left of the middle line, and that had been evidently developed at the expense of the peritoneum leading to the descending colon. There are three significant facts about the form of hernia that require notice. In the first place, the orifice of the sac was some way removed from the duodenal bend; in the next place, the commencement of the jejunum was not involved in the rupture; and thirdly, a branch of the inferior mesenteric artery ran along the free anterior margin of the orifice of the sac. Now, Treitz and others have endeavoured to show that this form of hernia also is developed from the duodenal fossa. If such were the case, it is hard to understand why the orifice should become so much displaced, and why the commence-

ment of the jejunum, that is theoretically the first segment of the gut to enter the sac, should be entirely free of it.

Treitz and his followers endeavour to explain this mesocolic hernia by asserting that the duodenal plica is often turned to the right, and the jejunum to the left. To aid the explanation they describe certain remarkable arrangements of the "horns" of the plica.

With regard to these statements, which are faithfully repeated in all accounts of these retroperitoneal hernia, I can only say that I have never found the plica or its fossa turned either to the right or to the left, nor have I ever met with the remarkable arrangements of the fold that have been so carefully described as essential to the production of some varieties of this hernia.

My statement is based upon a careful examination of forty-eight plicae, of each of which a drawing was made as the parts lay exposed *in situ*. In no instance was the upper margin of the plica other than horizontal, and all the chief deviations from the typical arrangement I have already described. I venture to think that the mesocolic hernia of the type described by Cooper, may be readily explained by a reference to Fig. 5 A, which was taken from the body of a female foetus at full term. It will here be seen that a deep pouch has been formed in the parietal peritoneum, leading to the descending colon by a branch of the inferior mesenteric artery. The mouth of the pouch is directed upwards. Such a fossa could readily engage a loop of small intestine, more readily even than the duodenal fossa; and, when a retroperitoneal hernia so originating had been fully formed, then the three features that I have already alluded to would be clearly marked; that is to say, the orifice of the sac would be some way from the duodenum, the commencement of the jejunum would not be involved, and a branch of the inferior mesenteric artery would be found skirting the orifice of the sac on its anterior aspect. Folds and pouches such as this, formed by blood-vessels, would appear to be not uncommon. I have met with two such in the mesentery of two fetuses at full term. They were in both instances formed by the ileo-colic artery, and in both examples the fold of the serous membrane bounded a not inconsiderable fossa, the mouth of which was directed downwards.

THE BROWN LECTURES ON PATHOLOGY.

Delivered at the University of London, December, 1884.

By VICTOR HORSLEY, B.S., M.B., F.R.C.S.,
Brown Professor of Pathology to the University.

LECTURE III.—TRAUMATIC FEVER: PYREXIA FOLLOWING SIMPLE FRACTURES.

ACCORDING to received notions, as far, indeed, as they are comprehensible, numerous kinds of fever, of utterly different natures, have for many years been grouped together under the term surgical or traumatic fever. Putting aside the first term, surgical fever, which has nothing in the world to recommend it, and everything to lead us to get rid of it, let us examine the second, as it has formed the subject of many monographs, to the substance of which I shall have to refer in describing my own observations.

The expression traumatic fever has invariably been employed to denote the febrile state which usually follows an injury to the body, with or without a lesion of the skin or mucous membranes. Now, if this febrile state were always the same, that is to say, if it invariably succeeded the injury at the same interval of time, and ran a course of the same character, although it might be of different degrees of severity, then we might justly say that here was a condition which might be termed traumatic fever, subject, of course, to certain considerations to be enumerated directly. But, at present, the term embraces fevers of widely different incubation-periods, and varied modes of progress and conclusion. This is easily comprehensible when we find that, in a paper written as lately as a few years ago, Lucas Championnière describes puerperal fever, pyæmic fever, that of septiciæmia, and urethral fever, all under the same title, namely, traumatic fever.

The most recent scientific work on this and kindred subjects (Brisson's *Science and Art of Surgery*, 8th edition, 1884) divides the septic and urethral fevers from the pyrexia following a simple injury under the headings septic, traumatic, and urethral fevers respectively.

Now, at the outset, I venture to suggest that the term traumatic fever should be restricted solely to those instances in which it can be