tasted of nastiness to her. All these things no doubt were symptoms of disturbed action of the special sense-areas of the cortex.

Those perverted painful sensations are not confined to the head, or to the special senses. They are referred in some case to the chest, to the abdomen, and to the limbs. Some melancholic patients complain of pains in the region of the heart, and some of feelings of constriction in the chest, as if they could not get breath. Most frequent of all sensations, except those in the head, are sensations of sinking and pain, of fulness, of "anxiety" in the epigastric region. Certain forms of melancholia are so characterised by abdominal symptoms, that they may be fitly called "visceral" cases of the disease. Such organic sensations of discomfort often precede the mental depression like the headache, but, unlike this, they seldom pass away when the mental pain begins. Some patients can tell when they are going to have an attack of melancholia, from the beginning of such sensations.

Spinal pains and discomfort, and even lumbago, are common. I have now a case of aggravated resistive melancholia under my care, whose attacks have always been preceded, and their earlier stages accompanied, by attacks of lumbago, which passed off when the mental pain became intense.

Pains resembling sciatica are not uncommon in the legs before melancholia supervenes; while feelings of stiffness and of cramps in the limbs I have also known. Perverted sensations with regard to the size of the head and limbs are by no means uncommon. I had a patient who said he felt as if his head were enormously large, and another who felt as if his legs had grown quite diminutive.

Perverted sensations of heat and cold are common. One melancholic says he has a constant feeling of cold, another that he feels hot subjectively—the actual bodily temperature in both being just under normal, as is usual in most varieties of melancholia, except the excited form. Sensations of prickling, of formication, of itching in the skin are not uncommon. I have seen many patients, again, who complained of "a want of feeling all over the body, a deadness," as if they carried about a corpse instead of a living body.

I have never been able to believe that such painful and perverted sensations commonly originate in vascular abnormalities. On the contrary, the facts seem to point to a purely nervous origin, but no doubt they are always accompanied, and commonly kept up, by vaso-motor abnormalities; though the facts of megrim and other neuroses seem to point to pain being readily caused, in some cases, by vaso-motor dilatation and paralysis.

My recent experience, since my attention was strongly directed to the close relationship of mental and bodily pain, has been that in nine out of every ten cases of melancholia, if carefully questioned during and after the attacks as to their sensory history, we shall discover, as a clinical fact, painful perversions of bodily sensibility in some shape or other, standing preliminary to or accompanying the attack. Melancholia is not a mere mental disease; it is a general disease of the brain-cortex, usually implying a combination of mental and bodily symptoms, the latter chiefly consisting of neuroses of sensibility, and commonly accompanied by vaso-motor trophic neuroses.

Many every-day facts show that mental and bodily pain can take the place of each other, and in many cases, indeed, cannot co-exist in any great intensity. I have known a painful burn stop the mental pain for the time in melancholia. All of us have seen the transient good effect in some cases of a blister; while, on the other hand, bad news, or even the painful mental anticipation of the dentist's forceps, will often cure toothache. Sometimes a normal and physiological mental pain, such as the news of the death of a child, will inhibit and cure the abnormal pain of melancholia.

Another relationship is frequent. Severe bodily pain long continued will frequently bring on melancholia. I have seen two cases lately of ladies, who, subsequently to painful surgical operations, became melancholic. I have seen melancholia come on after a painful attack of rheumatism. In such cases, no doubt the nutrition and the general nervous energy would be thus run down.

There can be no question whatever as to the general similarity of many of the typical cases of hereditary neuralgia and of hereditary melancholia in the bodily condition of the patients, both being thin, both bad sleepers, bad digesters of their food, both over-sensitive, both intellectual but wanting in staying power for continuous mental work and in surplus energy bodily and mental, and both manifest, in certain cases, a marked tendency to diurnal aggravations. Both may arise from local sources of irritation, bodily and mental. Both inhibit volitional action and energy, emotion and sexual feeling. And, unquestionably, the treatment found useful for both diseases is the same; change of air and scene, a generous fattening diet, cheerful company, systematic regimen and division of time, iron, quinine, tonic nerve stimulants, volitional energy rightly roused, and fixing the attention

on objective things of an agreeable kind. In short, whatever nourishes the body, increases the red blood-corpuscles, and causes functional hyperæmia, and improved nutrition of the brain-cortex, cures both diseases.

In certain hereditarily neuropathic families it is very common for one member to be subject to neuralgia, and another to melancholia; while we have all seen hereditary connections between the two diseases, such as a neuralgic mother having a melancholic son. I have seen several cases now of circular neurosis, where an intolerable neuralgia formed one part of the circle, and a deep mental misery formed another.

In hypochondriasis, we have a disease where there is neither acute bodily nor intense mental pain, but a combination of both in a modified and mixed-up way. No one can doubt who has seen a strong intellectual man pass through an attack of hypochondriasis, or of melancholia of the hypochondriacal type, and seen him recover again, that these were real diseases, caused by disordered energising of the brain-cortex, however much the symptoms during the attack seemed mere "imaginings," and the strange conduct appeared controllable by the will, if the patient had chosen to exercise it. But there are few mental diseases where a man's normal volition, his feelings, his habits, and his power of working, will be more inhibited than in this. The pains and discomforts, the imaginary disorders of function, and the general sense of organic ill-being of the hypochondriac, I look on as resulting from an equal disturbance of the mental and sensory portions of the brain-cortex; just as in ordinary melancholia there is an unequal disturbance, the mental portions being most affected. No doubt, the areas that control and specially represent the organic viscera, and their special nerve-ganglia are specially disturbed in function in these cases.

From the prophylactic point of view, the general association of melancholia with sensory disturbances has a special value. Some pain or sensory perversion usually preceding the mental disease, the physician can take early measures of prevention. How many attacks of melancholia could be averted, if proper treatment were only adopted in time?

It appears to me that physicians in general practice frequently do not realise the significance of slight mental depression, accompanying or following neuralgia or sensory perversions, and that psychiatrists often do not sufficiently think of the sensory symptoms in the study and treatment of melancholia. The borderland between neuralgia and melancholia I look on as a very slight one indeed. They are both manifestations of one type of the neurotic diathesis. The seat of both is in the brain-cortex, and in the same part of the cortex, though not in the same cells. The actual cortical conditions under which the mental and the bodily pains arise would appear to be the same; therefore, it seems clear that they should not be studied or treated apart from each other.

AN ADDRESS

DELIVERED AT THE OPENING OF

THE SECTION OF PATHOLOGY,

At the Annual Meeting of the British Medical Association held in Brighton, August, 1886.

By JULIUS DRESCHFELD, M.D., F.R.C.P.,

Professor of Pathology, Victoria University; Physician to the Royal Infirmary, Manchester; President of the Section.

THE RELATIONS OF PATHOLOGY; AND ITS STUDY

GENTLEMEN,—In taking the chair in this section, I wish in the first place to express my sincere thanks for the great honour conferred on me by my election to the presidency, a post which has been held by some of the most eminent men in this country. In spite of all my endeavours, I am afraid I shall be but a feeble successor to them. I will, however, do my best to prove myself worthy of the confidence placed in me; and I am sure I shall not ask in vain for your kind indulgence.

My first, and by no means easiest duty, is to say a few words introductory to our work. In doing so I have avoided taking up any special subject of pathology, as the discussions and papers already announced provide ample material in that direction. Moreover, as our JOURNAL gives from time to time an excellent retrospect of recent pathological researches, I do not propose to occupy your time with such a retrospect on the present occasion. I will only ask your attention for a very short time while I offer a few observations on the

relations of pathology to the other branches of medical science, and on the study of pathology in England compared with other countries.

There is no branch of medicine which has made such rapid strides, and which covers so large a field as pathology, and for reasons which are sufficiently obvious. Pathology, as the study of disease, stands in the same relation to the diseased organism as anatomy and physiology do to the healthy organism.

Pathological anatomy makes us acquainted with the changes in structure which a diseased part undergoes, and pathological physiology considers how and why the functions of such a part become altered. But pathology has also to inquire into the cause of disease; and it is in this particular branch—the etiology of diseases—as I need scarcely tell you, that the greatest activity is being displayed, and that the most important discoveries are being made.

Of these different subdivisions, pathological anatomy, the oldest of them, has received the largest share of attention. It may be said to have been really founded by John Hunter, who has bequeathed to England a museum superior to any in Europe, and which, thanks to the labours of Sir James Paget and his able fellow-workers, has been richly supplemented, and has kept pace with the advancement of pathological science. The study of the minute structure of the diseased organs has gone hand-in-hand with the progress of normal histology and embryology; and, though many points still require clearing up, a considerable number of the great problems of general pathology have been solved, and the gaps yet to be filled up are becoming smaller and smaller. It behoves the pathologist, however, to watch closely the progress of histology, for there are few discoveries in general and special histology which will not find their application in pathology. Take, for instance, the recent researches on the structure of the cell-nucleus, and its changes during the normal processes of cell-life. It has been shown by the older observations of Flemming, Klein, and others, supplemented by the more recent ones of Heuser and Rabl, that the nucleus of a cell—the most active and vital part of a cell—is a composite structure, made up of several constituents, of which one, chromatin, is morphologically the best defined, and which, during the process of cell-division, shows those remarkable changes of form known as karyokinetic figures. Pathology is already beginning to profit by this new line of research. Thus, the observations of Pfitzner show, among other things, that the cancer-cell has an embryonic character; its nucleus is much poorer in chromatin than the nucleus of a normal cell.¹ Again, according to Baumgarten's observations on the karyokinesis seen in the artificially produced tubercle of the cornea and iris, and Scheltema's researches on the inflammatory changes in the subcutaneous tissue, it becomes evident that, in inflammation, the fixed tissue-corpuscles play a much more prominent part than we have hitherto assumed.

I may be allowed to allude to another histological method, introduced quite recently by Ehrlich, which has opened out a new field for pathological research, and may lead to important results concerning the toxic action of many substances. Ehrlich found that methylin blue, when introduced into the living animal body, possesses apparently a peculiar predilection for certain nerve-terminations and nerve-elements, which it stains; while parts more or less analogous, such as the motor nerve-terminations, it leaves unstained. Ehrlich believes that this peculiar behaviour of the nerve-elements is partly due to the richness of oxygen, partly to their alkaline reaction. But even if this explanation be considered as yet too theoretical, this method, if further developed, may give us some insight into the more obscure phenomena of cell-life, the reaction of cells and their specific characters, and may throw further light on some of those peripheric paralyses due to chronic intoxication, which is to form the subject of one of our discussions.

Pathological anatomy, by itself, teaches us more than the mere appearances of the tissues or organs at the time of death; it often enables us to trace the successive steps of the process of disease, and may even lead us to recognise the cause of disease. No doubt the truth is only arrived at after long and tedious labour; often we have to retrace our steps, and often we have to unlearn what we have learned; but in this respect pathological anatomy does not differ materially from other experimental sciences. Nothing illustrates this better than the history of tuberculosis. Laënnec taught the unity of tubercle and phthisis. For many years, pathologists following the lead of Virchow and others, distinguished between tubercle as a new growth and phthisis as the result of inflammation. More minute examination into the histology of tubercle, of phthisis, scrofulosis, and lupus, showed their close relationship, if not actual identity; and independent observers, amongst whom Wilson Fox took the lead, preached

¹ It is interesting to note here that Savoy arrived at a similar conclusion by a somewhat different route.

again the doctrine of unity of phthisis and tuberculosis, and even before Koch demonstrated the actual cause of tuberculosis, and showed the same bacillus to be present in phthisis, scrofulosis, and lupus, Villemin, Cohnheim, and others believed the cause of tubercle to be some infective agent.

Much more productive than pathological anatomy in the extension of our knowledge of disease, has been pathological physiology or the experimental investigation of disease, a branch of research in which English pathologists are unfortunately prevented at present from successfully competing with their continental brethren. The importance of the pathological experiment did not escape the old observers, and here I may again cite Hunter as one of the pioneers. The commencement of experimental pathology, however, dates from the rise of experimental physiology, with which it is closely allied, and with which it has many points in common. The close relationship of these two sciences was so ably and forcibly brought before this Association by Professor Michael Foster, in the address he gave at the Cambridge meeting, and again by Professor Greenfield, at the Belfast meeting, that no further comments of mine are necessary. There is scarcely any subject of experimental physiology which has not found its application in pathology; and whether we take the researches on the function of the liver, the investigations on the functions of the kidney, the observations on the innervation of the heart, or the numerous experiments on the functions of the different parts of the nervous system, they have not only taught us the normal functions of the body, but thrown light on the morbid processes which underlie disease; hence, also, on its diagnosis.

Experimental inquiries are even of greater necessity in pathology than in physiology; and in support of this assertion I need only point to the observations on inflammation, thrombosis, and fever, and to the whole chapter of the so-called bacterial pathology, culminating in the most recent and most important researches of Pasteur on hydrophobia. It is especially the study of the relations of micro-organisms to disease, prosecuted with such vigour on the Continent, which has already given, and is likely to give, the most valuable results, not only to pathology, but also to hygiene and therapeutics.

We now possess reliable methods for the detection of micro-organisms; we know a good deal already of the specific character of some of them, the progressive virulence of some, and the innocuous nature of others. We have already learned how the virulence of some can be weakened, and how the organism can be made capable of resisting their poisonous action; and yet, though the pathological experiment has been shown not only to be of scientific interest, as furthering the progress of pathology, but also to have practical bearing on the prevention and cure of disease, we pathologists in England, from restrictions placed upon experimental research, labour under the greatest possible difficulties. As the law relating to vivisection stands at present, experimentation concerning some of the most important inquiries is impossible.

One instance I may perhaps be permitted to give. We had quite recently a case of hydrophobia in the Manchester Infirmary, and an opportunity was thus afforded to test one of Pasteur's assertions, namely, that if a portion of the fresh medulla from a patient that had died of hydrophobia be injected beneath the dura mater of a rabbit, death ensues in the rabbit after a certain time from a peculiar form of hydrophobia. As, from the date of admission to the death of the patient only two days elapsed, there was, of course, no time to apply for a licence, and hence the experiment could not be attempted. Now, as hydrophobia is for the time almost extinct in Germany and Austria, while in London alone there occurred, during the first ten months of last year, no fewer than sixteen cases of hydrophobia, Pasteur's inoculation experiments are of much more direct value to us than to the other large European countries; and experimentation from different independent observers is highly desirable.

In the application for a licence, the applicant has to state fully the reasons why he wants to conduct certain experiments, and to give the exact methods he wishes to employ. Now, although it is quite right that an experimenter should know exactly the object of his inquiry, and the methods he wishes to pursue—and it is well known that Claude Bernard, the *déte noire* of the antivivisectionists, and yet one whose contributions to physiology and pathology mark an era in both sciences, always had a full and detailed plan formed before he proceeded to his experimental inquiries—yet it will often happen that new points and different methods will suggest themselves as an experimental investigation proceeds, and yet, in consequence of the restrictions at present existing, it is impossible to deviate from the plan on which the licence is granted. In the interests, not more of the progress of medicine than of humanity, it is highly desirable that some of the restrictions should

be relaxed; and if experimentation were only allowed to be carried on in the physiological laboratories and in the few existing pathological institutes, we might yet hope to compete successfully with continental observers in this fruitful line of inquiry. A powerful society like the British Medical Association, which encourages original research so much, as seen by the foundation of research scholarships; and the grants given for original investigations, may yet, I hope, be able to bring about a relaxation of some of the restrictions in the laws regulating vivisection; and although a powerful appeal has already been made in vain, if another effort be made, especially as the public must now better appreciate the benefits conferred on humanity by experimental methods of research, I think a fresh appeal might not be altogether fruitless.

Fortunately, pathology covers a wider field, and includes more than pathological experimentation; it is closely connected with clinical medicine and surgery, and, in this association, it has been studied as diligently and as successfully in England as on the Continent.

Take, for instance, the pathology of the nervous diseases, in which such enormous progress has been made within the last ten to fifteen years, and where, amongst the many investigators, M. Charcot, whom we gladly welcome here to-day, stands out first and foremost; here careful clinical investigations, coupled with accurate *post mortem* observations, have brought knowledge, system, and order where there was before so much obscurity and disorder. These investigations have not only given us important pathological results, but physiology has largely gained by them; for our knowledge obtained by observations on man of many physiological questions, such as the trophic nerves, the localisation of centres for different functions, rests on a much more satisfactory basis than the results of experimental enquiries, though I do not by any means wish to depreciate the great value they also have. In fact, in every experimental science we require different methods, in order that one method may check another.

Again, the pathology of Bright's disease, which has been so extensively and carefully studied in this country, can only be correctly arrived at by a combination of exact clinical observation and accurate anatomical investigation; though here, too, experimental pathology has contributed much to the elucidation of many obscure points, and will yet solve some of the more intricate phenomena in connection with the subject. But even apart from *post mortem* examinations, there is a bed-side or clinical pathology in which English physicians have taken a leading share; as, for instance, the study of the altered arterial tension in certain diseases, which is of great therapeutic value, and has already led to important therapeutic results. Again, the causes and processes which underlie some forms of dyspepsia are now being better known by help of exact clinical observations; we know now that, in most cases of nervous dyspepsia, gastric digestion is completed in the normal time, and many of the symptoms are due to reflex irritation, in others there is increased secretion of hydrochloric acid in the stomach, whilst in others the gastric juice is deficient in hydrochloric acid, or contains some other acid, lactic or acetic, in abnormal proportions. For our knowledge of the so-called functional diseases, such as hysteria, and for many chapters of psychophysics, we are as yet entirely dependent on clinical observations. The work to be done here is so extensive that it has led to the establishment of many clinical laboratories on the Continent, for which, as far as I know, no adequate provision has yet been made in our large English hospitals; though I do not wish to ignore the fact that a good deal of such work is carried out privately by some English physicians in their own work-rooms.

From all the foregoing considerations it is evident that pathology covers an immense field; the question how it is best studied, and best taught, is by no means so easy to answer.

It is essential that the pathologist should keep himself thoroughly *au courant* with the progress of histology and physiology, and that he should be fully acquainted with what we have called clinical pathology. Should, then, the pathologist devote his whole time and energies to the *post mortem* room and pathological laboratory, and abstain altogether from active practice or clinical work? Would it not be more advantageous if he could still be engaged in clinical work? Theoretically, there is no reason why he should not be actively engaged in the exercise of his profession; indeed, for one line of pathology, clinical observation is absolutely essential. If we look round, we find that a great deal of the pathological work is done by clinicians; take, for instance, the pathology of nervous diseases, the relations of micro-organisms to diseases, and most chapters of surgical pathology. Our recent progress in the knowledge of these subjects we owe chiefly to the indefatigable labours of men engaged in clinical practice; and yet there are many and obvious reasons why there should be

pathologists who should confine their attention to pathology alone. The chief reason is that the whole field is too large for one man to take up; and, another reason that, if not allured away from his laboratory, the pathologist can devote much time to teaching, to original research, and to the assistance of those who are actively engaged in their profession, in the working out of problems which present themselves to them in the course of their clinical observations.

Nothing is more useful to the medical practitioner than a thorough knowledge of pathology, and nothing is more likely to make him a careful and thoughtful observer, and to stimulate him to work. In a science like ours, where we have to do at first with a number of detailed inquiries, from which eventually generalisations are deducted, everyone may contribute his quota. Koch laid the foundation of his great discoveries when still a general practitioner, and Stilling did some of his best work in a small work-room, when not yet engrossed by a laborious practice.

If we look at the many excellent clinicians which England has produced, we find that most of them have spent their earlier years in pathological work and pathological research. Amongst the long list of names, many of European fame, I may take one as an example, one who stands out pre-eminently, and who, even by his latest publications, shows us that he is as great a pathologist as he is a surgeon; I mean Sir James Paget. Now, whilst fully admitting the vast amount of pathological work done in England, as seen by that mine of wealth on pathological subjects, the *Transactions of the Pathological Society*, yet one cannot help feeling that still more would be done if we had institutes like those on the Continent, and if pathology entered more largely into the curriculum of our medical study. I think it will be admitted on all hands that it is chiefly by virtue of an arrangement for the complete and systematic study of pathology and physiology, that scientific medicine in Germany has received such a powerful impetus, and that it is the want of these facilities here in England which will make it difficult for us to keep up the race with our neighbours across the Channel.

What we then require in England, above all, is the establishment of well-endowed chairs of pathology, in connection with our leading medical schools. I know the objections there are to State help for such purposes, and the difficulty there exists for obtaining such assistance from the Government; yet, what has been done in recent years for physiology could also be done for pathology, and the example of Cambridge should be followed by the sister University and some of our large medical schools. At present, though the teaching is excellent, we miss the systematic instruction in pathology; the lectures in pathology are insufficient in number, and the teaching in the *post mortem* room is often left to inexperienced demonstrators. Yet there is no country where it is more essential for the ordinary practitioner to be well versed in *post mortem* examinations than in England, where every practitioner may be called upon, in medico-legal cases, to conduct necropsies, and give medical evidence thereon in a court of justice—a system which is very much to be condemned, for I feel sure it would be much better if the medico-legal *post mortem* examinations of one district were conducted by one well qualified person in a centrally situated institution, where there would be the necessary appliances for a detailed and careful examination, and to which students could also have access. The ends of justice would in this way, I think, be better served, and the advantages of such a system, from a scientific point of view, are equally evident.

If we had well appointed teachers of pathology, it would also soon follow that pathology proper would enter more largely into the curriculum of medical students, and form a more prominent part in the examinations for the various diplomas and degrees. Thus there is no separate examination in pathology at any of the examinations, either at the College of Surgeons or College of Physicians, and even the London University has no separate examiners in pathology, and joins together general pathology, general hygiene and therapeutics, all in one examination paper. It is a source of satisfaction that the Cambridge University has not only established an endowed chair of pathology, but gives also due weight to pathology in the medical examination; and I may be pardoned if I allude to the Victoria University, where separate examinations are held, both in general pathology and in pathological anatomy, and where the candidate for the degree has to bring proofs of having studied pathology practically in a laboratory.

However much legal restrictions may interfere with laboratory work of experimental character, yet the success of our Scotch universities, where pathological teaching is carried out much more systematically and thoroughly than with us, and the excellent work done in the Brown Institution, show that much may be achieved even in spite of these obstacles.

Let it not be supposed that the improved teaching of pathology would add much to the labours of the already overburdened medical student, or that it would have the tendency to make him a pathologist rather than a practitioner of medicine, whose chief duties, after all, will be to heal the sick. It would merely systematise his studies; he would be better prepared for his clinical work, and it would be of the greatest advantage to him afterwards in the exercise of his practice; for, as one of my predecessors in this chair has rightly said, "Pathology is the basis of every rational system of therapeutics."

After these few somewhat disjointed remarks, which express not only my views, but also the views of many practitioners whom I constantly meet, and who appreciate fully the advantages of a systematic training in pathology, I will ask you to turn your attention to the work of this Section, which, both as regards the subjects for discussion and the papers to be read, promises good results. In selecting subjects for discussion, we cannot be said to be behind the age, for as nerves and micro-organisms occupy now-a-days so much of our attention, peripheral neuritis, one of the most recent chapters of nerve-pathology, and the etiology of pneumonia, a disease which by many is now held to be parasitic and infectious, may be considered very fashionable subjects. In the third subject for discussion, aneurysm, we see the close relation of pathology to practical medicine and surgery. These discussions, in which some of our best and ablest workers have promised to take part, will leave but little time for the reading of papers; but these, though small in number, yet cover a very large ground, and are sure to add largely to the useful work on which this Section is about to enter.

AN ADDRESS

DELIVERED AT THE OPENING OF

THE SECTION OF PHARMACOLOGY AND THERAPEUTICS,

At the Annual Meeting of the British Medical Association held in Brighton, August, 1886.

By THOMAS LAUDER BRUNTON, M.D., F.R.S.,

Lecturer on Materia Medica and Therapeutics at St. Bartholomew's Hospital
President of the Section.

ON THE CONNECTION BETWEEN CHEMICAL CONSTITUTION AND PHYSIOLOGICAL ACTION.

THE meeting of the British Medical Association is not for mutual instruction only; it is also for recreation; and, probably, many members of this Association will utilise the opportunity which a meeting at the seaside, like the present one at Brighton, affords them of indulging in that excellent occupation for an idle man, of watching the waves on the seashore, and speculating how far each of them will come. If one have only half an hour to spare, it is difficult to say whether the tide is ebbing or flowing; it is only by watching for a longer time that one can be certain that the water is really moving in one direction or another. Probably, a great part of the charm which this occupation possesses is due to the resemblance which one involuntarily traces between the ebb and flow of waters and that of human affairs—individual, national, or racial. The life of a single man is very short in comparison with the history of race; and it is often very difficult to say whether mankind is advancing or retrograding, unless we compare his condition at epochs widely removed from one another.

On doing this, we find a general consensus of opinion, to the effect that civilisation has steadily advanced; and this advancement is usually divided into four stages, characterised by the nature of the tools or weapons employed. In the first, or Paleolithic age, man employed weapons or tools of flint roughly chipped into shape and unpolished. In the next, or Neolithic age, the implements consisted of stone, but they were polished. The next age is characterised by the employment of bronze as a material, and the fourth and highest stage by the employment of iron. These stages are not all marked off from one another, for we find them together in the same country or

in different countries. Thus, the age in which at present we live is recognised as the Iron age, on account of the large employment of that metal; but we find that in various countries stone, more or less rudely fashioned, is still used in the manufacture of weapons or tools. For example, when I was in the Colonial Exhibition lately with Mr. Norman Lockyer, he pointed out a kind of threshing implement, such as is now used in Cyprus. It consists of a flat board, in the under side of which are imbedded a number of stone celts exactly like those made by prehistoric man, and perhaps used by him for a similar purpose as well as for axes. In the same way that we recognise four stages in the development of the implements used by man in the arts or in warfare, we may, I think, recognise four stages in the development of the implements he has used in the treatment of disease. In the first stage, crude drugs were employed, prepared in the roughest manner, such as powdered cinchona or metallic antimony. In the next stage, these were converted into more active and more manageable forms, such as extracts or solutions, watery or alcoholic. In the third stage, the pure active principles, separated from the crude drugs, were employed, e.g., morphine and quinine. In the fourth stage, instead of attempting to extract our medicines from the natural products in which they are contained, we seek to make for ourselves such substances as shall possess the particular action we desire. Now, just as we find stone and iron implements occasionally used together in the same country, so we find that drugs belonging to the different stages mentioned, are used at the same time. For example, we may find crude powders, alcoholic extracts, and pure alkaloids all contained in the same pill. Nay more, we may sometimes give to the patient in addition to all these, a medicine made artificially. But while this condition still exists, we notice that crude drugs are being less and less used, and their place is gradually being taken by pure active principles. We may say, then, that we are passing at present from the Stone age into the Bronze age of Pharmacology; and may, indeed, be said to be just entering on the Iron age. This age may be said to have begun about twenty years ago, when the researches which my predecessor in this office, Dr. Fraser, made with Professor Crum Brown upon the connection between physiological action and chemical constitution, inaugurated a new era in Pharmacology. They found that, by modifying the chemical constitution of strychnine, they could also alter its physiological action, and convert it from a poison which would tetanise the spinal cord, into one which would paralyse the motor nerves.

We might perhaps date the beginning of this age from Blake's attempts to show that a connection exists between the form in which various bodies crystallise, and the mode in which they act upon an animal body. Richardson, too, had observed that, amongst various compounds of carbon, certain differences existed in physiological action which might be supposed to correspond to differences in their chemical composition. And at the same time that Crum Brown and Fraser were making their experiments, Schroff in Vienna and Jolyet and Cahours in France had independently arrived at somewhat similar conclusions; nevertheless, I think we may fairly say that it was the experiments of Crum Brown and Fraser which fairly started pharmacology in the new direction in which it has since been steadily advancing. It would be impossible for me to enter at all fully into the recent development of this branch of research, but I think it may be both interesting and useful to try to give you a short and popular account of the chief points already made out; and, in doing so, I may perhaps be excused for using, almost to the extent of abusing, similes which are not precisely exact, but which may be useful in giving you a rough idea of a somewhat complicated subject.

We have all heard of the "flesh-pots" of Egypt; but I find that everybody is not acquainted with the "flesh-pots of Shiloh," though "good little Samuel" has, probably been frequently held up before us as an example to be followed, and possibly the naughty sons of Eli as an example to be avoided. When these sons of Eli were priests in Shiloh, their custom was, when any man offered a sacrifice, to send their servants with a "flesh-hook" of three teeth, in his hand, which he struck into the pan, or kettle, or cauldron, or pot; and all that the flesh-hooks brought up, the priest took for himself.

It is obvious that what the priest's man brought up would depend very greatly on two things, viz., the contents of the pot and the nature of the hook—whether it were large or small, sharp or pointed, single-pronged or many-pronged. It is obvious, too, that a very slight alteration of the points, by the judicious application of a file or whetstone, might considerably influence the savouriness of the priest's dinner. With the small pots that they were likely to have in Shiloh, it would not matter much what the nature of the handle was; but it would matter very greatly if the priests had to go fishing in the brazen sea of Solomon, for there, with a short handle, they might not be able to reach the tit-bits in the middle, and if the handle were too long, they