these two diseases were very severe and fatal amongst a certain portion of the population.

Acute articular rheumatism is rare, and no cases of malaria are met with, as a rule, though, in days gone by, the neighbourhood of Lancing gave us cases of ague in its various forms.

The Benefit of the Place for Invalids.—There is no doubt in my own mind that the climate of Worthing is most fitted for those people somewhat advanced in years, who desire rest and an easy means of getting about. For children, too, of all ages up to puberty, the place is very well suited. Its free and open exposure to the rays of the sun; the soft balmy air from the sea, accompanied with a large and frequent supply of ozone; and the evaporation when the tide is out, thereby cooling the heated air of the summer, and tending to bring about a fresh breeze always some time throughout the day—all these are favourable for its position as a health-resort. Its perfectly level situation tends to recommend it to all those suffering from heart-disease; and the atmosphere appears to agree with most cases of chronic bronchitis and some cases of asthma. In the latter stage of phthisis, the atmosphere has a beneficial and soothing tendency; but, in the earlier stages, the climate, in my opinion, is too moist and relaxing. Cases, too, of whooping-cough do well here, especially if brought down in the later stage. Chronic rheumatism, and those suffering from chronic rheumatic arthritis, derive benefit as well by being able to be fairly exposed to the influences of the sea-air, and enabled thereby to take passive exercise in this free and open sunshine.

ILLUSTRATIONS OF EXCEPTIONAL SYMPTOMS AND EXAMPLES OF RARE FORMS OF DISEASE.

By JONATHAN HUTCHINSON, F.R.S., LL.D.,
Emeritus Professor of Surgery at the London Hospital.

(Continued from page 1212 of last volume.)

XIV. RECOVERY FROM ASCITES, AFTER PARACENTESIS, IN A YOUNG MAN, THE SUBJECT OF DISEASE OF THE LIVER: GOOD HEALTH SEVENTEEN YEARS AFTERWARDS.

I SAW, with Mr. Alfred Kennedy, at West Ham, in May, 1884, a case of very considerable interest in reference to prognosis. Our patient was a man, named G., now aged 35. The reason for my seeing him was that he had been recently injured in a railway accident, but the interest of his case belongs wholly to his former history. Seventeen years ago, when he was 18, he had a long illness which, he said, was diagnosed as "ascites, pleurisy, and diseased heart and liver." He was treated in St. Thomas's Hospital, under the care of Dr. Bristowe. Some scars under the umbilicus confirmed his statement that he was tapped more than once for ascites. The left side of his chest was decidedly contracted. Both legs were covered by dusky stains, as if he had formerly suffered from thrombotic purpura. His heart's action was very irregular. I could not detect any *bruit*, and there was no proof of hypertrophy, but the action was peculiar and jerky, possibly from adherent pericardium. I was told that, on some occasions, a murmur had been detected. His liver was still very much enlarged, its edge, thickened and bossy, could be very easily felt two inches below the ribs; it appeared to be equally enlarged in all parts. The man told us that he had enjoyed excellent health, in spite of the conditions mentioned, ever since his discharge from St. Thomas's Hospital. He had never had any return of the ascites or oedema of the legs. He had led an active life, and had only suffered from a little shortness of breath and palpitation, after making unusual exertion. He had been married for six years; he looked fairly healthy, but not robust. Since the railway accident, he had suffered from palpitation, with a tendency to faintness and to hysteria. There did not appear to be any definite explanation of his former illness, for he alleged that he had always led a temperate life.

Having forwarded the above narrative to my friend Dr. Bristowe, he was so very kind as to refer to the hospital records, write out the following notes, and permit their publication.

Dr. Bristowe's Notes.—"G. G., aged 20, came under my care in St. Thomas's Hospital, November 11th, 1869. The history he gave was to the effect that, two years previously, he had been laid up for three months in University College Hospital with pleurisy, from which he had recovered; that, three months later (eighteen months ago), he began to complain of swelling and tenderness in the epigastric region, and that, from that time, he had been gradually enlarging in the abdomen. He continued at work up to three weeks ago.

"On admission, he was pale and thin. The abdomen was large, containing so much fluid that only a very small area of resonance existed in front. There was no tenderness, and no definite enlargement of liver or tumour to be detected. The left side of the chest was much contracted, measuring, at the nipple-line, only 16½ inches in circumference (the right measuring 19). It was dull in its lower half, and the breath-sounds were feeble or absent; no cough; no signs of progressive disease; urine free from albumen; legs oedematous. In other respects he was healthy. The abdomen slowly increased in circumference (at level of navel) from 38 inches to 41, and, on January 6th, two months after admission, twenty-three pints of fluid were removed by paracentesis. He was relieved, but slowly refilled, until, on March 7th, twenty pints were removed. After this, he appeared to be filling again slowly, and he left the hospital on April 11th, still presenting some ascites; but his general health seemed good.

"I have no record of his occupation, or of his habits, or of any statement with respect to syphilis; and I never detected clear evidence of disease of the liver. I have not recorded my diagnosis; but I suspect that I attributed his affections, first, effusion into the pleura, and then ascites, to tuberculosis. During the greater part of his residence in hospital, he was treated with quinine and iron. There was no disease of the heart."

XV. SIMULATION OF BRACHIOPLEGIA BY RHEUMATISM.

When the patient to whose case the following statements refer entered my room, with his head bent forwards and his arms held stiffly by his sides, I certainly thought, as he did, that it was a case of incomplete paralysis of the upper extremities, probably consequent on an injury to the cervical spine; and it will be seen that the history which he gave was one which quite fitted with such a diagnosis. He said that his arms were powerless immediately after a fall from a swing, and that they had remained so ever since. On careful examination, however, I came to the conclusion that nothing had happened, excepting a severe contusion of the neck and shoulders, causing pain in motion, and that the enforced rest, subsequently, had induced rheumatic stiffening of both shoulder-joints. On careful inquiry, it appeared that at no time had there been any sensory paralysis. There had never been any numbness in the limbs, or "pins and needles." Although all the muscles were wasted and thin, none were especially so, excepting the deltoids. He could move all the joints of his forearms, hands, and fingers quite well. The deltoids were so wasted, that all the bony prominences were easily seen. Both shoulders were stiff, permitting only the very slightest degree of motion; in both, there had been much aching, especially about the insertion of the deltoid, as usual in cases of rheumatic disease of the joint.

I inquired whether there was any tendency to rheumatic gout in his family, and found that his father had had gout, and that he himself had suffered from lumbago. From the manner in which he held his head stiffly forwards, and from his complaint of pain in the back of the neck and shoulders, it seemed probable that rheumatic changes were in progress in the joints of the vertebrae, tending to the production of spondylitis deformans. It is possible that he may have had some fracture of the laminae; but, whether this occurred or not, it is clear that there were never any real indications of injury to nerves. After the fall, he had much pain in his neck, and could not use his arms. He could walk, and was able to ride home, and walked upstairs, holding his arms across his chest and his head bent forwards. He was nearly two months in bed after the accident, with pain in the neck and arms. During that time, he lay almost constantly on his back in one position.

XVI. NEURALGIA OF THE SCALE, PROBABLY OF GOUTY ORIGIN: DETAILED DESCRIPTION OF THE PAIN.

A gentleman, possessed of some anatomical knowledge, and almost hypochondriacal in his tendency to self-observation, described to me the following symptoms. Early in the morning of March 2nd, he experienced over the middle of his left parietal bone a sharp explosion of lightning-like pain. It began exactly in the middle, and passed forwards nearly to the forehead, and backwards nearly to the occiput, being most intense in the centre. It radiated a little laterally, but not nearly so far as it went backwards and forwards, and it never crossed

¹ The notes, from which I quote, are in my own handwriting, and I believe that the absence of reference to these points means that I elicited nothing worthy of record, or that seemed to throw light on the nature of his case. My notes are difficult to read. On looking through them again, I find that, after each tapping, I examined the abdomen; that, after the first, I did not recognise hepatic enlargement; but that, after the second, I remarked that there was dullness for two inches below the right ribs, and that, in the scrobiculus, there was a distinct hard edge feeling like that of liver, but moving with the abdominal walls.—J. S. B.

the middle line. It did not run in straight lines, but spread out somewhat like a coruscation of sheet lightning. It was over in a few seconds, but quivered a little, not being absolutely momentary. After one explosion, another, and often a third or fourth, would follow in quick succession, and then he would be quite free for half an hour or an hour. The scalp was not in the least tender. The pain was quite bearable; in fact, at first it rather amused than hurt him; but, as the day went on, it became worse, and the explosions were attended by a feeling as if the left half of the head were being suddenly crushed. In the evening, the explosions were, both more frequent and more prolonged. On some occasions, there was the suggestion of a pain passing into the cheek and lower jaw, involving, in fact, the whole half of the head; but, in these parts, its course could not be traced with accuracy. Neither in the upper nor in the lower jaw did it run along the course of known nerve-trunks. The bicuspid tooth of the lower jaw was aching a little all day; but it was a steady ache, attended with a feeling of general numb swelling about it, and with tenderness on pressure, which prevented eating, but without any explosions of severe pain. My friend became very nervous in the evening, feeling sure that he was about to be the victim of "tic-douloureux."

I found that, on the previous day, he drank a bottle of claret instead of his usual modicum of a third; that he had eaten much more liberally of meat than was his wont; and that, on the day on which his neuralgia began, he also felt his ankles and wrists puffy and aching; and, finally, that his urine began on that day to deposit uric acid freely. Knowing that he had for long been threatened with gout, I assured him that the neuralgia was gouty, and made him take a full dose of bicarbonate of potash at bedtime. On the next morning, both the toothache and the sheet-lightning pains in the head had ceased, and nothing remained but a very slight perception of general ache in that side of the head.

Although it is easy, in this instance, to feel confident as to the influences which were at work, and to say that the neuralgic shoots, and probably the tooth-ache also, were caused by a gouty state of the blood (lithæmia), yet it is more difficult to explain the local mechanism of the pain. The scalp-pain always began in one and the same spot, and that not one to which any anatomical peculiarities attach. Was it a periosteal pain, and due to changes (neuritis?) in some ganglia, or other peripheral nerve-organ? Such seems the most probable hypothesis.

I could sympathise much with my patient, having often experienced pain of a somewhat similar kind. From my own observations, I can quite confirm the accuracy of his distinction between pains which are like forked lightning, and those to which the comparison of sheet-lightning may be applied. The latter are less severe, as a rule, and always less momentary than the others. They spread over a wider area, and do not feel to run in such straight lines. Both of them differ from another kind of explosive pain, in which the shoots extend for equal distances around the centre of starting. This might be called the star-shaped pain, or bomb-shell explosion; but, if the latter term be used, it must clearly be understood that the pain denoted, although always radiating, is by no means always of any terrible degree of severity.

THE AIMS AND CLAIMS OF THE BRITISH MEDICAL TEMPERANCE ASSOCIATION.

An Abstract of a Paper read at the Tenth Annual Meeting of the Association.

By J. J. RIDGE, M.D., B.S., B.A.Lond., Honorary Secretary.

AFTER reference to the great increase in the number of medical abstainers since the formation of the Association, many more being still unenrolled than the 341 who have already joined, Dr. Ridge proceeded:

"Our first aim is to form a rallying point for medical abstainers through the length and breadth of the land. We are constantly discovering in busy cities, or in country towns and villages, medical men who are total abstainers, but whose fame as such is not known far beyond the neighbourhood in which they dwell. It is gratifying to all who are interested in the temperance cause to find so many of these solitary adherents, and a source of help and encouragement to them to learn that they are not alone, but are associated with an ever increasing number of their medical brethren. The Association makes such men acquainted with each other, and mutual sympathy strengthens individual zeal.

The object of the Association is thus very clearly stated in the

constitution: "to advance the practice of total abstinence in and through the medical profession, and to promote investigation as to the action of alcohol in health and disease." We aim to stimulate the interest of medical practitioners in the subject of total abstinence. The future of the temperance movement depends on the adoption of right views of the position and action of alcohol on the human organism, both mind and body. What those views should be, ought to be decided by medical scientists. The effect of alcohol on body and mind is part of that physiology which is the "proper study" of the profession. The world waits for a clear and authoritative declaration. There has been too often given it hitherto the "uncertain sound" of contradictory statements, a display as confusing, and often amusing, to the public, as it is injurious and discreditable to us. Practical men have adopted total abstinence, temporarily or altogether, to enable them to accomplish feats of strength, skill, or endurance. Soldiers, engineers, labour-masters have enforced total abstinence on occasion for the purpose of attaining most easily the highest efficiency of their men, and have been delighted with the result. If the profession, or any members of it, set their faces against a principle thus practically tested and triumphant, it is high time that they realised that they are running their heads against a rock, and must suffer the inevitable consequences. It is necessary to declare that the highest development of health and strength, the soundest condition of mind and body, are only possible when intoxicating liquors are abstained from. The Association enables the profession to declare this in the best and most natural way, by showing that each member practices what he preaches.

For, after all, and very properly, the public will judge us more by what we do than by what we say. Our special knowledge involves a special duty. The medical abstainer is a living beacon, whose silent yet eloquent example influences many far beyond his ken, saving them from the alcoholic rocks and quicksands, on which some would otherwise be inevitably shipwrecked.

We advocate total abstinence because we are in earnest, and practical reformers, dealing with human nature as we find it. Our ultimate aim is the extinction of all the special evils which the action of alcohol is able to produce, and the diminution of all other vices, diseases, and troubles by how much they are now aggravated by alcohol. The method which commends itself to our common sense is ourselves to discontinue to drink alcoholic liquors, and to induce all whom we can influence likewise to abstain. Our end is attained in proportion as our means are adopted. A medical abstainer has more influence than any: "Truth, from his lips, prevails with double sway."

I need hardly remind medical men that the harm arising from the use of alcohol is not to be measured by the unsteadiness of the legs or the thickness of the speech. We are all aware that a great deal of perversion of function takes place before these occur. We know that it affects the brain, upon which it exerts a paralysing action, weakening first of all the highest powers of the mind, those which are least automatic, those which are most connected with the individual, or are the latest acquisitions of the race. It produces progressive paralysis of the judgment on the sensory side, and of the will on the motor. It lengthens the time required for a voluntary effort; and, while reducing self-control (that is, temperance), leaves the lower reflex centres to act for a time more automatically; these reflex actions are, with larger doses, one by one weakened and extinguished—the most complex first, until at last only respiration and circulation remain, or may themselves be narcotised.

I am convinced that all will admit that this gradual paralysis of mental and moral control is of more importance than any paralysis of mere muscular power or co-ordination. Hence it is not desirable to practice any habit which can produce such results. This is a clear, safe, and easily understood practice, such as the profession should recommend for the moral elevation and physical development of the masses.

Some contend that medical men have no special concern with morals; but we are still men, bound to promote the good of others, and surely unable to repudiate our obligation, because we have special knowledge and greater influence. The fact that morality is affected by a material agent—alcohol—makes it peculiarly the province of the physician. We ought to let our fellow-creatures know that alcohol saps the foundation of morality, by rendering self-control more difficult, and at last impossible. Alcohol degrades a man, or a nation. It takes the individual back through the stages of youth and childhood, to the unconscious and merely animal condition of the new-born babe; and it has a greater effect on the young, and on those low in the scale of intelligence and general self-control. The same applies to nations; its power being more irresistible and destructive, the lower a nation is in the scale of civilisation. Physicians, then, ought to exert their