may be treated successfully without a great deal of attention, there are others—and their number is large—in which no trouble must be spared to effect the desired cure. The adaptation of the apparatus may require frequent alteration; the position of the patient during recumbency, and even the management of every detail of his life, may have to be supervised, or at least directed, by the surgeon. We must not rest content with saying that some cases will recover and others will die. I would rather urge my belief that every case, if dealt with with sufficient energy, and if taken in hand before the constitution is undermined, may be cured provided no intercurrent disease be present to complicate the case.

Mr. JESSOP remarked upon one or two signs of carious spine, on which, in cases of difficulty and obscurity, he relied with some confidence. First came the power of the patient to twist his spine, as in looking behind him, whilst his feet were kept in apposition and motionless. Then there was the effect of pressure upon the head by the hands of the surgeon, a means of diagnosis which, if attended with some danger in advanced cases, was safely applicable in those which were obscure; again, there was the pain elicited by jumping from a chair or table. He mentioned these chiefly because Mr. Smith, in his remarks, seemed to condemn them, and he desired it to be understood that he applied them only in cases of obscurity, and therefore presumably of early disease.

EXAMINATION OF THE NASAL CAVITIES FROM THE FRONT.

Read in the Section of Otolary at the Annual Meeting of the British Medical Association in Brighton.

By E. CRESSWELL BABER, M.B. LOND.,

Surgeon to the Brighton and Sussex Throat and Ear Dispensary.

EXAMINATION of the nasal cavities from the front has long been a favourite method of ascertaining the condition of the interior of the nose. For two reasons, however, there is often, especially to the beginner, considerable difficulty in recognising the different structures presented to the eye. They are—first, the perspective view of many of the parts which alone can be obtained; and, secondly, the variations in appearance which these parts present in the same individual at short intervals of time. In order to facilitate the examination, and to render a record of its results more easy, I have made some sketches of the views obtained by this method, which are, as far as I know, the first representation of these parts as seen in the living subject.

The detailed study of the subject necessary to produce the drawings has brought out some points which I think may be of interest to the Section, although it is, of course, impossible to enter fully into the subject in the limited time now at my disposal.

The extent to which the deeper parts of the nasal cavity are visible from the front depends largely upon the state of the erectile body on the anterior extremity of the inferior turbinated bone; in other

words, upon whether this structure is in a collapsed or in a swollen condition.

1. *View when the Anterior Erectile Body on the Inferior Turbinated Bone is collapsed.*—When the inferior turbinated body is collapsed (Fig. 1), its anterior extremity (*a*) forms a somewhat irregular undulating surface, terminating below in a more or less sharp margin, which overhangs the inferior meatus. Above, it tapers gradually into the outer wall of the nasal cavity, forming, as viewed from the front, a kind of “neck” (*l*). Its inner surface is seen greatly fore-shortened, and appears either as a shallow concavity or as a slightly rounded eminence (Fig. 1*b*). When examined with a probe, the bone can be felt under the mucous membrane, and the instrument does not sink in to any depth.

Middle Turbinated Body.—Above, and at a considerably greater distance from the nasal orifice, is seen the middle turbinated body (Figs. 1 and 2).

A consideration of the anatomy of the nose will show that, provided there be nothing to intercept the view, the parts of the middle spongy bone visible will be the anterior and inferior borders, the angle of junction of the two being the part which is nearest to the observer's eye; each of these borders presents an inner and an outer lip. Moreover, as the surgeon's eye is usually closer to the septum than to the outer wall of the nasal fossa, a portion of the inner surface of the bone will also be visible.

Theoretically, these are the parts which should be seen, but, practically, a considerable portion of the middle turbinated body is hidden by the tubercle of the septum (Fig. 1, *d*, Fig. 2, *A e*). The tubercle of the septum, which was first described by Morgagni, and has been recently figured by Zuckerkandl, consists of a thickening of the mucous membrane, containing glandular elements, at the front entrance to the olfactory slit. The view obtained, therefore, depends to a great extent, upon the shape and size of this structure. The three diagrams in Fig. 2 have been drawn to show the different appearances which are met with.

When the tubercle is absent, or so small as not to interfere with the view (Fig. 2, Diagram C), the anterior margin of the body is seen terminating above in the “neck” of the bone, that is, the narrow por-

Fig. 2.—Diagrams illustrating different appearances presented by the tubercle of the septum. (Right anterior rhinoscopic view.) *a*. Outer lip of anterior border of middle turbinated body. *b*. Angle of middle turbinated body. *c*. Lower border of middle turbinated body. *d*. Inferior turbinated body. *e*. Tubercle of septum. *f*. Outer wall of nasal cavity. In diagrams A and B, the margins of the middle turbinated body, hidden by the tubercle, are indicated by dotted lines.

tion where it joins the body of the ethmoid; below this, it runs downwards and, usually, outwards, increasing in breadth as it descends. Beneath the angle of the body, and rapidly receding into the distance, and, therefore, less highly illuminated is the inferior border. The outer lip of the anterior border is separated from the outer wall of the nasal cavity by a deep groove of greater or less width (Fig. 1, *k*). This groove, into which a probe can be inserted, is of importance, as it leads into the hiatus semilunaris. A little of the internal, or median, surface can sometimes be seen. Between it and the septum is the olfactory slit. The whole body, when viewed from the front, often presents a sigmoid form, being convex above, and concave below on its inner or septal aspect. When the tubercle is small, we have the appearance shown in Fig. 1. The inner lip of the anterior margin of the middle turbinated body is, to a great extent, hidden, but above the tubercle is seen the “neck” of the body. A more prominent tubercle will hide the whole inner lip of the middle turbinated body, but a part of the neck remains visible (Fig. 2, B).

Diagram A shows the parts as they very commonly appear. The tubercle, which is large, and extends to the top of the field of vision, conceals the neck and the whole of the inner lip of the anterior border of the body, leaving only a small portion of the outer lip of the an-

Fig. 1.—Left anterior rhinoscopic view in a Woman, aged 26. *a*. Anterior extremity (collapsed) of inferior turbinated body. *b*. Inner surface of same, seen in perspective, showing a rounded swelling. *c*. Anterior border of middle turbinated body, ending below in the angle, which in this case presents a puffy swelling. Above, the anterior border terminates in the neck of the body (*e*). *d*. Tubercle of the septum. *e*. Lower border of middle turbinated body. *g k*. Depressions on either side of the neck; *g* forming part of the olfactory slit, and *k* a portion of the middle meatus. *h*. A moderately deep-seated swelling on the septum. *i*. Ditto, much further inwards. *j*. Inferior meatus, through which the movements of the palatal muscles can be seen. *l*. “Neck” of the inferior turbinated body.

terior border (*a*), the angle (*b*), and the lower border of the body (*c*), exposed to sight. The outer lip of the anterior border is sometimes hidden by a swelling of the outer wall of the nasal cavity at *f*, which swelling may be temporary in character. The view may also be rendered more difficult to understand by puffiness of the mucous membrane of the middle turbinated body in this situation, partly hiding the neck of the body.

All these appearances are met with in practice, but, in my experience, it is more rare to find the tubercle flat or absent, than to see it hiding a considerable portion of the middle concha.

Not having found any references to the tubercle of the septum in the living subject, I was for some time at a loss to account for this swelling, so constantly present on the septum. On studying Zuckerkandl's description of the tubercle in the dead body, however, I recognised that it was the tubercle that I really had before me.

The tubercle must be distinguished from a deflection of the septum, which, in the living subject, it resembles in appearance. It is commonly seen on both sides in the same individual, and by inserting one blade of a Fränkel's speculum into each nostril, the two tubercles can be compared. At the same time, it is important to remember that a deflection of the septum in this situation, which is of common occurrence, may modify the aspect presented by the tubercles, making one appear more, the other less, prominent than usual. Like the inferior turbinated body, though in a much less degree, the tubercle appears capable of undergoing temporary swelling; at least it can be reduced by the application of cocaine. It is soft to the touch of the probe, by which it can, to a certain extent, be pushed aside. Touching it often gives rise to sneezing, under apparently normal circumstances. It may be fairly suggested, I think, that the tubercle guards and protects the anterior extremity of the olfactory slit, in the same manner that the anterior end of the inferior turbinated body, especially when erected, guards the respiratory region, the protecting agency of the tubercle being enhanced by its tendency on contact with a foreign body, to give rise to the violent expiratory act of sneezing.

Besides these usual appearances of the middle turbinated body, it will be readily understood that others occur which it is impossible to detail here.

The mucous membrane covering the middle turbinated body is paler than that on the inferior. It is normally smooth and shiny in appearance, and often presents a very minute red and yellow mottling, which is best seen in reflected sunlight.

The Septum.—Having described the tubercle, I need say but little else regarding the septum. The frequency with which it presents irregularities and deflections renders it necessary to estimate its straightness by a comparison of the appearance on the two sides. It is very usual to meet with a small conical eminence opposite the anterior inferior end of the lower spongy bone (see Fig. 2). A ridge is sometimes seen running upwards and backwards from this, and corresponding to the anterior margin of the vomer. Other prominences occur higher up and farther back on the septum (Fig. 1, *h*, *i*). When the septum is much deflected, not only the lower, but also the middle turbinated body in the larger nasal cavity, often exceed their fellow in size.

By projecting a strong light through the inferior meatus (Fig. 1, *j*), or through the space between the inferior turbinated body and the septum, the movements of the palate during deglutition or phonation can be very commonly seen. When the palate is at rest, the posterior wall of the naso-pharynx is visible. It presents a nodulated, hillocky aspect, often with several light spots, and, being seen *en face*, is more highly illuminated than the deeper parts of the septum and inferior turbinated body. In fairly wide nasal cavities, the upper margin of the choana, with its two arches, one immediately above the other, is sometimes seen, also the swelling on the septum close to its posterior margin, which is so often observed by posterior rhinoscopy.

2. View when the Anterior Erectile Body is Swollen.—We have hitherto considered the appearance presented when the anterior erectile body on the inferior turbinated bone is collapsed. We must now turn to the view when this structure is swollen.

Fig. 3 is taken from a man, aged 24, in whom the erectile body was seen to swell while he was under examination. Twenty minutes previously, the inferior turbinated body presented an appearance very similar to that shown in Fig. 2, Diagram A. The middle turbinated body was plainly seen, also the palatal movements. In its swollen state, on the contrary, you see the inferior turbinated body practically occludes the inferior meatus, preventing all view of the naso-pharynx, and allowing only a glimpse of the middle concha. When more swollen still, as often happens, the inferior turbinated body hides the latter completely, being in close contact with the septum. The inferior

turbinated body itself appears as a round shining eminence, with a smooth surface; above, it is prolonged into a broad neck (*b*). This neck is formed by erection of the mucous membrane above the inferior turbinated bone, rather than of that on the bone itself, and is of great practical importance, on account of the way in which it hides the upper part of the nasal cavity. The sensation produced by touching the swollen inferior turbinated body with a probe is that of a soft cushion, which can be deeply indented by pressure, but rapidly regains its original globular shape. This characteristic serves to distinguish it from true hypertrophy of the turbinated body. Its colour does not necessarily differ from that of the non-erected body, but it may be either paler, or more florid in appearance.

Fig. 3.—Right Anterior Rhinoscopic View.—*a*, erected inferior turbinated body terminating above in the swollen "neck," *b*, which almost entirely conceals the middle turbinated body. The upper part of the drawing is slightly diagrammatic.

The view of the deeper parts is, in my experience, much more commonly interfered with by swelling of the anterior erectile body than by malformation of the septum or inferior turbinated bone. If necessary, however, the swollen tissues can be pressed aside with a probe, blunt hook, or other instrument, and a limited view of the deeper parts obtained, or, better still, the erection may be reduced by cocaine. Cocaine, as you know, possesses the property of contracting the swollen erectile tissue in the nose, and is therefore valuable for distinguishing between erection and hypertrophy of the inferior turbinated body, and for rendering the nasal cavity clearer both for purposes of respiration and for facilitating anterior rhinoscopy. Its action appears, however, to vary in degree in different cases.

Except when it is extreme, erection of the anterior part of the inferior turbinated body alone does not produce marked nasal obstruction. I cannot here enumerate the different causes which give rise to this temporary erection, but in its slighter stages it is no doubt a physiological phenomenon, the entrance of irritating particles into the nose being thereby prevented.

A few words in regard to the use of the nasal probe. This instrument is, of course, only to be employed under careful guidance of the eye. Besides giving information about the shape, attachments and consistence of growths, especially of small swellings on the turbinated bodies, it is indispensable for ascertaining the reflex irritability of the different intra-nasal structures. By stroking gently in turn the inferior and middle turbinated bodies and septum, and by observing whether any reflex phenomena—such as sneezing, watery secretion, erection of the inferior turbinated body, or cough—be produced, important information may be gained. Touching the anterior extremity of the inferior turbinated body produces, in my experience, no reflex phenomena as a rule; exceptionally it gives rise to cough. The effect of touching the tubercle of the septum has already been mentioned.

In conclusion, I would mention my conviction that in recording the appearances presented by anterior rhinoscopy, a rough sketch of the view obtained, accompanied by a few notes, is far more effective than many lines of description, and sketching these parts is facilitated by the use of the nasal outlines which have been prepared at my suggestion by Messrs. Danielsson.

It is, of course, only possible to speak from one's own experience, but I have found graphic representation of so much practical value, that since making the original drawings in 1884, I find I have drawn between three and four hundred anterior rhinoscopic sketches in the notes of cases under my care.

BEQUESTS AND DONATIONS.—Mr. William Hayes, of Clarence Place, Cork, bequeathed £300 to St. Patrick's Hospital, and £50 to the North Infirmary, both at Cork.—Mr. Charles Thomas Wakefield Parry, J.P., bequeathed £100 to the Chester Infirmary.—The Leicester Infirmary has received £100 under the will of Mr. W. Bowman.