day, the same experiment with asparagus was tried on Eng, with similar results. They have not passed entirely through life without a quarrel. Their mother spoke of one occurring when they were eight years of age; and again, while on their first voyage from Siam to America, one of them wished to bathe, as was their custom, to which the other objected, as the day was cold. The difference was soon adjusted; those rare exceptions proving the rule rather than the reverse.

THE ETIOLOGY OF PHTHISIS.*

By PETER EADE, M.D.Lond., M.R.C.P.,
Physician to the Norfolk and Norwich Hospital, etc.

THE subject of consumption and of tubercular disease in general has always attracted a very large amount of attention on the part of the profession; and this may well be the case, seeing how large a proportion of the human family is cut off by this intractable disease. Of late, the interest attaching to it has received a further sharp, and somewhat sudden stimulus from the views which have been advanced respecting it by Dr. W. Budd on the one hand, and on the other from the published experiments of Villemin, Simon, Clark, Burdon Sanderson, Wilson Fox, and others, showing that the disease, or a condition undistinguishable from it, is capable of being produced in some animals by the inoculation of tubercular or other matters, or in rodents merely by the irritative-inflammation set up by the presence of setons beneath the skin. The addition to our knowledge which we have thus gained is immense; and I think it is not going too far to say, of the minute anatomical investigations of Dr. Wilson Fox in particular, that they have thrown a bright stream of light upon this part of the subject.

But still, even if we could assume that the intrinsic element of tubercle and tubercular disease is an excessive development of lymph-cells and lymphatic products, and could take it for proved—which, however, is not proved—that this process of abnormal development may be set up in man as in rodents by the introduction into the system by inoculation of tubercular or allied matters, it is very clear, as is fully stated by Dr. Fox himself, and endorsed by Drs. Cotton, Wilks, Marston, and others, that the demonstration of these facts is not the whole of the case, and that, in a large majority of instances of tuberculosis, some other etiological element or condition is at work to cause or bring about the fully formed disease; for not only do we find that individuals develop phthisis or other forms of tuberculosis where no suspicion of inoculation or contagion can exist, and where ordinary causes have evidently produced the disease, but also that infants are born into the world with the tubercular diathesis so strongly stamped upon them from the very first, that there can be no doubt that this condition is one which has grown with their growth and development, and is, in fact, a congenital, hereditary, and intrinsic part of their animal formation.

Tubercle, according to present evidence, would appear to consist essentially of cells usually more or less degenerating, and of nuclei; these cell-elements being generally formed in small rounded masses or aggregations, or, in many of these, coalesced to form larger agglomerations. Further, the recent researches by means of inoculation have proved almost to a certainty that these or similar cells or nuclei once formed, are capable, under certain suitable conditions, of multiplying and reproducing themselves—i.e., are endowed with a sort of independent vitality analogous to that seen in other forms of cell-life, and in this respect fairly comparable to the property of proliferation possessed by the cells of cancerous, recurrent fibroid, and other tumours. But the recent investigations of Waller, Cohnheim, and others, have shown that the cells of cancer, like those of all other animal tissue, are not abnormal developments or proliferations of other cells or corpuscles, and there is strong reason further to believe that they are in fact but vagaries of natural elements, unrestrained and unguided growths, as it were, of normal, glandular, epithelial, or connective tissue-cells or corpuscles.

In the *Lancet* for 1863, I called attention to the well-known fact that, in a very large number of instances, primary cancers follow closely upon some injury to the part from which the tumours spring; with a view to the expression of an opinion that the absence of controlling nervous force from the effused and organising lymph-elements was the main cause of their permanently stopping short at the lowest form of life—that of cells. I would venture, to-day, to give some reasons for thinking that a peculiar form of deficiency of nervous power may be an important element in the production of the superabundant cells which constitute

the various forms of tubercular disease, and that this disease may, in fact, be but an expression of a feeble vitality—feeble especially, and primarily, as regards the constitution of the nervous system, and secondarily, as regards the tissues which, from want of due and vigorous innervation, have been either imperfectly and loosely constructed from the first, or have become disordered in their action at a later period.

Cells more or less simple are, there is no doubt, the original and lowest form in which animal life is presented, and the type under which it approximates most closely to the vegetable world. This form of life, too, is the one which is capable of the quickest and easiest multiplication. Higher forms of animal bodies are formed out of such cells, variously modified, elaborated, and combined, to form tissues endowed with properties fitting them for the respective functions they are required to perform.

Every increase in the elaboration of the animal tissues is accompanied by a proportionate increase in the quantity and complexity of the nervous system; and it is most noteworthy, in reference to my present hypothesis, that the development of the nervous centres (especially as seen in the highest animal, man) actually precedes the formation of any other part of the body. There is strong reason, therefore, to believe that those centres exercise a governing and controlling power over the development of the other tissues, all of which are of later formation than themselves. Any original defect, therefore, of this primal nervous governing centre, will permit the body to be less perfectly formed, or, in other words, will tend to allow its tissues to be created on a degenerated type, with a constant tendency to revert more or less to the elements of which they are composed—i.e., cells. A body so created, and gradually growing and enlarging under such defective influences, is exactly the one which would eventuate in the soft, flabby, feebly constituted animal framework which we see in the typical rickety or scrofulous subject, the chief characteristics of whose structures are their general weakness and want of vigour, or rather power of endurance, and their proneness to the development of cells, as shown by ulceration—i.e., a ready disintegration or separation of the cell-elements of the feebly formed tissues; by glandular enlargement—i.e., distension of the lymphatic tubes by the hyperdevelopment of lymph-cells and gland-cells; and by purulent discharges—i.e., the excessive development of cells of pus from the mucous membranes.

This, then, would constitute an explanation of the congenital form of the tuberculous or strumous diathesis—a cachexia in which the body throughout is organised with a less perfect type of tissue, and with its nervous centres unequal to their proper work, either from original defect of formation, or from their being so feebly constituted as to be unable to appropriate to themselves the requisite nerve-force-developing pabulum. In the worst cases of strumous cachexia, the feeble nervous power secreted and available is insufficient from the first for the ordinary and daily demands of the parts which it should supply, and such subjects do rarely reach adult life. In somewhat better types, the nerve-power is equal to this work, but it is immediately used up by the slight extra demand made upon it by organs or tissues, with the result that some form of strumous disease immediately ensues, as is well illustrated in the proneness of the most worked parts of the developing body to degenerate first—i.e., to develop cellular products.

But tubercular disease is not only hereditary and congenital; it may be also acquired, even by the strongest and most robust, when they are placed in suitable circumstances. But it is noticeable here, that these vigorously formed subjects rarely develop other ordinary forms of strumous disease, though they do frequently become the victims of pulmonary phthisis.

I have said that the necessary effect of partial removal of nerve-force is to allow the parts supplied by it to take on a lower form of vital action; and the degeneration will, of course, vary with the cause of the deprival, with its intensity and seat, as well as with the constitution of the individual. Ordinary complete removal of motor or sensitive nerve-influence of course produces ordinary paralysis; but removal of nutritive nerve-influence produces other and distinct effects, as shown by the experiments of Reid and others; whilst the diminution of the combined spinal and nutritive nerve-influence gives rise to what is called suppurative or destructive inflammation—i.e., a process in which destruction pure and simple does not take place, but in which a rapid form of a lower type of growth or development—viz., that of cells, of pus-cells—is allowed to take place, at the expense of the more highly formed tissues. This effect is familiarly seen in the so-called catarrhal inflammation of the bladder which ensues upon spinal disease or injury, and in the suppurative disease which, as has been pointed out by Drs. G. Budd, Sims, and others, takes place in a lung whose nerves have been invaded or compressed by a cancerous tumour in the chest.

The aspect of the subjects of congenital and hereditary tuberculosis appears to me to favour this nerve-view of the disease; for such individ-

* Read in the Medical Section at the Annual Meeting of the British Medical Association in Oxford, August 1863.

duals are generally softly and delicately formed, confessedly weak in resisting power, and prone to have parts or organs take on tuberculous action whenever these are called upon for any large amount of exercise or exertion. This is well illustrated on a large scale by the fact pointed out by Dr. Barlow, and now well recognised, that the three great cavities of the body are most liable to take on this diseased action in the order of time in which they are respectively called upon for active exertion during the process of growth—viz., the head in infancy, the belly in early youth, and the lungs in later periods of life: the irregular and excessive development of the nervous centres being further shown by the frequently disproportionate strength of the mental powers in infancy, and their sudden failure under the exertion which they crave, and for which, indeed, they are temporarily fitted.

I have already alluded to the proneness of these individuals to ulcerations, to suppurations, and to diseases in which cell-growth, and that of an imperfect kind, is the prevailing type; and I would merely mention one other very familiar symptom of the tuberculous cachexia, viz., the tumid puffy upper lip—a condition which I believe to be very perfectly imitated, for the time being, by the prominent pouting of this part during a paroxysm of acute hysteria, in which it is well known that voluntary controlling nervous power is temporarily wanting.

Again, I think that the phenomena of the invasion of tubercular diseases in those originally strong and healthy agree with this view of its originating in depraved, or depressed, or irregular nervous power; and we must never forget that, whatever be the predisposing causes of disease, the nervous tissue, as the most highly and delicately organised, are nearly sure to be the first to suffer in nutrition, and consequently in integrity, and therefore are preeminently likely to be the starting-point of the disease. The predisposing causes are undoubtedly such as cause a depressed condition of system—such as mental anxiety; exhaustion from overwork or discipline; privation; poverty or neglect; intemperance; want of fresh air or exercise. It is during the operation of these that what is called the pretubercular state arises; and it is when to the depressed state of the general nervous power induced by them that local abstraction or exhaustion of nerve-force is superadded, as by over-exertion of the lungs, or the application of cold or wet to the skin, or by the continued irritation of dust or rough particles inhaled into the lungs, that the local tubercular disease springs up—i. e., that erring local life, or local cell-growth of a defective character, is allowed to begin.

Again: it appears to me that the results of the treatment of this disease are all in favour of this view. I need only allude to the negative evidence afforded by the disastrous results of the treatment of phthisis by "coddling" and hot rooms, and to the far better results of the opposite mode of treatment by fresh air, exercise, and tonics with good food; but I would especially point to the fact that all the medicinal remedies of any acknowledged efficacy (with the single exception of cod-liver oil, whose modus operandi is unknown) belong to the tonic class, and nearly all to the mineral kingdom. It may be fanciful to suppose that their efficacy depends upon their furnishing fresh electrolytic food or the wanting positive or negative elements to the minute nerve-batteries or electro-voltaic arrangements aggregated in the nerve-cells of the great nerve-centres; but I cannot help thinking that, even if fanciful, it is very likely true.

Nerve-force and galvanic force are, as is well known, not identical, but closely allied forces. Again, heat and electricity are also nearly allied forces, and not only so correlated, as it is called, but actually convertible. It is almost certain, therefore, that heat and nerve-force are also both correlated and convertible terms; and we can thus understand the way in which not only wet, by rendering clothes better conductors, can abstract the nerve-force of the body as it would electricity, and so destroy its balance and make its working irregular, but also how the mere abstraction of heat from the body by the influence of currents of air can effect the same result.

Now, when a deprivation of nerve-force is effected on a large scale in a healthy, i. e., a non-cachectic or non-predisposed, individual by the application of cold or wet to the surface of the body, and especially of the chest, ordinary acute inflammation of the lungs is probably the result. The exhaustion of the nervous twigs, spread out upon the skin, being conveyed along the afferent nerve-trunks to the nerve-centres, these themselves suffer, and then the exhaustion is propagated by a further reflex action along the efferent nerve-trunks to the internal mucous surface supplied by these nerves, and, this being thus deprived of nerve-force, a perverted vital action takes place; the mucous membrane, as a result of this, secretes gelatinous or plastic albuminous matter, and the ordinary condition of pneumonia is then established. But if the individual, instead of being healthy, be lowered in vital health, or cachectic, the secretion, as is well known, instead of being mainly composed of healthy plastic matter, is constituted of material formed on a lower type. Or,

again, if the exhausting influence, instead of being exerted suddenly, is exercised partially or more gradually, then a different kind of action ensues. The nerve-centres are reflexly exhausted only partially or irregularly; their influence on the pulmonary membranes is cut off only in spots or localities; and then, where this takes place, the irregular vital action which results eventuates in the production or secretion of nodules or masses composed of imperfect cells and nuclei and other germinal lymph-matters, having the well-known tendency to degenerate and decay—in fact, of ordinary tubercles.

In cases of phthisis, originating in mechanical irritation of the lungs by flour, grinder's dust, etc., the exhaustion is probably of a somewhat different character. In this case, the peripheral extremities of the nerves of the pulmonary mucous membrane would be the parts primarily exhausted of their force, and the effect would be exerted either directly, or reflexly, or in both ways upon this surface, producing either partial con-olidations (mixed quasi-cellular and albuminous deposits), or, where true tubercle is found, the small punctate masses of this substance known as milary tubercle.

The necessity for making my paper as short as possible, and my unwillingness to take up the valuable time of the Association at any great length, have obliged me to omit many references to distinguished authors which I should otherwise have desired to make, and have only enabled me to bring before you the most meagre outline of the subject. Indeed, it is only the fact that this topic is now uppermost in the professional mind, and my conviction, which I have held increasingly for some time past, that the truth of the great problem of tuberculous lies somewhat in this direction, that has induced me to bring before you today my views upon the subject.

The facts of the almost universality of the disease, the construction of a large class of individuals on the tubercular type, the acknowledged imperfection of structure of even the best formed of our bodies, the proneness of that part of the body to become tubercular which is being most used at the time of its development, all point to the insufficiency of any theory of mere lymphatic disorder adequately to account for all its phenomena, and to the necessity of some more intimate and vital cause to explain these. On the other hand, the close relationship of inflammatory and tubercular disease of the lungs (a relationship acknowledged by Dr. Williams, in his late most masterly account of this disease, in the *Lancet*); the acknowledged origin of the former in cold and other exhausting agencies applied to the surface—in other words, in the abstraction of vital heat or nerve-force; and the adequacy of this view of reflex nervous exhaustion to explain so very many of the phenomena of consumptive disease—justify me, I think, in endeavouring to refer them both to such a cause. I have, perhaps, to apologise not only for the imperfection of my views and arguments, but also for venturing to handle at all a subject which has engaged the thoughts and pens of many able observers. But I feel that, in spite of all that has been done, and although we seem to be becoming well acquainted with the histological characters of pulmonary and other tubercle, yet its intimate causation and origin, its meaning or relation to the basement membrane or germinal tissue which produces it, are still quite undetermined.

Although some vague expressions of belief in the possible origin of phthisis in defect of the nervous system have been used by some authors, yet, I believe, no definite attempt has hitherto been made to show how this might take place. To point out a way in which I believe its morbid conditions may be accounted for, through its agency, is the utmost that I have aimed at to-day; and I can only hope that these suggestions of mine, however crude, may, at least, bear some fruit, by turning thought in a new direction, and may so contribute, in however small a degree, to the elucidation of a disease which has hitherto foiled all our efforts either to unravel its nature, to cure its lesions, or, I might almost say, even to mitigate its ravages.

I will, in conclusion, thus briefly summarise what I have said.—1. The congenital tubercular diathesis, or cachexia, is essentially due to original defect or disease of the nervous system.—2. It consists in a development of the tissues of the body on an imperfect or lower type, owing to the absence of controlling nervous influence of a standard sufficiently high and perfect for the wants of the growing body.—3. The actual appearance of strumous disease is due to the exhaustion of the nerve-power of the part in which the disease commences.—4. In acquired tuberculous, the cell-development is the result of the removal of the nerve-power of the part affected, usually by reflex exhaustion, propagated from the part upon which the morbid causative influence is exerted.—5. There is a very close relationship between the causes of inflammatory and tubercular disease of the lungs; to which I would add—6. The possible inoculability of tubercle matter in man as well as animals does not militate against this theory, since, if true, it only proves that tubercular, like other cells, when once formed, are capable of proliferation and reproduction.