APPENDIX.

After the foregoing paper had been read at Oxford, a second paper was read on the same subject, by Mr. Lockhart Clarke, and it gave me great pleasure to find that he took a view of the reasons of locomotor ataxy substantially like my own. Dr. Duchenne of Boulogne, who was present, differed from both of us and strongly urged his own original interpretation. Both publicly and privately Dr. Duchenne was most unwilling to allow that there was any loss of motor power in the disease; but of this I have no doubt, and I think that Mr. Clarke agrees with me. I have subsequently found that Oppolzer also enumerates "weakness of the lower extremities" among the early symptoms.* My own opinion on this point was formed independently, as I have repeatedly proved it during the last four years to the Leeds students, and have stated it in clinical lectures. In his paper, in the *St. George's Reports* for 1867, I see that Mr. Clarke does not distinctly lay down weakness of the lower extremities as a symptom of locomotor ataxy; and it is distinctly denied by Dr. Radcliffe.† I attach much importance, however, to this symptom, for, I believe, in opposition to many much better men than myself, that the disease of the posterior columns is chiefly the cause of this weakness, and that it has not much to do directly with the incoordination. Indirectly, it has much to do with it; and that, as I have said, by its encroachment upon the fibres of the posterior roots; for, herein, namely, in our interpretation of the way in which this encroachment causes incoordination, lies the difference between the view I venture still to hold and that of my friend Mr. Lockhart Clarke. My own view is simply this: that the loss of tactile sensibility is sufficient to cause loss of coordination, that the tactile sensibility is lost, and, therefore, that the loss of tactile sensibility is probably the cause of the parallel loss of coordination; that, in short the extravagant gait results from loss of consciousness of the relations in space between the body and the medium.‡ Lockhart Clarke's view differs importantly in detail from this, and his view seemed to me at first so suggestive and so ingenious withal, that, on the reading of his paper, I felt it only remained for me to pass on my own to the tallow-chandler or to the British Museum. On subsequent reflection, however, unfortunately for the reader, I have recanted, and, like other converts, am now more than ever disposed to preach my former doctrine. Clarke's view I take to be this (if I misinterpret him I trust that the publication of his paper will set me right); that the muscle may, with sufficient accuracy for the present discussion, be said to complete the circle formed, partly by a sensitive nerve and partly again by a motor nerve; that the completeness of this circuit is necessary for the activity of its component parts, and, in particular, let us say, of the muscle; that the disease, involving the posterior roots, destroys their nerve-fibres, not all at once, but separately, or by a few at a time; that in consequence there is at any given moment a rupture of a certain number of such circuits corresponding to the number of ruptured fibres, while at the same time there remains as great a number of unbroken circuits as there are fibres as yet uninvaded; that there are, therefore, muscles still forming parts of complete circuits, and, still therefore, quick, and muscles no longer forming parts of such complete circuits, and therefore deadened; that, as the progress of the disease is irregular, there is consequently, in all movements of the limb, an irregular predominance of the quick muscles over those rendered sluggish or inactive, which change of conditions must end in spasm and disorder. This theory, though fascinating and suggestive, does not, I now think, coincide with the facts; and for these reasons: firstly, the spasmodic disorder of the limbs is not an early symptom, coinciding with the early anæsthesia and with the commencement of the attack upon the posterior fibres in detail, but it is a late symptom and increases with the more and more complete disorganisation of the whole roots. This I have verified by the microscope in one case in which death occurred at the height of the "spasmodic" stage, and it is borne out by the far more valuable investigations of Clarke himself, of Vulpian, and of other observers. Were the wild muscles those whose connections remained entire, their wildness should gradually cease as these connections are cut through; whereas the contrary is nearer the truth, that the muscles become wilder and wilder as the roots become structureless. Secondly, there is no uniformity in the abnormal movements in any given

case at a given time. The disease is often of very slow progress, and, as the destruction of sensitive nerve-fibres slowly advances or remains stationary, so we ought to have mutiny in particular muscles or sets of muscles and uniform deviation from normal movements. But the movements, so far as I can make out, and I have looked into this again in the present aspect of the inquiry, are wholly capricious, and cannot even from one minute to another be seen to form any calculable combinations.

Thirdly, and on the other hand, I consider that the tactile insensibility is amply sufficient to explain not only the maladaptations, but the "spasmodic" movements also; and certainly the spasmodic movements do increase as the tactile æsthesis decreases. If, for example, all the members of a crowd, excited to some common purpose, were to become blind, they would first show indecision in their common movement; and secondly, losing relations among themselves, they would, if still excited, lose all combination with each other. The blinder the men, the blinder the fury. That which keeps muscles or crowds in action as groups is their combination for some common purpose. The common purpose, recognised by the æsthesia, and striven after by impulse, is the nexus, not only between the group and the object, but also between the members of the group themselves. If, while the impulse be continued, the perception of a common purpose be dimmed, the combination will be weakened, and the end unattained. Let it be cut off, and not only will the end be wholly lost sight of, but all combination to that end will cease, and individuals will take the uppermost, according to the momentary and varying states of their tension, their innervation, their size, their lever-ages, and other incidental attributes of men or of muscles. Confusion only can be the result of excitement aroused in a multitude, unless an unifying impression from without can make its way through general or special channels to the centre, be that centre a ganglion, a Napier, or a Beales.

CASES OF SALIVARY CALCULUS:
WITH REMARKS.

By JOSEPH BELL, F.R.C.S.Ed.,
Lecturer on Surgery, Edinburgh.

THE following cases seem to be tolerably characteristic examples of a disease not very common, and regarding which little specific information is given in the more familiar text-books.

CASE I.—A. B., aged 19, called on me in April 1868, in great mental distress. She was a strong, healthy-looking, and very attractive young lady. For two years she had noticed a tumour under her tongue to the right side of the middle line; but, as it seemed stationary and gave her no pain, she had never mentioned its existence to any one; for some months, however, it had been rapidly increasing, and within the last few weeks still more rapidly, and had given her most acute pain. She was now certain that she was suffering from cancer of the tongue.

Present Condition.—On opening the mouth, there was seen on the right side of the tongue a red, fungating, angry-looking tumour as large as a walnut, which forced the tongue upwards so as to interfere with speech, and certainly had a most malignant-looking aspect. It was of stony hardness, and contained no fluid. Trusting to her appearance of perfect health, I excised the projecting portion of the tumour with probe-pointed scissors. Considerable hæmorrhage followed, and no cavity or calculus was exposed as I had hoped. However, by cautiously continuing my incision into the densely fibrous substance of the tumour, I was at last rewarded by extracting an oval calculus, about three-quarters of an inch in length. The relief to the pain was instantaneous, and in a very few days the tumour had entirely disappeared. There has been no return.

CASE II.—Miss F. M., aged 30, consulted me in December 1868. She stated that for the last eight years she had noticed a tumour under the right side of her tongue, which had been frequently the seat of uneasy sensations; at times it had given acute lancinating pain, and latterly, from its size, had interfered with speech and deglutition. All the symptoms had been increasing during the last twelve months. She had consulted at different times two different surgeons, who had both counselled delay. I found that the tumour was of stony hardness, and that a small portion of it projected forwards into the mouth. A free incision enabled me very easily to remove a calculus of peculiar shape, twisted like a ram's-horn, and about an inch and a half in length, and of a bright orange colour. All the symptoms disappeared along with the tumour.

REMARKS.—Salivary calculi are generally found at the orifices of the Whartonian ducts, or in those of the smaller ducts of the sublingual glands. (Craigie's *Elements of General and Pathological Anatomy*, p. 829). They are very rarely found in the duct of the parotid. In most

* Vide *Wien. Med. Wochens.*, 26-28, 1866; quoted *Virch. Jahresh.*, 1867, Band ii, p. 49.

† As my dissent from some points in Dr. Radcliffe's article called forth my own paper, I have seemed to stand in constant opposition to him. I should much regret it, however, if I thus appeared to speak with anything less than the highest respect for Dr. Radcliffe's writings in general, for few writings have given me more pleasure, and I have found few so fruitful.

‡ I gather from an indirect source that this explanation of the phenomena called "locomotor ataxy" has been already given by Professor Leyden, of Königsberg, in *Virchow's Archives*; but, owing to the carelessness of a reference, I have not yet found his paper. What will be the doom of authors who have wasted the time and energy of their fellow-creatures by giving insufficient or inaccurate references?

cases, their presence is easily detected by a probe or even by the finger, as they project close to the extremity of the duct, and an incision over them renders removal very easy. Occasionally, even in this comparatively superficial position, they have been the cause of so much irritation as to be mistaken for cancer. (Holmes's *Dictionary of Surgery*, vol. iii, p. 907). Salivary calculi are occasionally described as either complications or causes of ranula; this seems not to be a correct way of putting the case; for, though some cases are seen in which the salivary calculus has caused by its mechanical obstruction of the duct a temporary collection of saliva behind it, the fluid which is liberated by its removal has not the consistence or appearance of the fluid met with in ordinary cases of ranula. In the two cases recorded above, the calculi were more deeply seated, being rather in the substance of the gland than in the duct, and the second one especially is remarkable for the long duration of the symptoms. In both, the walls of the tumour in which the calculi were imbedded were very firm and hard; in neither was there the slightest accumulation of either saliva or the glairy fluid of a ranula. A case recorded by Gross of Philadelphia (Gross's *Surgery*, 3rd ed., vol. ii, p. 479) seems to have resembled mine in the obscurity of some of the earlier symptoms. Probably, if left alone, the natural course of such calculi is, as South says (South's *Chelius*, vol. ii, p. 406), to escape by ulceration into the cavity of the mouth; but my cases showed that this may, in some cases, involve a very long and a very painful process, before any progress in such ulceration results. Excision in every case is easy, painless, and satisfactory. I have preferred, for the sake of brevity, rather to refer to authorities on this subject than to quote them.

ENCHONDROMA OF THE HAND IN A CHILD: REMOVAL OF THE TUMOUR WITH TWO METACARPAL BONES, THE FINGERS BEING PRESERVED.

By JOHN K. BARTON, M.D., F.R.C.S.I.,
Surgeon to the Adelaide Hospital, Dublin.

SARAH J. C., a year and a half old, was brought to me in January 1868 by her mother, with a note from my friend Dr. Duke, requesting my opinion of the case. I found a tumour occupying the back of the left hand of about the size of a large walnut. It was fixed, widest below, where it passed into the metacarpal bones of the middle and ring fingers, and quite smooth on its surface. It felt very firm and osseous at its base, somewhat softer on its sides, and at its apex soft and semifluctuating. In the palm there was a perceptible swelling corresponding to the two metacarpal bones affected. There seemed to be no pain, even when the tumour was freely handled. It was slightly red at its most prominent part.

The history given by the mother was, that about six months previously, the father of the child, when playing with her, had accidentally struck the back of the child's hand against a hard piece of furniture, which hurt the child a good deal at the time, and soon produced a swelling, which had remained, and gradually increased in hardness and in size; that there did not at any subsequent time appear to be pain in the tumour, as the child played about and used the affected hand freely; but that latterly she could not close her fingers. The child had always enjoyed good health. From the peculiar feel of the tumour, which was yielding at the sides and apex, but firm at its base, and also from the fact that it was certainly continuous with the shafts of the two metacarpal bones, I concluded that it was an enchondroma springing from these bones. However, as the apex was somewhat red and fluctuating, I introduced a fine trocar at this spot into the tumour. A very little blood alone escaped. At a subsequent examination, I found that supuration to a limited extent was taking place where the puncture had been made. After a few days' poulticing, an opening being made, a small quantity of purulent matter escaped; but the bulk of the tumour was not at all diminished. The child was now taken to the country, and given cod-liver oil. When I again had an opportunity of examining it, I found the little patient in excellent health, but the tumour decidedly larger than when it was first seen. Under these circumstances, I advised the removal of the tumour with the bones from which it sprang. The mother and child were consequently admitted to the Adelaide Hospital upon March 10th, 1868. The metacarpal bones of the middle and ring fingers were manifestly part of the tumour; their removal was therefore imperative. The question whether by any possibility the two fingers could be saved was then considered; and, in consultation with my colleagues, I determined to make the attempt, arguing that, even if the fingers proved useless, their presence would lessen the deformity of the hand. Accordingly, upon March 18th, the child being under the

influence of chloroform, I laid bare the tumour, and removed it with the two metacarpal bones to which it was firmly attached. During this process, the extensor and flexor tendons were carefully held out of the way of the knife. The metacarpal bones were cut with the bone-forceps at the carpal extremities, while they were disarticulated from the phalanges. The middle and ring fingers were thus left without any osseous support. The sides of the wound were then drawn together, and the hand and fingers supported with a splint. The little patient did very well, and was able to leave the hospital upon April 13th; the wound in the back of the hand granulating, while the whole hand and fingers were still supported with a splint.

Upon examination of the tumour, it was found to be a true enchondroma, softened in the centre, where it was filled with a substance exactly like sago made with wine, soft, granular, and of a red colour. The walls of the tumour, particularly where it joined the bone, presented the peculiar bluish-white tint of foetal cartilage characteristic of enchondromatous growths. The two metacarpal bones were rough and denuded upon the inner surfaces, where they formed part of the growth, which extended from their palmar, as well as their dorsal, aspects. Microscopic examination of the walls and contents of the tumour showed cartilage-cells in abundance.

Three months elapsed after the child left the hospital before the wound was completely healed; but long before this she was using the hand freely. In November last, the child was brought to me, and I had an opportunity of carefully examining the hand. I found the hand somewhat smaller, not much, however, than the other; the middle and ring fingers somewhat retracted towards the hand, so that their tips were not quite so low as those of the other; but, with this exception, they presented a perfectly natural appearance. With regard to the power of moving, I found that a pencil was grasped and held firmly by all the fingers together; but the two unsupported fingers were not extended again as quickly as the others. Sensation was perfect in them, and they appeared to have grown in proportion with the other fingers.

This case may claim some attention from the rarity of enchondroma in so young a child; but the success which followed the attempt to save the fingers, while their metacarpal bones were necessarily removed, gives it its chief interest. I confess that I felt very doubtful as to either the usefulness or beauty of fingers so left, and only made the attempt to save them, as the mother begged that if possible they might be left. The result more than fulfilled my expectations, and is quite sufficient to encourage a similar attempt in other cases.

CARBOLIC ACID: ITS DOSES AND MEDICINAL VALUE.

By HENRY WILLIAM FULLER, M.D. CANTAB.,
Physician to St. George's Hospital.

FOUR years ago, being desirous of testing the value of carbolic acid as an internal remedy in the treatment of disease, I determined to administer it experimentally in various cases at St. George's Hospital. My first object was to ascertain the maximum dose in which it could be taken without inconvenience, to note the symptoms, if any, produced by full doses, to determine its action on the pulse and the secretions, and to observe whether any evidence could be obtained of its cumulative action. Never having given it internally before, and never having heard of its having been administered in more than two minim-doses, I began by the exhibition of two minims dissolved in a drachm of glycerine and eleven drachms of water. This was given three and ultimately four times in the twenty-four hours. Finding that no appreciable effects resulted from that dose, I added one minim to each dose on alternate days, until the patients were either unable to swallow a stronger dose or complained of unpleasant symptoms. Thus, as far as the mere dose was concerned, I found that some adults—especially men who had been spirit-drinkers—could take ten or twelve minims without inconvenience and notwithstanding the occurrence of a certain degree of discomfort, could take doses of fifteen minims three or four times a day for many days consecutively; but that most persons, especially women, began to complain when the dose had been increased to eight or ten minims, and found six or seven minims a full dose.

The disagreeable symptoms produced by full or overfull doses were, firstly, a sense of burning in the throat on swallowing the draught; and, secondly, a giddiness and fulness, or peculiar feeling in the head—a feeling which occurred in some persons within two minutes after taking the acid, and in others not until the expiration of six or eight minutes. In some persons this giddiness passed off in ten or fifteen minutes, and in others lasted nearly an hour. In no instance was there